



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶: C12N 15/90, 15/85	A2	(11) International Publication Number: WO 97/40183 (43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/US97/05911 (22) International Filing Date: 10 April 1997 (10.04.97) (30) Priority Data: 629,822 10 April 1996 (10.04.96) US 682,080 15 July 1996 (15.07.96) US 695,191 7 August 1996 (07.08.96) US (71) Applicants (for all designated States except US): THE BIOLOGICAL RESEARCH CENTER OF THE HUNGARIAN ACADEMY OF SCIENCES [HU/HU]; P.O. Box 521, H-6701 Szeged (HU). LOMA LINDA UNIVERSITY [US/US]; Loma Linda, CA 92350 (US). AMERICAN GENE THERAPY, INC. [CA/CA]; 5022 154th Street, Edmonton, Alberta T6H 5P6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): HADLACZKY, Gyula [HU/HU]; Szamos U.I.A. IX. 36., H-6723 Szeged (HU). SZALAY, Aladar, A. [US/US]; 7327 Fairwood, Highland, CA 92346 (US). (74) Agent: SEIDMAN, Stephanie, L.; Brown, Martin, Haller & McClain, 1660 Union Street, San Diego, CA 92101-2926 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU. ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>Without international search report and to be republished upon receipt of that report.</i>

(54) Title: ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS FOR PREPARING ARTIFICIAL CHROMOSOMES

(57) Abstract

Methods for preparing cell lines that contain artificial chromosomes, methods for preparation of artificial chromosomes, methods for purification of artificial chromosomes, methods for targeted insertion of heterologous DNA into artificial chromosomes, and methods for delivery of the chromosomes to selected cells and tissues are provided. Also provided are cell lines for use in the methods, and cell lines and chromosomes produced by the methods. In particular, satellite artificial chromosomes that, except for inserted heterologous DNA, are substantially composed of heterochromatin, are provided. Methods for use of the artificial chromosomes, including for gene therapy, production of gene products and production of transgenic plants and animals are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

-1-

**ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS FOR
PREPARING ARTIFICIAL CHROMOSOMES**

RELATED APPLICATIONS

For U.S. national stage purposes, this application is a continuation-
5 in-part of copending U.S. application Serial No. 08/695,191, filed August
7, 1996 by GYULA HADLACZKY and ALADAR SZALAY, entitled
*ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS FOR
PREPARING ARTIFICIAL CHROMOSOMES*. This application is also
continuation-in-part of copending U.S. application Serial No. 08/682,080,
10 filed July 15, 1996 by GYULA HADLACZKY and ALADAR SZALAY,
entitled *ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS
FOR PREPARING ARTIFICIAL CHROMOSOMES*, and is also a
continuation-in-part of copending U.S. application Serial No. 08/629,822,
filed April 10, 1996 by GYULA HADLACZKY and ALADAR SZALAY,
15 entitled *ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS
FOR PREPARING ARTIFICIAL CHROMOSOMES*.

For international purposes, the benefit of priority to each of these
application is claimed and the subject matter of that application is
incorporated herein in its entirety.

20 U.S. application Serial No. 08/695,191 is a continuation-in-part of
U.S. application Serial No. 08/682,080 and also is a continuation-in-part
of U.S. application Serial No. 08/629,822. U.S. application Serial No.
08/682,080 is a continuation-in-part of U.S. application Serial No.
08/629,822.

25 This application is related to U.S. application Serial No.
07/759,558, now U.S. Patent No. 5,288,625, is related to U.S.
application Serial No. 08/734,344, filed October 21, 1996, and is related
to allowed U.S. application Serial No. 08/375,271, filed 1/19/95, which
is a continuation of U.S. application Serial No. 08/080,097, filed 6/23/93

-2-

which is a continuation of U.S. application Serial No. 07/892,487, filed 6/3/92, which is a continuation of U.S. application Serial No. 07/521,073, filed 5/9/90.

To the extent permitted, the subject matter of each of U.S. application Serial Nos. 08/734,344, 08/695,191, 08/682,080, 08/629,822, 08/375,271, 08/080,097, 07/892,487, and 07/521,073, and U.S. Patent No. 5,288,625 is incorporated in its entirety by reference thereto.

FIELD OF THE INVENTION

10 The present invention relates to methods for preparing cell lines that contain artificial chromosomes, methods for isolation of the artificial chromosomes, targeted insertion of heterologous DNA into the chromosomes, delivery of the chromosomes to selected cells and tissues and methods for isolation and large-scale production of the
15 chromosomes. Also provided are cell lines for use in the methods, and cell lines and chromosomes produced by the methods. Further provided are cell-based methods for production of heterologous proteins, gene therapy methods and methods of generating transgenic animals, particularly non-human transgenic animals, that use artificial
20 chromosomes.

BACKGROUND OF THE INVENTION

Several viral vectors, non-viral, and physical delivery systems for gene therapy and recombinant expression of heterologous nucleic acids have been developed [see, e.g., Mitani et al. (1993) Trends Biotech.
25 11:162-166]. The presently available systems, however, have numerous limitations, particularly where persistent, stable, or controlled gene expression is required. These limitations include: (1) size limitations because there is a limit, generally on order of about ten kilobases [kB], at most, to the size of the DNA insert [gene] that can be accepted by viral

-3-

vectors, whereas a number of mammalian genes of possible therapeutic importance are well above this limit, especially if all control elements are included; (2) the inability to specifically target integration so that random integration occurs which carries a risk of disrupting vital genes or cancer suppressor genes; (3) the expression of randomly integrated therapeutic genes may be affected by the functional compartmentalization in the nucleus and are affected by chromatin-based position effects; (4) the copy number and consequently the expression of a given gene to be integrated into the genome cannot be controlled. Thus, improvements in gene delivery and stable expression systems are needed [see, e.g., Mulligan (1993) Science 260:926-932].

In addition, safe and effective vectors and gene therapy methods should have numerous features that are not assured by the presently available systems. For example, a safe vector should not contain DNA elements that can promote unwanted changes by recombination or mutation in the host genetic material, should not have the potential to initiate deleterious effects in cells, tissues, or organisms carrying the vector, and should not interfere with genomic functions. In addition, it would be advantageous for the vector to be non-integrative, or designed for site-specific integration. Also, the copy number of therapeutic gene(s) carried by the vector should be controlled and stable, the vector should secure the independent and controlled function of the introduced gene(s); and the vector should accept large (up to Mb size) inserts and ensure the functional stability of the insert.

The limitations of existing gene delivery technologies, however, argue for the development of alternative vector systems suitable for transferring large [up to Mb size or larger] genes and gene complexes together with regulatory elements that will provide a safe, controlled, and persistent expression of the therapeutic genetic material.

-4-

At the present time, none of the available vectors fulfill all these requirements. Most of these characteristics, however, are possessed by chromosomes. Thus, an artificial chromosome would be an ideal vector for gene therapy, as well as for stable, high-level, controlled production of gene products that require coordination of expression of numerous genes or that are encoded by large genes, and other uses. Artificial chromosomes for expression of heterologous genes in yeast are available, but construction of defined mammalian artificial chromosomes has not been achieved. Such construction has been hindered by the lack of an isolated, functional, mammalian centromere and uncertainty regarding the requisites for its production and stable replication. Unlike in yeast, there are no selectable genes in close proximity to a mammalian centromere, and the presence of long runs of highly repetitive pericentric heterochromatic DNA makes the isolation of a mammalian centromere using presently available methods, such as chromosome walking, virtually impossible. Other strategies are required for production of mammalian artificial chromosomes, and some have been developed. For example, U.S. Patent No. 5,288,625 provides a cell line that contains an artificial chromosome, a minichromosome, that is about 20 to 30 megabases. Methods provided for isolation of these chromosomes, however, provide preparations of only about 10-20% purity. Thus, development of alternative artificial chromosomes and perfection of isolation and purification methods as well as development of more versatile chromosomes and further characterization of the minichromosomes is required to realize the potential of this technology.

Therefore, it is an object herein to provide mammalian artificial chromosomes and methods for introduction of foreign DNA into such chromosomes. It is also an object herein to provide methods of isolation and purification of the chromosomes. It is also an object herein to

-5-

provide methods for introduction of the mammalian artificial chromosome into selected cells, and to provide the resulting cells, as well as transgenic non-human animals, birds, fish and plants that contain the artificial chromosomes. It is also an object herein to provide methods for

5 gene therapy and expression of gene products using artificial chromosomes. It is a further object herein to provide methods for constructing species-specific artificial chromosomes de novo. Another object herein is to provide methods to generate de novo mammalian artificial chromosomes.

10 SUMMARY OF THE INVENTION

Mammalian artificial chromosomes [MACs] are provided. Also provided are artificial chromosomes for other higher eukaryotic species, such as insects, birds, fowl and fish, produced using the MACS and methods provided herein. Methods for generating and isolating such

15 chromosomes are provided. Methods using the MACs to construct artificial chromosomes from other species, such as insect, bird, fowl and fish species are also provided. The artificial chromosomes are fully functional stable chromosomes. Two types of artificial chromosomes are provided. One type, herein referred to as SATACs [satellite artificial

20 chromosomes or satellite DNA based artificial chromosomes (the terms are used interchangeably herein)] are stable heterochromatic chromosomes, and the other type are minichromosomes based on amplification of euchromatin.

Artificial chromosomes provide an extra-genomic locus for targeted

25 integration of megabase [Mb] pair size DNA fragments that contain single or multiple genes, including multiple copies of a single gene operatively linked to one promoter or each copy or several copies linked to separate promoters. Thus, methods using the MACs to introduce the genes into cells, tissues, and animals, as well as species such as birds, fowl, fish

-6-

and plants, are also provided. The artificial chromosomes with integrated heterologous DNA may be used in methods of gene therapy, in methods of production of gene products, particularly products that require expression of multigenic biosynthetic pathways, and also are intended for
5 delivery into the nuclei of germline cells, such as embryo-derived stem cells [ES cells], for production of transgenic (non-human) animals, birds, fowl and fish. Transgenic plants, including monocots and dicots, are also contemplated herein.

Mammalian artificial chromosomes provide extra-genomic specific
10 integration sites for introduction of genes encoding proteins of interest and permit megabase size DNA integration so that, for example, genes encoding an entire metabolic pathway or a very large gene, such as the cystic fibrosis [CF; ~250 kb] genomic DNA gene, several genes, such as multiple genes encoding a series of antigens for preparation of a
15 multivalent vaccine, can be stably introduced into a cell. Vectors for targeted introduction of such genes, including the tumor suppressor genes, such as p53, the cystic fibrosis transmembrane regulator cDNA [CFTR], and the genes for anti-HIV ribozymes, such as an anti-HIV gag ribozyme gene, into the artificial chromosomes are also provided.

20 The chromosomes provided herein are generated by introducing heterologous DNA that includes DNA encoding one or multiple selectable marker(s) into cells, preferably a stable cell line, growing the cells under selective conditions, and identifying from among the resulting clones those that include chromosomes with more than one centromere and/or
25 fragments thereof. The amplification that produces the additional centromere or centromeres occurs in cells that contain chromosomes in which the heterologous DNA has integrated near the centromere in the pericentric region of the chromosome. The selected clonal cells are then used to generate artificial chromosomes.

-7-

Although non-targeted introduction of DNA, which results in some frequency of integration into appropriate loci, targeted introduction is preferred. Hence, in preferred embodiments, the DNA with the selectable marker that is introduced into cells to initiate generation of
5 artificial chromosomes includes sequences that target it to the an amplifiable region, such as the pericentric region, heterochromatin, and particularly rDNA of the chromosome. For example, vectors, such as pTEMPUD and pHASPUd [provided herein], which include such DNA specific for mouse satellite DNA and human satellite DNA, respectively,
10 are provided. The plasmid pHASPUd is a derivative of pTEMPUD that contains human satellite DNA sequences that specifically target human chromosomes. Preferred targeting sequences include mammalian ribosomal RNA (rRNA) gene sequences (referred to herein as rDNA) which target the heterologous DNA to integrate into the rDNA region of
15 those chromosomes that contain rDNA. For example, vectors, such as pTERPUD, which include mouse rDNA, are provided. Upon integration into existing chromosomes in the cells, these vectors can induce the amplification that results in generation of additional centromeres.

Artificial chromosomes are generated by culturing the cells with
20 the multicentric, typically dicentric, chromosomes under conditions whereby the chromosome breaks to form a minichromosome and formerly dicentric chromosome. Among the MACs provided herein are the SATACs, which are primarily made up of repeating units of short satellite DNA and are nearly fully heterochromatic, so that without
25 insertion of heterologous or foreign DNA, the chromosomes preferably contain no genetic information or contain only non-protein-encoding gene sequences such as rDNA sequences. They can thus be used as "safe" vectors for delivery of DNA to mammalian hosts because they do not contain any potentially harmful genes. The SATACs are generated, not

-8-

from the minichromosome fragment as, for example, in U.S. Patent No. 5,288,625, but from the fragment of the formerly dicentric chromosome.

In addition, methods for generating euchromatic minichromosomes and the use thereof are also provided herein. Methods for generating one type of MAC, the minichromosome, previously described in U.S. Patent No. 5,288,625, and the use thereof for expression of heterologous DNA are provided. In a particular method provided herein for generating a MAC, such as a minichromosome, heterologous DNA that includes mammalian rDNA and one or more selectable marker genes is introduced into cells which are then grown under selective conditions. Resulting cells that contain chromosomes with more than one centromere are selected and cultured under conditions whereby the chromosome breaks to form a minichromosome and a formerly multicentric (typically dicentric) chromosome from which the minichromosome was released.

Cell lines containing the minichromosome and the use thereof for cell fusion are also provided. In one embodiment, a cell line containing the mammalian minichromosome is used as recipient cells for donor DNA encoding a selected gene or multiple genes. To facilitate integration of the donor DNA into the minichromosome, the recipient cell line preferably contains the minichromosome but does not also contain the formerly dicentric chromosome. This may be accomplished by methods disclosed herein such as cell fusion and selection of cells that contain a minichromosome and no formerly dicentric chromosome. The donor DNA is linked to a second selectable marker and is targeted to and integrated into the minichromosome. The resulting chromosome is transferred by cell fusion into an appropriate recipient cell line, such as a Chinese hamster cell line [CHO]. After large-scale production of the cells carrying the engineered chromosome, the chromosome is isolated. In particular, metaphase chromosomes are obtained, such as by addition of colchicine,

-9-

and they are purified from the cell lysate. These chromosomes are used for cloning, sequencing and for delivery of heterologous DNA into cells.

- Also provided are SATACs of various sizes that are formed by repeated culturing under selective conditions and subcloning of cells that
- 5 contain chromosomes produced from the formerly dicentric chromosomes. The exemplified SATACs are based on repeating DNA units that are about 15 Mb [two ~ 7.5 Mb blocks]. The repeating DNA unit of SATACs formed from other species and other chromosomes may vary, but typically would be on the order of about 7 to about 20 Mb.
- 10 The repeating DNA units are referred to herein as megareplicons, which in the exemplified SATACs contain tandem blocks of satellite DNA flanked by non-satellite DNA, including heterologous DNA and non-satellite DNA. Amplification produces an array of chromosome segments [each called an amplicon] that contain two inverted megareplicons
- 15 bordered by heterologous ["foreign"] DNA. Repeated cell fusion, growth on selective medium and/or BrdU [5-bromodeoxyuridine] treatment or other treatment with other genome destabilizing reagent or agent, such as ionizing radiation, including X-rays, and subcloning results in cell lines that carry stable heterochromatic or partially heterochromatic
- 20 chromosomes, including a 150-200 Mb "sausage" chromosome, a 500-1000 Mb gigachromosome, a stable 250-400 Mb megachromosome and various smaller stable chromosomes derived therefrom. These chromosomes are based on these repeating units and can include heterologous DNA that is expressed.
- 25 Thus, methods for producing MACs of both types (i.e., SATACS and minichromosomes) are provided. These methods are applicable to the production of artificial chromosomes containing centromeres derived from any higher eukaryotic cell, including mammals, birds, fowl, fish, insects and plants.

-10-

The resulting chromosomes can be purified by methods provided herein to provide vectors for introduction of heterologous DNA into selected cells for production of the gene product(s) encoded by the heterologous DNA, for production of transgenic (non-human) animals, birds, fowl, fish and plants or for gene therapy.

In addition, methods and vectors for fragmenting the minichromosomes and SATACs are provided. Such methods and vectors can be used for in vivo generation of smaller stable artificial chromosomes. Vectors for chromosome fragmentation are used to produce an artificial chromosome that contains a megareplicon, a centromere and two telomeres and will be between about 7.5 Mb and about 60 Mb, preferably between about 10 Mb-15 Mb and 30-50 Mb. As exemplified herein, the preferred range is between about 7.5 Mb and 50 Mb. Such artificial chromosomes may also be produced by other methods.

Isolation of the 15 Mb [or 30 Mb amplicon containing two 15 Mb inverted repeats] or a 30 Mb or higher multimer, such as 60 Mb, thereof should provide a stable chromosomal vector that can be manipulated in vitro. Methods for reducing the size of the MACs to generate smaller stable self-replicating artificial chromosomes are also provided.

Also provided herein, are methods for producing mammalian artificial chromosomes, including those provided herein, in vitro, and the resulting chromosomes. The methods involve *in vitro* assembly of the structural and functional elements to provide a stable artificial chromosome. Such elements include a centromere, two telomeres, at least one origin of replication and filler heterochromatin, e.g., satellite DNA. A selectable marker for subsequent selection is also generally included. These specific DNA elements may be obtained from the artificial chromosomes provided herein such as those that have been

-11-

- generated by the introduction of heterologous DNA into cells and the subsequent amplification that leads to the artificial chromosome, particularly the SATACs. Centromere sequences for use in the *in vitro* construction of artificial chromosomes may also be obtained by
- 5 employing the centromere cloning methods provided herein. In preferred embodiments, the sequences providing the origin of replication, in particular, the megareplicator, are derived from rDNA. These sequences preferably include the rDNA origin of replication and amplification promoting sequences.
- 10 Methods and vectors for targeting heterologous DNA into the artificial chromosomes are also provided as are methods and vectors for fragmenting the chromosomes to produce smaller but stable and self-replicating artificial chromosomes.
- The chromosomes are introduced into cells to produce stable
- 15 transformed cell lines or cells, depending upon the source of the cells. Introduction is effected by any suitable method including, but not limited to electroporation, direct uptake, such as by calcium phosphate precipitation, uptake of isolated chromosomes by lipofection, by microcell fusion, by lipid-mediated carrier systems or other suitable method. The
- 20 resulting cells can be used for production of proteins in the cells. The chromosomes can be isolated and used for gene delivery. Methods for isolation of the chromosomes based on the DNA content of the chromosomes, which differs in MACs versus the authentic chromosomes, are provided. Also provided are methods that rely on
- 25 content, particularly density, and size of the MACs.

-12-

These artificial chromosomes can be used in gene therapy, gene product production systems, production of humanized genetically transformed animal organs, production of transgenic plants and animals (non-human), including mammals, birds, fowl, fish, invertebrates, 5 vertebrates, reptiles and insects, any organism or device that would employ chromosomal elements as information storage vehicles, and also for analysis and study of centromere function, for the production of artificial chromosome vectors that can be constructed in vitro, and for the preparation of species-specific artificial chromosomes. The artificial 10 chromosomes can be introduced into cells using microinjection, cell fusion, microcell fusion, electroporation, nuclear transfer, electrofusion, projectile bombardment, nuclear transfer, calcium phosphate precipitation, lipid-mediated transfer systems and other such methods. Cells particularly suited for use with the artificial chromosomes include, 15 but are not limited to plant cells, particularly tomato, arabidopsis, and others, insect cells, including silk worm cells, insect larvae, fish, reptiles, amphibians, arachnids, mammalian cells, avian cells, embryonic stem cells, haematopoietic stem cells, embryos and cells for use in methods of genetic therapy, such as lymphocytes that are used in methods of adop- 20 tive immunotherapy and nerve or neural cells. Thus methods of producing gene products and transgenic (non-human) animals and plants are provided. Also provided are the resulting transgenic animals and plants.

Exemplary cell lines that contain these chromosomes are also provided.

25 Methods for preparing artificial chromosomes for particular species and for cloning centromeres are also provided. For example, two exemplary methods provided for generating artificial chromosomes for use in different species are as follows. First, the methods herein may be applied to different species. Second, means for generating species-

-13-

specific artificial chromosomes and for cloning centromeres are provided. In particular, a method for cloning a centromere from an animal or plant is provided by preparing a library of DNA fragments that contain the genome of the plant or animal and introducing each of the fragments
5 into a mammalian satellite artificial chromosome [SATAC] that contains a centromere from a species, generally a mammal, different from the selected plant or animal, generally a non-mammal, and a selectable marker. The selected plant or animal is one in which the mammalian species centromere does not function. Each of the SATACs is
10 introduced into the cells, which are grown under selective conditions, and cells with SATACs are identified. Such SATACS should contain a centromere encoded by the DNA from the library or should contain the necessary elements for stable replication in the selected species.

Also provided are libraries in which the relatively large fragments
15 of DNA are contained on artificial chromosomes.

Transgenic (non-human) animals, invertebrates and vertebrates, plants and insects, fish, reptiles, amphibians, arachnids, birds, fowl, and mammals are also provided. Of particular interest are transgenic (non-human) animals and plants that express genes that confer resistance or
20 reduce susceptibility to disease. For example, the transgene may encode a protein that is toxic to a pathogen, such as a virus, bacterium or pest, but that is not toxic to the transgenic host. Furthermore, since multiple genes can be introduced on a MAC, a series of genes encoding an antigen can be introduced, which upon expression will serve to
25 immunize [in a manner similar to a multivalent vaccine] the host animal against the diseases for which exposure to the antigens provide immunity or some protection.

-14-

Also of interest are transgenic (non-human) animals that serve as models of certain diseases and disorders for use in studying the disease and developing therapeutic treatments and cures thereof. Such animal models of disease express genes [typically carrying a disease-associated
5 mutation], which are introduced into the animal on a MAC and which induce the disease or disorder in the animal. Similarly, MACs carrying genes encoding antisense RNA may be introduced into animal cells to generate conditional "knock-out" transgenic (non-human) animals. In such animals, expression of the antisense RNA results in decreased or
10 complete elimination of the products of genes corresponding to the antisense RNA. Of further interest are transgenic mammals that harbor MAC-carried genes encoding therapeutic proteins that are expressed in the animal's milk. Transgenic (non-human) animals for use in xenotransplantation, which express MAC-carried genes that serve to
15 humanize the animal's organs, are also of interest. Genes that might be used in humanizing animal organs include those encoding human surface antigens.

Methods for cloning centromeres, such as mammalian centromeres, are also provided. In particular, in one embodiment, a
20 library composed of fragments of SATACs are cloned into YACs [yeast artificial chromosomes] that include a detectable marker, such as DNA encoding tyrosinase, and then introduced into mammalian cells, such as albino mouse embryos. Mice produced from embryos containing such YACs that include a centromere that functions in mammals will express
25 the detectable marker. Thus, if mice are produced from albino mouse embryos into which a functional mammalian centromere was introduced, the mice will be pigmented or have regions of pigmentation.

-15-

A method for producing repeated tandem arrays of DNA is provided. This method, exemplified herein using telomeric DNA, is applicable to any repeat sequence, and in particular, low complexity repeats. The method provided herein for synthesis of arrays of tandem

5 DNA repeats are based in a series of extension steps in which successive doublings of a sequence of repeats results in an exponential expansion of the array of tandem repeats. An embodiment of the method of synthesizing DNA fragments containing tandem repeats may generally be described as follows. Two oligonucleotides are used as starting

10 materials. Oligonucleotide 1 is of length k of repeated sequence (the flanks of which are not relevant) and contains a relatively short stretch (60-90 nucleotides) of the repeated sequence, flanked with appropriately chosen restriction sites:-

5'-S1 > S2__-3'

15 where S1 is restriction site 1 cleaved by E1, S2 is a second restriction site cleaved by E2 > represents a simple repeat unit, and '__' denotes a short (8-10) nucleotide flanking sequence complementary to oligonucleotide 2:

3'-_____S3-5'

20 where S3 is a third restriction site for enzyme E3 and which is present in the vector to be used during the construction. The method involves the following steps: (1) oligonucleotides 1 and 2 are annealed; (2) the annealed oligonucleotides are filled-in to produce a double-stranded (ds) sequence; (3) the double-stranded DNA is cleaved with restriction

25 enzymes E1 and E3 and subsequently ligated into a vector (e.g., pUC19 or a yeast vector) that has been cleaved with the same enzymes E1 and E3; (4) the insert is isolated from a first portion of the plasmid by digesting with restriction enzymes E1 and E3, and a second portion of the plasmid is cut with enzymes E2 (treated to remove the 3'-overhang)

-16-

- and E3, and the large fragment (plasmid DNA plus the insert) is isolated; (5) the two DNA fragments (the S1-S3 insert fragment and the vector plus insert) are ligated; and (6) steps 4 and 5 are repeated as many times as needed to achieve the desired repeat sequence size. In each
- 5 extension cycle, the repeat sequence size doubles, i.e., if m is the number of extension cycles, the size of the repeat sequence will be $k \times 2^m$ nucleotides.

DESCRIPTION OF THE DRAWINGS

- Figure 1 is a schematic drawing depicting formation of the
- 10 MMCneo [the minichromosome] chromosome. A-G represents the successive events consistent with observed data that would lead to th formation and stabilization of the minichromosome.

- Figure 2 shows a schematic summary of the manner in which the observed new chromosomes would form, and the relationships among
- 15 the different *de novo* formed chromosomes. In particular, this figure shows a schematic drawing of the *de novo* chromosome formation initiated in the centromeric region of mouse chromosome 7. (A) A single E-type amplification in the centromeric region of chromosome 7 generates a neo-centromere linked to the integrated "foreign" DNA, and
- 20 forms a dicentric chromosome. Multiple E-type amplification forms the λ neo-chromosome, which separates from the remainder of mouse chromosome 7 through a specific breakage between the centromeres of the dicentric chromosome and which was stabilized in a mouse-hamster hybrid cell line; (B) Specific breakage between the centromeres of a
- 25 dicentric chromosome 7 generates a chromosome fragment with the neo-centromere, and a chromosome 7 with traces of heterologous DNA at the end; (C) Inverted duplication of the fragment bearing the neo-centrom re results in the formation of a stable neo-minichromosome; (D) Integration of exogenous DNA into the heterologous DNA region of the formerly

-17-

dicentric chromosome 7 initiates H-type amplification, and the formation of a heterochromatic arm. By capturing a euchromatic terminal segment, this new chromosome arm is stabilized in the form of the "sausage" chromosome; (E) BrdU [5-bromodeoxyuridine] treatment and/or drug
5 selection induce further H-type amplification, which results in the formation of an unstable gigachromosome: (F) Repeated BrdU treatments and/or drug selection induce further H-type amplification including a centromere duplication, which leads to the formation of another heterochromatic chromosome arm. It is split off from the chromosome 7
10 by chromosome breakage, and by acquiring a terminal segment, the stable megachromosome is formed.

Figure 3 is a schematic diagram of the replicon structure and a scheme by which a megachromosome could be produced.

Figure 4 sets forth the relationships among some of the exemplary
15 cell lines described herein.

Figure 5 is a diagram of the plasmid pTEMPUD.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

Definitions

Unless defined otherwise, all technical and scientific terms used
20 herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. All patents and publications referred to herein are incorporated by reference.

As used herein, a mammalian artificial chromosome (MAC) is a piece of DNA that can stably replicate and segregate alongside
25 endogenous chromosomes. It has the capacity to accommodate and express heterologous genes inserted therein. It is referred to as a mammalian artificial chromosome because it includes an active mammalian centromere(s). Plant artificial chromosomes, insect artificial chromosomes and avian artificial chromosomes refer to chromosomes

-18-

that include plant and insect centromeres, respectively. A human artificial chromosome [HAC] refers to chromosomes that include human centromeres, BUGACs refer to insect artificial chromosomes, and AVACs refer to avian artificial chromosomes. Among the MACs provided herein
5 are SATACs, minichromosomes, and in vitro synthesized artificial chromosomes. Methods for construction of each type are provided herein.

As used herein, in vitro synthesized artificial chromosomes are artificial chromosomes that is produced by joining the essential
10 components (at least the centromere, and origins of replication) in vitro.

As used herein, endogenous chromosomes refer to genomic chromosomes as found in the cell prior to generation or introduction of a MAC.

As used herein, stable maintenance of chromosomes occurs when
15 at least about 85%, preferably 90%, more preferably 95%, of the cells retain the chromosome. Stability is measured in the presence of a selective agent. Preferably these chromosomes are also maintained in the absence of a selective agent. Stable chromosomes also retain their structure during cell culturing, suffering neither intrachromosomal nor
20 interchromosomal rearrangements.

As used herein, growth under selective conditions means growth of a cell under conditions that require expression of a selectable marker for survival.

As used herein, an agent that destabilizes a chromosome is any
25 agent known by those of skill in the art to enhance amplification events, mutations. Such agents, which include BrdU, are well known to those of skill in the art.

-19-

As used herein, *de novo* with reference to a centromere, refers to generation of an excess centromere as a result of incorporation of a heterologous DNA fragment using the methods herein.

As used herein, euchromatin and heterochromatin have their
5 recognized meanings, euchromatin refers to chromatin that stains
diffusely and that typically contains genes, and heterochromatin refers to
chromatin that remains unusually condensed and that has been thought
to be transcriptionally inactive. Highly repetitive DNA sequences
[satellite DNA], at least with respect to mammalian cells, are usually
10 located in regions of the heterochromatin surrounding the centromere
[pericentric heterochromatin]. Constitutive heterochromatin refers to
heterochromatin that contains the highly repetitive DNA which is
constitutively condensed and genetically inactive.

As used herein, BrdU refers to 5-bromodeoxyuridine, which during
15 replication is inserted in place of thymidine. BrdU is used as a mutagen; it
also inhibits condensation of metaphase chromosomes during cell
division.

As used herein, a dicentric chromosome is a chromosome that
contains two centromeres. A multicentric chromosome contains more
20 than two centromeres.

As used herein, a formerly dicentric chromosome is a chromosome
that is produced when a dicentric chromosome fragments and acquir s
new telomeres so that two chromosomes, each having one of the
centromeres, are produced. Each of the fragments are replicable
25 chromosomes. If one of the chromosomes undergoes amplification of
euchromatic DNA to produce a fully functional chromosome that contains
the newly introduced heterologous DNA and primarily [at least more than
50%] euchromatin, it is a minichromosome. The remaining chromosome
is a formerly dicentric chromosome. If one of the chromosomes

-20-

undergoes amplification, whereby heterochromatin [satellite DNA] is amplified and a euchromatic portion [or arm] remains, it is referred to as a sausage chromosome. A chromosome that is substantially all heterochromatin, except for portions of heterologous DNA, is called a

5 SATAC. Such chromosomes [SATACs] can be produced from sausage chromosomes by culturing the cell containing the sausage chromosome under conditions, such as BrdU treatment and/or growth under selective conditions, that destabilize the chromosome so that a satellite artificial chromosomes [SATAC] is produced. For purposes herein, it is

10 understood that SATACs may not necessarily be produced in multiple steps, but may appear after the initial introduction of the heterologous DNA and growth under selective conditions, or they may appear after several cycles of growth under selective conditions and BrdU treatment.

As used herein, a SATAC refers to a chromosome that is

15 substantially all heterochromatin, except for portions of heterologous DNA. Typically, SATACs are satellite DNA based artificial chromosomes, but the term encompasses any chromosome made by the methods herein that contains more heterochromatin than euchromatin.

As used herein, amplifiable, when used in reference to a

20 chromosome, particularly the method of generating SATACs provided herein, refers to a region of a chromosome that is prone to amplification. Amplification typically occurs during replication and other cellular events involving recombination. Such regions are typically regions of the chromosome that include tandem repeats, such as satellite DNA, rDNA

25 and other such sequences.

As used herein, amplification, with reference to DNA, is a process in which segments of DNA are duplicated to yield two or multiple copies of identical or nearly identical DNA segments that are typically joined as substantially tandem or successive repeats or inverted repeats.

-21-

As used herein an amplicon is a repeated DNA amplification unit that contains a set of inverted repeats of the megareplicon. A megareplicon represents a higher order replication unit. For example, with reference to the SATACs, the megareplicon contains a set of

5 tandem DNA blocks each containing satellite DNA flanked by non-satellite DNA. Contained within the megareplicon is a primary replication site, referred to as the megareplicator, which may be involved in organizing and facilitating replication of the pericentric heterochromatin and possibly the centromeres. Within the megareplicon there may be

10 smaller [e.g., 50-300 kb in some mammalian cells] secondary replicons. In the exemplified SATACS, the megareplicon is defined by two tandem ~ 7.5 Mb DNA blocks [see, e.g., Fig. 3]. Within each artificial chromosome [AC] or among a population thereof, each amplicon has the same gross structure but may contain sequence variations. Such

15 variations will arise as a result of movement of mobile genetic elements, deletions or insertions or mutations that arise, particularly in culture. Such variation does not affect the use of the ACs or their overall structure as described herein.

As used herein, ribosomal RNA [rRNA] is the specialized RNA that

20 forms part of the structure of a ribosome and participates in the synthesis of proteins. Ribosomal RNA is produced by transcription of genes which, in eukaryotic cells, are present in multiple copies. In human cells, the approximately 250 copies of rRNA genes per haploid genome are spread out in clusters on at least five different chromosomes

25 (chromosomes 13, 14, 15, 21 and 22). In mouse cells, the presence of ribosomal DNA [rDNA] has been verified on at least 11 pairs out of 20 mouse chromosomes [chromosomes 5, 6, 9, 11, 12, 15, 16, 17, 18, 19 and X][see e.g., Rowe et al. (1996) Mamm. Genome 7:886-889 and Johnson et al. (1993) Mamm. Genome 4:49-52]. In eukaryotic cells, the

-22-

multiple copies of the highly conserved rRNA genes are located in a tandemly arranged series of rDNA units, which are generally about 40-45 kb in length and contain a transcribed region and a nontranscribed region known as spacer (i.e., intergenic spacer) DNA which can vary in length and sequence. In the human and mouse, these tandem arrays of rDNA units are located adjacent to the pericentric satellite DNA sequences (heterochromatin). The regions of these chromosomes in which the rDNA is located are referred to as nucleolar organizing regions (NOR) which loop into the nucleolus, the site of ribosome production within the cell nucleus.

As used herein, the minichromosome refers to a chromosome derived from a multicentric, typically dicentric, chromosome [see, e.g., FIG. 1] that contains more euchromatic than heterochromatic DNA.

As used herein, a megachromosome refers to a chromosome that, except for introduced heterologous DNA, is substantially composed of heterochromatin. Megachromosomes are made of an array of repeated amplicons that contain two inverted megareplicons bordered by introduced heterologous DNA [see, e.g., Figure 3 for a schematic drawing of a megachromosome]. For purposes herein, a megachromosome is about 50 to 400 Mb, generally about 250-400 Mb. Shorter variants are also referred to as truncated megachromosomes [about 90 to 120 or 150 Mb], dwarf megachromosomes [~ 150-200 Mb] and cell lines, and a micro-megachromosome [~ 50-90 Mb, typically 50-60 Mb]. For purposes herein, the term megachromosome refers to the overall repeated structure based on an array of repeated chromosomal segments [amplicons] that contain two inverted megareplicons bordered by any inserted heterologous DNA. The size will be specified.

-23-

As used herein, genetic therapy involves the transfer or insertion of heterologous DNA into certain cells, target cells, to produce specific gene products that are involved in correcting or modulating disease. The DNA is introduced into the selected target cells in a manner such that the

5 heterologous DNA is expressed and a product encoded thereby is produced. Alternatively, the heterologous DNA may in some manner mediate expression of DNA that encodes the therapeutic product. It may encode a product, such as a peptide or RNA, that in some manner mediates, directly or indirectly, expression of a therapeutic product.

10 Genetic therapy may also be used to introduce therapeutic compounds, such as TNF, that are not normally produced in the host or that are not produced in therapeutically effective amounts or at a therapeutically useful time. Expression of the heterologous DNA by the target cells within an organism afflicted with the disease thereby enables modulation

15 of the disease. The heterologous DNA encoding the therapeutic product may be modified prior to introduction into the cells of the afflicted host in order to enhance or otherwise alter the product or expression thereof.

As used herein, heterologous or foreign DNA and RNA are used interchangeably and refer to DNA or RNA that does not occur naturally

20 as part of the genome in which it is present or which is found in a location or locations in the genome that differ from that in which it occurs in nature. It is DNA or RNA that is not endogenous to the cell and has been exogenously introduced into the cell. Examples of heterologous DNA include, but are not limited to, DNA that encodes a

25 gene product or gene product(s) of interest, introduced for purposes of gene therapy or for production of an encoded protein. Other examples of heterologous DNA include, but are not limited to, DNA that encodes traceable marker proteins, such as a protein that confers drug resistance, DNA that encodes therapeutically effective substances, such as anti-

-24-

cancer agents, enzymes and hormones, and DNA that encodes other types of proteins, such as antibodies. Antibodies that are encoded by heterologous DNA may be secreted or expressed on the surface of the cell in which the heterologous DNA has been introduced.

5 As used herein, a therapeutically effective product is a product that is encoded by heterologous DNA that, upon introduction of the DNA into a host, a product is expressed that effectively ameliorates or eliminates the symptoms, manifestations of an inherited or acquired disease or that cures said disease.

10 As used herein, transgenic plants refer to plants in which heterologous or foreign DNA is expressed or in which the expression of a gene naturally present in the plant has been altered.

 As used herein, operative linkage of heterologous DNA to regulatory and effector sequences of nucleotides, such as promoters, enhancers, transcriptional and translational stop sites, and other signal sequences refers to the relationship between such DNA and such sequences of nucleotides. For example, operative linkage of heterologous DNA to a promoter refers to the physical relationship between the DNA and the promoter such that the transcription of such DNA is initiated from the promoter by an RNA polymerase that specifically recognizes, binds to and transcribes the DNA in reading frame. Preferred promoters include tissue specific promoters, such as mammary gland specific promoters, viral promoters, such as TK, CMV, adenovirus promoters, and other promoters known to those of skill in the art.

15
20
25

 As used herein, isolated, substantially pure DNA refers to DNA fragments purified according to standard techniques employed by those skilled in the art, such as that found in Maniatis et al. [(1982) Molecular

-25-

Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY].

As used herein, expression refers to the process by which nucleic acid is transcribed into mRNA and translated into peptides, polypeptides, or proteins. If the nucleic acid is derived from genomic DNA, expression may, if an appropriate eukaryotic host cell or organism is selected, include splicing of the mRNA.

As used herein, vector or plasmid refers to discrete elements that are used to introduce heterologous DNA into cells for either expression of the heterologous DNA or for replication of the cloned heterologous DNA. Selection and use of such vectors and plasmids are well within the level of skill of the art.

As used herein, transformation/transfection refers to the process by which DNA or RNA is introduced into cells. Transfection refers to the taking up of exogenous nucleic acid, e.g., an expression vector, by a host cell whether or not any coding sequences are in fact expressed. Numerous methods of transfection are known to the ordinarily skilled artisan, for example, by direct uptake using calcium phosphate [CaPO₄; see, e.g., Wigler et al. (1979) Proc. Natl. Acad. Sci. U.S.A. 76:1373-1376], polyethylene glycol [PEG]-mediated DNA uptake, electroporation, lipofection [see, e.g., Strauss (1996) Meth. Mol. Biol. 54:307-327], microcell fusion [see, EXAMPLES, see, also Lambert (1991) Proc. Natl. Acad. Sci. U.S.A. 88:5907-5911; U.S. Patent No. 5,396,767, Sawford et al. (1987) Somatic Cell Mol. Genet. 13:279-284; Dhar et al. (1984) Somatic Cell Mol. Genet. 10:547-559; and McNeill-Killary et al. (1995) Meth. Enzymol. 254:133-152], lipid-mediated carrier systems [see, e.g., Teifel et al. (1995) Biotechniques 19:79-80; Albrecht et al. (1996) Ann. Hematol. 72:73-79; Holmen et al. (1995) In Vitro Cell Dev. Biol. Anim. 31:347-351; Remy et al. (1994) Bioconjug. Chem. 5:647-654; Le Bolch

-26-

et al. (1995) Tetrahedron Lett. 36:6681-6684; Loeffler et al. (1993) Meth. Enzymol. 217:599-618] or other suitable method. Successful transfection is generally recognized by detection of the presence of the heterologous nucleic acid within the transfected cell, such as any

5 indication of the operation of a vector within the host cell.

Transformation means introducing DNA into an organism so that the DNA is replicable, either as an extrachromosomal element or by chromosomal integration.

As used herein, injected refers to the microinjection [use of a small
10 syringe] of DNA into a cell.

As used herein, substantially homologous DNA refers to DNA that includes a sequence of nucleotides that is sufficiently similar to another such sequence to form stable hybrids under specified conditions.

It is well known to those of skill in this art that nucleic acid
15 fragments with different sequences may, under the same conditions, hybridize detectably to the same "target" nucleic acid. Two nucleic acid fragments hybridize detectably, under stringent conditions over a sufficiently long hybridization period, because one fragment contains a segment of at least about 14 nucleotides in a sequence which is
20 complementary [or nearly complementary] to the sequence of at least one segment in the other nucleic acid fragment. If the time during which hybridization is allowed to occur is held constant, at a value during which, under preselected stringency conditions, two nucleic acid fragments with exactly complementary base-pairing segments hybridize
25 detectably to each other, departures from exact complementarity can be introduced into the base-pairing segments, and base-pairing will nonetheless occur to an extent sufficient to make hybridization detectable. As the departure from complementarity between the base-pairing segments of two nucleic acids becomes larger, and as conditions

-27-

of the hybridization become more stringent, the probability decreases that the two segments will hybridize detectably to each other.

Two single-stranded nucleic acid segments have "substantially the same sequence," within the meaning of the present specification, if

- 5 (a) both form a base-paired duplex with the same segment, and (b) the melting temperatures of said two duplexes in a solution of 0.5 X SSPE differ by less than 10°C. If the segments being compared have the same number of bases, then to have "substantially the same sequence", they will typically differ in their sequences at fewer than 1 base in 10.
- 10 Methods for determining melting temperatures of nucleic acid duplexes are well known [see, e.g., Meinkoth and Wahl (1984) Anal. Biochem. 138:267-284 and references cited therein].

- As used herein, a nucleic acid probe is a DNA or RNA fragment that includes a sufficient number of nucleotides to specifically hybridize
- 15 to DNA or RNA that includes identical or closely related sequences of nucleotides. A probe may contain any number of nucleotides, from as few as about 10 and as many as hundreds of thousands of nucleotides. The conditions and protocols for such hybridization reactions are well known to those of skill in the art as are the effects of probe size,
 - 20 temperature, degree of mismatch, salt concentration and other parameters on the hybridization reaction. For example, the lower the temperature and higher the salt concentration at which the hybridization reaction is carried out, the greater the degree of mismatch that may be present in the hybrid molecules.

- 25 To be used as a hybridization probe, the nucleic acid is generally rendered detectable by labelling it with a detectable moiety or label, such as ^{32}P , ^3H and ^{14}C , or by other means, including chemical labelling, such as by nick-translation in the presence of deoxyuridylate biotinylated at the 5'-position of the uracil moiety. The resulting probe includes the

-28-

biotinylated uridylate in place of thymidylate residues and can be detected [via the biotin moieties] by any of a number of commercially available detection systems based on binding of streptavidin to the biotin. Such commercially available detection systems can be obtained, for

5 example, from Enzo Biochemicals, Inc. [New York, NY]. Any other label known to those of skill in the art, including non-radioactive labels, may be used as long as it renders the probes sufficiently detectable, which is a function of the sensitivity of the assay, the time available [for culturing cells, extracting DNA, and hybridization assays], the quantity of DNA or

10 RNA available as a source of the probe, the particular label and the means used to detect the label.

Once sequences with a sufficiently high degree of homology to the probe are identified, they can readily be isolated by standard techniques, which are described, for example, by Maniatis et al. ((1982) Molecular

15 Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY).

As used herein, conditions under which DNA molecules form stable hybrids and are considered substantially homologous are such that DNA molecules with at least about 60% complementarity form stable

20 hybrids. Such DNA fragments are herein considered to be "substantially homologous". For example, DNA that encodes a particular protein is substantially homologous to another DNA fragment if the DNA forms stable hybrids such that the sequences of the fragments are at least about 60% complementary and if a protein encoded by the DNA retains

25 its activity.

For purposes herein, the following stringency conditions are defined:

- 1) high stringency: 0.1 x SSPE, 0.1% SDS, 65 °C
- 2) medium stringency: 0.2 x SSPE, 0.1% SDS, 50°C

-29-

3) low stringency: 1.0 x SSPE, 0.1% SDS, 50°C
or any combination of salt and temperature and other reagents that result
in selection of the same degree of mismatch or matching.

As used herein, immunoprotective refers to the ability of a vaccine
5 or exposure to an antigen or immunity-inducing agent, to confer upon a
host to whom the vaccine or antigen is administered or introduced, the
ability to resist infection by a disease-causing pathogen or to have
reduced symptoms. The selected antigen is typically an antigen that is
presented by the pathogen.

10 As used herein, all assays and procedures, such as hybridization
reactions and antibody-antigen reactions, unless otherwise specified, are
conducted under conditions recognized by those of skill in the art as
standard conditions.

A. Preparation of cell lines containing MACs

15 **1. The megareplicon**

The methods, cells and MACs provided herein are produced by
virtue of the discovery of the existence of a higher-order replication unit
[megareplicon] of the centromeric region. This megareplicon is delimited
by a primary replication initiation site [megareplicator], and appears to
20 facilitate replication of the centromeric heterochromatin, and most likely,
centromeres. Integration of heterologous DNA into the megareplicator
region or in close proximity thereto, initiates a large-scale amplification of
megabase-size chromosomal segments, which leads to *de novo*
chromosome formation in living cells.

25 DNA sequences that provide a preferred megareplicator are the
rDNA units that give rise to ribosomal RNA (rRNA). In mammals,
particularly mice and humans, these rDNA units contain specialized
elements, such as the origin of replication (or origin of bidirectional
replication, i.e., OBR, in mouse) and amplification promoting sequences

-30-

- (APS) and amplification control elements (ACE) (see, e.g., Gogel et al. (1996) Chromosoma 104:511-518; Coffman et al. (1993) Exp. Cell. Res. 209:123-132; Little et al. (1993) Mol. Cell. Biol. 13:6600-6613; Yoon et al. (1995) Mol. Cell. Biol. 15:2482-2489; Gonzalez and Sylvester (1995) Genomics 27:320-328; Miesfeld and Arnheim (1982) Nuc. Acids Res. 10:3933-3949)); Maden et al. (1987) Biochem. J. 246:519-527).

- As described herein, without being bound by any theory, these specialized elements may facilitate replication and/or amplification of megabase-size chromosomal segments in the *de novo* formation of
- 10 chromosomes, such as those described herein, in cells. These specialized elements are typically located in the nontranscribed intergenic spacer region upstream of the transcribed region of rDNA. The intergenic spacer region may itself contain internally repeated sequences which can be classified as tandemly repeated blocks and nontandem blocks (see
- 15 e.g., Gonzalez and Sylvester (1995) Genomics 27:320-328). In mouse rDNA, an origin of bidirectional replication may be found within a 3-kb initiation zone centered approximately 1.6 kb upstream of the transcription start site (see, e.g., Gogel et al. (1996) Chromosoma 104:511-518). The sequences of these specialized elements tend to
- 20 have an altered chromatin structure, which may be detected, for example, by nuclease hypersensitivity or the presence of AT-rich regions that can give rise to bent DNA structures. An exemplary sequence encompassing an origin of replication is shown in SEQ ID NO. 16 and in GENBANK accession no. X82564 at about positions 2430-5435.
- 25 Exemplary sequences encompassing amplification-promoting sequences include nucleotides 690-1060 and 1105-1530 of SEQ ID NO. 16.

In human rDNA, a primary replication initiation site may be found a few kilobase pairs upstream of the transcribed region and secondary initiation sites may be found throughout the nontranscribed intergenic

-31-

spacer region (see, e.g., Yoon et al. (1995) Mol. Cell. Biol. 15:2482-2489). A complete human rDNA repeat unit is presented in GENBANK as accession no. U13369 and is set forth in SEQ ID NO. 17 herein.

Another exemplary sequence encompassing a replication initiation site
5 may be found within the sequence of nucleotides 35355-42486 in
SEQ ID NO. 17 particularly within the sequence of nucleotides 37912-
42486 and more particularly within the sequence of nucleotides 37912-
39288 of SEQ ID NO. 17 (see Coffman et al. (1993) Exp. Cell. Res.
209:123-132).

10 Cell lines containing MACs can be prepared by transforming cells,
preferably a stable cell line, with a heterologous DNA fragment that
encodes a selectable marker, culturing under selective conditions, and
identifying cells that have a multicentric, typically dicentric, chromosome.
These cells can then be manipulated as described herein to produce the
15 minichromosomes and other MACs, particularly the heterochromatic
SATACs, as described herein.

Development of a multicentric, particularly dicentric, chromosome
typically is effected through integration of the heterologous DNA in the
pericentric heterochromatin, preferably in the centromeric regions of
20 chromosomes carrying rDNA sequences. Thus, the frequency of
incorporation can be increased by targeting to these regions, such as by
including DNA, including, but not limited to, rDNA or satellite DNA, in the
heterologous fragment that encodes the selectable marker. Among the
preferred targeting sequences for directing the heterologous DNA to the
25 pericentromeric heterochromatin are rDNA sequences that target
centromeric regions of chromosomes that carry rRNA genes. Such
sequences include, but are not limited to, the DNA of SEQ ID NO. 16 and
GENBANK accession no. X82564 and portions thereof, the DNA of SEQ
ID NO. 17 and GENBANK accession no. U13369 and portions thereof,

-32-

and the DNA of SEQ ID NOS. 18-24. A particular vector incorporating DNA from within SEQ ID NO. 16 for use in directing integration of heterologous DNA into chromosomal rDNA is pTERPUD (see Example 12). Satellite DNA sequences can also be used to direct the

5 heterologous DNA to integrate into the pericentric heterochromatin. For example, vectors pTEMPUD and pHASPUD, which contain mouse and human satellite DNA, respectively, are provided herein (see Example 12) as exemplary vectors for introduction of heterologous DNA into cells for *de novo* artificial chromosome formation.

10 The resulting cell lines can then be treated as the exemplified cells herein to produce cells in which the dicentric chromosome has fragmented. The cells can then be used to introduce additional selective markers into the fragmented dicentric chromosome (*i.e.*, formerly dicentric chromosome), whereby amplification of the pericentric

15 heterochromatin will produce the heterochromatic chromosomes.

The following discussion describes this process with reference to the EC3/7 line and the resulting cells. The same procedures can be applied to any other cells, particularly cell lines to create SATACs and euchromatic minichromosomes.

20 2. Formation of *de novo* chromosomes

De novo centromere formation in a transformed mouse LMTK-fibroblast cell line [EC3/7] after cointegration of λ constructs [λ CM8 and λ gtWESneo] carrying human and bacterial DNA [Hadlaczy *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110 and U.S.

25 application Serial No. 08/375,271] has been shown. The integration of the "heterologous" engine red human, bacterial and phage DNA, and the subsequent amplification of mouse and heterologous DNA that led to the formation of a dicentric chromosome, occurred at the centromeric region of the short arm of a mouse chromosome. By G-banding, this

-33-

chromosome was identified as mouse chromosome 7. Because of the presence of two functionally active centromeres on the same chromosome, regular breakages occur between the centromeres. Such specific chromosome breakages gave rise to the appearance [in
5 approximately 10% of the cells] of a chromosome fragment carrying the neo-centromere. From the EC3/7 cell line [see, U.S. Patent No. 5,288,625, deposited at the European Collection of Animal Cell Culture (hereinafter ECACC) under accession no. 90051001; see, also Hadlaczký et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110, and U.S.
10 application Serial No. 08/375,271 and the corresponding published European application EP 0 473 253, two sublines [EC3/7C5 and EC3/7C6] were selected by repeated single-cell cloning. In these cell lines, the neo-centromere was found exclusively on a minichromosome [neo-minichromosome], while the formerly dicentric chromosome carried
15 traces of "heterologous" DNA.

It has now been discovered that integration of DNA encoding a selectable marker in the heterochromatic region of the centromere led to formation of the dicentric chromosome.

3. The neo-minichromosome

20 The chromosome breakage in the EC3/7 cells, which separates the neo-centromere from the mouse chromosome, occurred in the G-band positive "heterologous" DNA region. This is supported by the observation of traces of λ and human DNA sequences at the broken end of the formerly dicentric chromosome. Comparing the G-band pattern of the
25 chromosome fragment carrying the neo-centromere with that of the stable neo-minichromosome, it is apparent that the neo-minichromosome is an inverted duplicate of the chromosome fragment that bears the neo-centromere. This is supported by the observation that although the neo-minichromosome carries only one functional centromere, both ends of

-34-

the minichromosome are heterochromatic, and mouse satellite DNA sequences were found in these heterochromatic regions by *in situ* hybridization.

Mouse cells containing the minichromosome, which contains
5 multiple repeats of the heterologous DNA, which in the exemplified embodiment is λ DNA and the neomycin-resistance gene, can be used as recipient cells in cell transformation. Donor DNA, such as selected heterologous DNA containing λ DNA linked to a second selectable marker, such as the gene encoding hygromycin phosphotransferase
10 which confers hygromycin resistance [hyg], can be introduced into the mouse cells and integrated into the minichromosomes by homologous recombination of λ DNA in the donor DNA with that in the minichromosomes. Integration is verified by *in situ* hybridization and Southern blot analyses. Transcription and translation of the heterologous
15 DNA is confirmed by primer extension and immunoblot analyses.

For example, DNA has been targeted into the neo-minichromosome in EC3/7C5 cells using a λ DNA-containing construct [pNem1ruc] that also contains DNA encoding hygromycin resistance and the *Renilla* luciferase gene linked to a promoter, such as the cytomegalovirus [CMV]
20 early promoter, and the bacterial neomycin resistance-encoding DNA. Integration of the donor DNA into the chromosome in selected cells [designated PHN4] was confirmed by nucleic acid amplification [PCR] and *in situ* hybridization. Events that would produce a neo-minichromosome are depicted in Figure 1.

25 The resulting engineered minichromosome that contains the heterologous DNA can then be transferred by cell fusion into a recipient cell line, such as Chinese hamster ovary cells [CHO] and correct expression of the heterologous DNA can be verified. Following production of the cells, metaphase chromosomes are obtained, such as

-35-

by addition of colchicine, and the chromosomes purified by addition of AT- and GC-specific dyes on a dual laser beam based cell sorter (see Example 10 B for a description of methods of isolating artificial chromomsomes). Preparative amounts of chromosomes [5×10^4 - 5×10^7 chromosomes/ml] at a purity of 95% or higher can be obtained. The resulting chromosomes are used for delivery to cells by methods such as microinjection and liposome-mediated transfer.

Thus, the neo-minichromosome is stably maintained in cells, replicates autonomously, and permits the persistent long-term expression of the *neo* gene under non-selective culture conditions. It also contains megabases of heterologous known DNA [λ DNA in the exemplified embodiments] that serves as target sites for homologous recombination and integration of DNA of interest. The neo-minichromosome is, thus, a vector for genetic engineering of cells. It has been introduced into SCID mice, and shown to replicate in the same manner as endogenous chromosomes.

The methods herein provide means to induce the events that lead to formation of the neo-minichromosome by introducing heterologous DNA with a selective marker [preferably a dominant selectable marker] into cells and culturing the cells under selective conditions. As a result, cells that contain a multicentric, e.g., dicentric chromosome, or fragments thereof, generated by amplification are produced. Cells with the dicentric chromosome can then be treated to destabilize the chromosomes with agents, such as BrdU and/or culturing under selective conditions, resulting in cells in which the dicentric chromosome has formed two chromosomes, a so-called minichromosome, and a formerly dicentric chromosome that has typically undergone amplification in the heterochromatin where the heterologous DNA has integrated to produce a SATAC or a sausage chromosome [discussed below]. These cells can

-36-

be fused with other cells to separate the minichromosome from the formerly dicentric chromosome into different cells so that each type of MAC can be manipulated separately.

4. Preparation of SATACs

- 5 An exemplary protocol for preparation of SATACs is illustrated in Figure 2 [particularly D, E and F] and FIGURE 3 [see, also the EXAMPLES, particularly EXAMPLES 4-7].

To prepare a SATAC, the starting materials are cells, preferably a stable cell line, such as a fibroblast cell line, and a DNA fragment that
10 includes DNA that encodes a selective marker. The DNA fragment is introduced into the cell by methods of DNA transfer, including but not limited to direct uptake using calcium phosphate, electroporation, and lipid-mediated transfer. To insure integration of the DNA fragment in the heterochromatin, it is preferable to start with DNA that will be targeted
15 to the pericentric heterochromatic region of the chromosome, such as λ CM8 and vectors provided herein, such as pTEMPUD [Figure 5] and pHASPU (see Example 12) that include satellite DNA, or specifically into rDNA in the centromeric regions of chromosomes containing rDNA sequences. After introduction of the DNA, the cells are grown under
20 selective conditions. The resulting cells are examined and any that have multicentric, particularly dicentric, chromosomes [or heterochromatic chromosomes or sausage chromosomes or other such structure; see, Figure 2D, 2E and 2F] are selected.

In particular, if a cell with a dicentric chromosome is selected, it
25 can be grown under selective conditions, or, preferably, additional DNA encoding a second selectable marker is introduced, and the cells grown under conditions selective for the second marker. The resulting cells should include chromosomes that have structures similar to those depicted in Figures 2D, 2E, 2F. Cells with a structure, such as the

-37-

sausage chromosome, Figure 2D, can be selected and fused with a second cell line to eliminate other chromosomes that are not of interest. If desired, cells with other chromosomes can be selected and treated as described herein. If a cell with a sausage chromosome is selected, it can

5 be treated with an agent, such as BrdU, that destabilizes the chromosome so that the heterochromatic arm forms a chromosome that is substantially heterochromatic [i.e., a megachromosome, see, Figure 2F]. Structures such as the gigachromosome in which the heterochromatic arm has amplified but not broken off from the

10 euchromatic arm, will also be observed. The megachromosome is a stable chromosome. Further manipulation, such as fusions and growth in selective conditions and/or BrdU treatment or other such treatment, can lead to fragmentation of the megachromosome to form smaller chromosomes that have the amplicon as the basic repeating unit.

15 The megachromosome can be further fragmented in vivo using a chromosome fragmentation vector, such as pTEMPUD [see, Figure 5 and EXAMPLE 12], pHASPUd or pTERPUd (see Example 12) to ultimately produce a chromosome that comprises a smaller stable replicable unit, about 15 Mb-60 Mb, containing one to four megareplicons.

20 Thus, the stable chromosomes formed *de novo* that originate from the short arm of mouse chromosome 7 have been analyzed. This chromosome region shows a capacity for amplification of large chromosome segments, and promotes *de novo* chromosome formation. Large-scale amplification at the same chromosome region leads to the

25 formation of dicentric and multicentric chromosomes, a minichromosome, the 150-200 Mb size λ neo-chromosome, the "sausage" chromosome, the 500-1000 Mb gigachromosome, and the stable 250-400 Mb megachromosome.

-38-

A clear segmentation is observed along the arms of the megachromosome, and analyses show that the building units of this chromosome are amplicons of ~30 Mb composed of mouse major satellite DNA with the integrated "foreign" DNA sequences at both ends.

5 The ~30 Mb amplicons are composed of two ~15 Mb inverted doublets of ~7.5 Mb mouse major satellite DNA blocks, which are separated from each other by a narrow band of non-satellite sequences [see, e.g., Figure 3]. The wider non-satellite regions at the amplicon borders contain integrated, exogenous [heterologous] DNA, while the narrow

10 bands of non-satellite DNA sequences within the amplicons are integral parts of the pericentric heterochromatin of mouse chromosomes. These results indicate that the ~7.5 Mb blocks flanked by non-satellite DNA are the building units of the pericentric heterochromatin of mouse chromosomes, and the ~15 Mb size pericentric regions of mouse

15 chromosomes contain two ~7.5 Mb units.

Apart from the euchromatic terminal segments, the whole megachromosome is heterochromatic, and has structural homogeneity. Therefore, this large chromosome offers a unique possibility for obtaining information about the amplification process, and for analyzing some basic

20 characteristics of the pericentric constitutive heterochromatin, as a vector for heterologous DNA, and as a target for further fragmentation.

As shown herein, this phenomenon is generalizable and can be observed with other chromosomes. Also, although these *de novo* formed chromosome segments and chromosomes appear different, there are

25 similarities that indicate that a similar amplification mechanism plays a role in their formation: (i) in each case, the amplification is initiated in the centromeric region of the mouse chromosomes and large (Mb size) amplicons are formed; (ii) mouse major satellite DNA sequences are constant constituents of the amplicons, either by providing the bulk of

-39-

the heterochromatic amplicons [H-type amplification], or by bordering the
aeuchromatic amplicons [E-type amplification]; (iii) formation of inverted
segments can be demonstrated in the λ neo-chromosome and
megachromosome; (iv) chromosome arms and chromosomes formed by
5 the amplification are stable and functional.

The presence of inverted chromosome segments seems to be a
common phenomenon in the chromosomes formed *de novo* at the
centromeric region of mouse chromosome 7. During the formation of the
neo-minichromosome, the event leading to the stabilization of the distal
10 segment of mouse chromosome 7 that bears the neo-centromere may
have been the formation of its inverted duplicate. Amplicons of the
megachromosome are inverted doublets of ~7.5 Mb mouse major
satellite DNA blocks.

5. Cell lines

15 Cell lines that contain MACs, such as the minichromosome, the λ -
neo chromosome, and the SATACs are provided herein or can be
produced by the methods herein. Such cell lines provide a convenient
source of these chromosomes and can be manipulated, such as by cell
fusion or production of microcells for fusion with selected cell lines, to
20 deliver the chromosome of interest into hybrid cell lines. Exemplary cell
lines are described herein and some have been deposited with the
ECACC.

a. EC3/7C5 and EC3/7C6

Cell lines EC3/7C5 and EC3/7C6 were produced by single cell
25 cloning of EC3/7. For exemplary purposes EC3/7C5 has been deposited
with the ECACC. These cell lines contain a minichromosome and the
formerly dicentric chromosome from EC3/7. The stable mini-
chromosomes in cell lines EC3/7C5 and EC3/7C6 appear to be the same
and they seem to be duplicated derivatives of the ~10-15 Mb

-40-

"broken-off" fragment of the dicentric chromosome. Their similar size in these independently generated cell lines might indicate that ~ 20-30 Mb is the minimal or close to the minimal physical size for a stable minichromosome.

5 b. TF1004G19

Introduction of additional heterologous DNA, including DNA encoding a second selectable marker, hygromycin phosphotransferase, i.e., the hygromycin-resistance gene, and also a detectable marker, β -galactosidase (i.e., encoded by the lacZ gene), into the EC3/7C5 cell line
10 and growth under selective conditions produced cells designated TF1004G19. In particular, this cell line was produced from the EC3/7C5 cell line by cotransfection with plasmids pH132, which contains an anti-HIV ribozyme and hygromycin-resistance gene, pCH110 [encodes β -galactosidase] and λ phage [λ cl 875 Sam 7] DNA and selection with
15 hygromycin B.

Detailed analysis of the TF1004G19 cell line by *in situ* hybridization with λ phage and plasmid DNA sequences revealed the formation of the sausage chromosome. The formerly dicentric chromosome of the EC3/7C5 cell line translocated to the end of another
20 acrocentric chromosome. The heterologous DNA integrated into the pericentric heterochromatin of the formerly dicentric chromosome and is amplified several times with megabases of mouse pericentric heterochromatic satellite DNA sequences [Fig. 2D] forming the "sausage" chromosome. Subsequently the acrocentric mouse chromosome was
25 substituted by a euchromatic telomere.

In situ hybridization with biotin-labeled subfragments of the hygromycin-resistance and β -galactosidase genes resulted in a hybridization signal only in the heterochromatic arm of the sausage

-41-

chromosome, indicating that in TF1004G19 transformant cells these genes are localized in the pericentric heterochromatin.

A high level of gene expression, however, was detected. In general, heterochromatin has a silencing effect in *Drosophila*, yeast and
5 on the HSV-tk gene introduced into satellite DNA at the mouse centromere. Thus, it was of interest to study the TF1004G19 transformed cell line to confirm that genes located in the heterochromatin were indeed expressed, contrary to recognized dogma.

For this purpose, subclones of TF1004G19, containing a different
10 sausage chromosome [see Figure 2D], were established by single cell cloning. Southern hybridization of DNA isolated from the subclones with subfragments of hygromycin phosphotransferase and lacZ genes showed a close correlation between the intensity of hybridization and the length of the sausage chromosome. This finding supports the conclusion that
15 these genes are localized in the heterochromatic arm of the sausage chromosome.

(1) TF1004G-19C5

TF1004G-19C5 is a mouse LMTK⁻ fibroblast cell line containing neo-minichromosomes and stable "sausage" chromosomes. It is a
20 subclone of TF1004G19 and was generated by single-cell cloning of the TF1004G19 cell line. It has been deposited with the ECACC as an exemplary cell line and exemplary source of a sausage chromosome. Subsequent fusion of this cell line with CHO K20 cells and selection with hygromycin and G418 and HAT (hypoxanthine, aminopterin, and
25 thymidine medium; see Szybalski *et al.* (1962) Proc. Natl. Acad. Sci. 48:2026) resulted in hybrid cells (designated 19C5xHa4) that carry the sausage chromosome and the neo-minichromosome. BrdU treatment of the hybrid cells, followed by single cell cloning and selection with G418

-42-

and/or hygromycin produced various cells that carry chromosomes of interest, including GB43 and G3D5.

(2) other subclones

Cell lines GB43 and G3D5 were obtained by treating 19C5xHa4
5 cells with BrdU followed by growth in G418-containing selective medium
and retreatment with BrdU. The two cell lines were isolated by single
cell cloning of the selected cells. GB43 cells contain the neo-
minichromosome only. G3D5, which has been deposited with the
ECACC, carries the neo-minichromosome and the megachromosome.
10 Single cell cloning of this cell line followed by growth of the subclones in
G418- and hygromycin-containing medium yielded subclones such as the
GHB42 cell line carrying the neo-minichromosome and the
megachromosome. H1D3 is a mouse-hamster hybrid cell line carrying
the megachromosome, but no neo-minichromosome, and was generated
15 by treating 19C5xHa4 cells with BrdU followed by growth in hygromycin-
containing selective medium and single cell subcloning of selected cells.
Fusion of this cell line with the CD4⁺ HeLa cell line that also carries DNA
encoding an additional selection gene, the neomycin-resistance gene,
produced cells [designated H1xHE41 cells] that carry the
20 megachromosome as well as a human chromosome that carries CD4neo.
Further BrdU treatment and single cell cloning produced cell lines, such
as 1B3, that include cells with a truncated megachromosome.

5. DNA constructs used to transform the cells

Heterologous DNA can be introduced into the cells by transfection
25 or other suitable method at any stage during preparation of the
chromosomes [see, e.g., FIG. 4]. In general, incorporation of such DNA
into the MACs is assured through site-directed integration, such as may
be accomplished by inclusion of λ -DNA in the heterologous DNA (for the
exemplified chromosomes), and also an additional selective marker gene.

-43-

For example, cells containing a MAC, such as the minichromosome or a SATAC, can be cotransfected with a plasmid carrying the desired heterologous DNA, such as DNA encoding an HIV ribozyme, the cystic fibrosis gene, and DNA encoding a second selectable marker, such as

5 hygromycin resistance. Selective pressure is then applied to the cells by exposing them to an agent that is harmful to cells that do not express the new selectable marker. In this manner, cells that include the heterologous DNA in the MAC are identified. Fusion with a second cell line can provide a means to produce cell lines that contain one particular

10 type of chromosomal structure or MAC.

Various vectors for this purpose are provided herein [see, Examples] and others can be readily constructed. The vectors preferably include DNA that is homologous to DNA contained within a MAC in order to target the DNA to the MAC for integration therein. The vectors also

15 include a selectable marker gene and the selected heterologous gene(s) of interest. Based on the disclosure herein and the knowledge of the skilled artisan, one of skill can construct such vectors.

Of particular interest herein is the vector pTEMPUD and derivatives thereof that can target DNA into the heterochromatic region of selected

20 chromosomes. These vectors can also serve as fragmentation vectors [see, e.g., Example 12].

Heterologous genes of interest include any gene that encodes a therapeutic product and DNA encoding gene products of interest. These genes and DNA include, but are not limited to: the cystic fibrosis gene

25 [CF], the cystic fibrosis transmembrane regulator (CFTR) gene [see, e.g., U.S. Patent No. 5,240,846; Rosenfeld et al. (1992) Cell 68:143-155; Hyde et al. (1993) Nature 362: 250-255; Kerem et al. (1989) Science 245:1073-1080; Riordan et al. (1989) Science 245:1066-1072; Rommens et al. (1989) Science 245:1059-1065; Osborne et al. (1991)

-44-

Am. J. Hum. Genetics 48:6089-6122; White et al. (1990) Nature 344:665-667; Dean et al. (1990) Cell 61:863-870; Erlich et al. (1991) Science 252:1643; and U.S. Patent Nos. 5,453,357, 5,449,604, 5,434,086, and 5,240,846, which provides a retroviral vector encoding
5 the normal CFTR gene].

B. Isolation of artificial chromosomes

The MACs provided herein can be isolated by any suitable method known to those of skill in the art. Also, methods are provided herein for effecting substantial purification, particularly of the SATACs. SATACs
10 have been isolated by fluorescence-activated cell sorting [FACS]. This method takes advantage of the nucleotide base content of the SATACs, which, by virtue of their high heterochromatic DNA content, will differ from any other chromosomes in a cell. In particular embodiment, metaphase chromosomes are isolated and stained with base-specific
15 dyes, such as Hoechst 33258 and chromomycin A3. Fluorescence-activated cell sorting will separate the SATACs from the endogenous chromosomes. A dual-laser cell sorter [FACS Vantage Becton Dickinson Immunocytometry Systems] in which two lasers were set to excite the dyes separately, allowed a bivariate analysis of the chromosomes by
20 base-pair composition and size. Cells containing such SATACs can be similarly sorted.

Additional methods provided herein for isolation of artificial chromosomes from endogenous chromosomes include procedures that are particularly well suited for large-scale isolation of artificial
25 chromosomes such as SATACs. In these methods, the size and density differences between SATACs and endogenous chromosomes are exploited to effect separation of these two types of chromosomes. Such methods involve techniques such as swinging bucket centrifugation, zonal rotor centrifugation, and velocity sedimentation. Affinity-,

-45-

particularly immunoaffinity-, based methods for separation of artificial from endogenous chromosomes are also provided herein. For example, SATACs, which are predominantly heterochromatin, may be separated from endogenous chromosomes through immunoaffinity procedures

- 5 involving antibodies that specifically recognize heterochromatin, and/or the proteins associated therewith, when the endogenous chromosomes contain relatively little heterochromatin, such as in hamster cells.

C. *In vitro* construction of artificial chromosomes

- Artificial chromosomes can be constructed in vitro by assembling
- 10 the structural and functional elements that contribute to a complete chromosome capable of stable replication and segregation alongside endogenous chromosomes in cells. The identification of the discrete elements that in combination yield a functional chromosome has made possible the in vitro generation of artificial chromosomes. The process of
- 15 in vitro construction of artificial chromosomes, which can be rigidly controlled, provides advantages that may be desired in the generation of chromosomes that, for example, are required in large amounts or that are intended for specific use in transgenic animal systems.

- For example, in vitro construction may be advantageous when
- 20 efficiency of time and scale are important considerations in the preparation of artificial chromosomes. Because in vitro construction methods do not involve extensive cell culture procedures, they may be utilized when the time and labor required to transform, feed, cultivate, and harvest cells used in in vivo cell-based production systems is
- 25 unavailable.

In vitro construction may also be rigorously controlled with respect to the exact manner in which the several elements of the desired artificial chromosome are combined and in what sequence and proportions they are assembled to yield a chromosome of precise specifications. These

-46-

aspects may be of significance in the production of artificial chromosomes that will be used in live animals where it is desirable to be certain that only very pure and specific DNA sequences in specific amounts are being introduced into the host animal.

5 The following describes the processes involved in the construction of artificial chromosomes in vitro, utilizing a megachromosome as exemplary starting material.

1. **Identification and isolation of the components of the artificial chromosome**

10 The MACs provided herein, particularly the SATACs, are elegantly simple chromosomes for use in the identification and isolation of components to be used in the in vitro construction of artificial chromosomes. The ability to purify MACs to a very high level of purity, as described herein, facilitates their use for these purposes. For
15 example, the megachromosome, particularly truncated forms thereof [i.e. cell lines, such as 1B3 and mM2C1, which are derived from H1D3 (deposited at the European Collection of Animal Cell Culture (ECACC) under Accession No. 96040929, see EXAMPLES below) serve as starting materials.

20 For example, the mM2C1 cell line contains a micro-megachromosome (~ 50-60 kB), which advantageously contains only one centromere, two regions of integrated heterologous DNA with adjacent rDNA sequences, with the remainder of the chromosomal DNA being mouse major satellite DNA. Other truncated megachromosomes can
25 serve as a source of telomeres, or telomeres can be provided (see, Examples below regarding construction of plasmids containing tandemly repeated telomeric sequences). The centromere of the mM2C1 cell line contains mouse minor satellite DNA, which provides a useful tag for isolation of the centromeric DNA.

-47-

Additional features of particular SATACs provided herein, such as the micro-megachromosome of the mM2C1 cell line, that make them uniquely suited to serve as starting materials in the isolation and identification of chromosomal components include the fact that the

5 centromeres of each megachromosome within a single specific cell line are identical. The ability to begin with a homogeneous centromere source (as opposed to a mixture of different chromosomes having differing centromeric sequences) greatly facilitates the cloning of the

10 truncated megachromosomes, such as the micro-megachromosome, with appropriate restriction endonucleases and cloning the fragments into the commercially available and well known YAC vectors (see, e.g., Burke et al. (1987) Science 236:806-812), BAC vectors (see, e.g., Shizuya et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89: 8794-8797 bacterial artificial

15 chromosomes which have a capacity of incorporating 0.9 - 1 Mb of DNA) or PAC vectors (the P1 artificial chromosome vector which is a P1 plasmid derivative that has a capacity of incorporating 300 kb of DNA and that is delivered to E. coli host cells by electroporation rather than by bacteriophage packaging; see, e.g., Ioannou et al. (1994) Nature

20 Genetics 6:84-89; Pierce et al. (1992) Meth. Enzymol. 216:549-574; Pierce et al. (1992) Proc. Natl. Acad. Sci. U.S.A. 89:2056-2060; U.S. Patent No. 5,300,431 and International PCT application No. WO 92/14819) vectors, it is possible for as few as 50 clones to represent the entire micro-megachromosome.

25 a. Centromeres

An exemplary centromere for use in the construction of a mammalian artificial chromosome is that contained within the megachromosome of any of the megachromosome-containing cell lines provided herein, such as, for example, H1D3 and derivatives thereof,

-48-

such as mM2C1 cells. Megachromosomes are isolated from such cell lines utilizing, for example, the procedures described herein, and the centromeric sequence is extracted from the isolated megachromosomes. For example, the megachromosomes may be separated into fragments
5 utilizing selected restriction endonucleases that recognize and cut at sites that, for instance, are primarily located in the replication and/or heterologous DNA integration sites and/or in the satellite DNA. Based on the sizes of the resulting fragments, certain undesired elements may be separated from the centromere-containing sequences. The centromere-
10 containing DNA, which could be as large as 1 Mb.

Probes that specifically recognize the centromeric sequences, such as mouse minor satellite DNA-based probes [see, e.g., Wong et al. (1988) Nucl. Acids Res. 16:11645-11661], may be used to isolate the centromere-containing YAC, BAC or PAC clones derived from the
15 megachromosome. Alternatively, or in conjunction with the direct identification of centromere-containing megachromosomal DNA, probes that specifically recognize the non-centromeric elements, such as probes specific for mouse major satellite DNA, the heterologous DNA and/or rDNA, may be used to identify and eliminate the non-centromeric DNA-
20 containing clones.

Additionally, centromere cloning methods described herein may be utilized to isolate the centromere-containing sequence of the megachromosome. For example, Example 12 describes the use of YAC vectors in combination with the murine tyrosinase gene and NMRI/Han
25 mice for identification of the centromeric sequence.

Once the centromere fragment has been isolated, it may be sequenced and the sequence information may in turn be used in PCR amplification of centromere sequences from megachromosomes or other sources of centromeres. Isolated centromeres may also be tested for

-49-

function in vivo by transferring the DNA into a host mammalian cell. Functional analysis may include, for example, examining the ability of the centromere sequence to bind centromere-binding proteins. The cloned centromere will be transferred to mammalian cells with a selectable
5 marker gene and the binding of a centromere-specific protein, such as anti-centromere antibodies (e.g., LU851, see, Hadlaczky et al. (1986) Exp. Cell Res. 167:1-15) can be used to assess function of the centromeres.

b. Telomeres

10 Preferred telomeres are the 1 kB synthetic telomere provided herein (see, Examples). A double synthetic telomere construct, which contains a 1 kB synthetic telomere linked to a dominant selectable marker gene that continues in an inverted orientation may be used for ease of manipulation. Such a double construct contains a series of
15 TTAGGG repeats 3' of the marker gene and a series of repeats of the inverted sequence, i.e., GGGATT, 5' of the marker gene as follows: (GGGATTT)_n---dominant marker gene---(TTAGGG)_n. Using an inverted marker provides an easy means for insertion, such as by blunt end ligation, since only properly oriented fragments will be selected.

20 c. Megareplicator

The megareplicator sequences, such as the rDNA, provided herein are preferred for use in in vitro constructs. The rDNA provides an origin of replication and also provides sequences that facilitate amplification of the artificial chromosome in vivo to increase the size of the chromosome
25 to, for example accommodate increasing copies of a heterologous gene of interest as well as continuous high levels of expression of the heterologous genes.

-50-

d. Filler heterochromatin

Filler heterochromatin, particularly satellite DNA, is included to maintain structural integrity and stability of the artificial chromosome and provide a structural base for carrying genes within the chromosome. The
5 satellite DNA is typically A/T-rich DNA sequence, such as mouse major satellite DNA, or G/C-rich DNA sequence, such as hamster natural satellite DNA. Sources of such DNA include any eukaryotic organisms that carry non-coding satellite DNA with sufficient A/T or G/C composition to promote ready separation by sequence, such as by FACS,
10 or by density gradients. The satellite DNA may also be synthesized by generating sequence containing monotone, tandem repeats of highly A/T- or G/C-rich DNA units.

The most suitable amount of filler heterochromatin for use in construction of the artificial chromosome may be empirically determined
15 by, for example, including segments of various lengths, increasing in size, in the construction process. Fragments that are too small to be suitable for use will not provide for a functional chromosome, which may be evaluated in cell-based expression studies, or will result in a chromosome of limited functional lifetime or mitotic and structural
20 stability.

e. Selectable marker

Any convenient selectable marker may be used and at any convenient locus in the MAC.

2. Combination of the isolated chromosomal elements

25 Once the isolated elements are obtained, they may be combined to generate the complete, functional artificial chromosome. This assembly can be accomplished for example, by in vitro ligation either in solution, LMP agarose or on microbeads. The ligation is conducted so that one end of the centromere is directly joined to a telomere. The other

-51-

end of the centromere, which serves as the gene-carrying chromosome arm, is built up from a combination of satellite DNA and rDNA sequence and may also contain a selectable marker gene. Another telomere is joined to the end of the gene-carrying chromosome arm. The gene-carrying arm is the site at which any heterologous genes of interest, for example, in expression of desired proteins encoded thereby, are incorporated either during *in vitro* construction of the chromosome or sometime thereafter.

3. Analysis and testing of the artificial chromosome

Artificial chromosomes constructed *in vitro* may be tested for functionality in *in vivo* mammalian cell systems, using any of the methods described herein for the SATACs, minichromosomes, or known to those of skill in the art.

4. Introduction of desired heterologous DNA into the *in vitro* synthesized chromosome

Heterologous DNA may be introduced into the *in vitro* synthesized chromosome using routine methods of molecular biology, may be introduced using the methods described herein for the SATACs, or may be incorporated into the *in vitro* synthesized chromosome as part of one of the synthetic elements, such as the heterochromatin. The heterologous DNA may be linked to a selected repeated fragment, and then the resulting construct may be amplified *in vitro* using the methods for such *in vitro* amplification provided herein (see the Examples).

D. Introduction of artificial chromosomes into cells, tissues, animals and plants

Suitable hosts for introduction of the MACs provided herein, include, but are not limited to, any animal or plant, cell or tissue thereof, including, but not limited to: mammals, birds, reptiles, amphibians, insects, fish, arachnids, tobacco, tomato, wheat, plants and algae. The MACs, if contained in cells, may be introduced by cell fusion or microcell

-52-

fusion or, if the MACs have been isolated from cells, they may be introduced into host cells by any method known to those of skill in this art, including but not limited to: direct DNA transfer, electroporation, lipid-mediated transfer, e.g., lipofection and liposomes, microprojectile bombardment, microinjection in cells and embryos, protoplast regeneration for plants, and any other suitable method [see, e.g., Weissbach et al. (1988) *Methods for Plant Molecular Biology*, Academic Press, N.Y., Section VIII, pp. 421-463; Grierson et al. (1988) *Plant Molecular Biology*, 2d Ed., Blackie, London, Ch. 7-9; see, also U.S. Patent Nos. 5,491,075; 5,482,928; and 5,424,409; see, also, e.g., U.S. Patent No. 5,470,708, which describes particle-mediated transformation of mammalian unattached cells].

Other methods for introducing DNA into cells include nuclear microinjection and bacterial protoplast fusion with intact cells.

Polycations, such as polybrene and polyornithine, may also be used. For various techniques for transforming mammalian cells, see e.g., Keown et al. *Methods in Enzymology* (1990) Vol. 185, pp. 527-537; and Mansour et al. (1988) *Nature* 336:348-352.

For example, isolated, purified artificial chromosomes can be injected into an embryonic cell line such as a human kidney primary embryonic cell line [ATCC accession number CRL 1573] or embryonic stem cells [see, e.g., Hogan et al. (1994) *Manipulating the Mouse Embryo, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, see, especially, pages 255-264 and Appendix 3].

Preferably the chromosomes are introduced by microinjection, using a system such as the Eppendorf automated microinjection system, and grown under selective conditions, such as in the presence of hygromycin B or neomycin.

1. Methods for introduction of chromosomes into hosts

Depending on the host cell used, transformation is done using standard techniques appropriate to such cells. These methods include any, including those described herein, known to those of skill in the art.

5 a. DNA uptake

For mammalian cells that do not have cell walls, the calcium phosphate precipitation method for introduction of exogenous DNA [see, e.g., Graham et al. (1978) Virology 52:456-457; Wigler et al. (1979) Proc. Natl. Acad. Sci. U.S.A. 76:1373-1376; and Current Protocols in
10 Molecular Biology, Vol. 1, Wiley Inter-Science, Supplement 14, Unit 9.1.1-9.1.9 (1990)] is often preferred. DNA uptake can be accomplished by DNA alone or in the presence of polyethylene glycol [PEG-mediated gene transfer], which is a fusion agent, or by any variations of such methods known to those of skill in the art [see, e.g., U.S. Pat. No.
15 4,684,611].

Lipid-mediated carrier systems are also among the preferred methods for introduction of DNA into cells [see, e.g., Teifel et al. (1995) Biotechniques 19:79-80; Albrecht et al. (1996) Ann. Hematol. 72:73-79; Holmen et al. (1995) In Vitro Cell Dev. Biol. Anim. 31:347-351; Remy et
20 al. (1994) Bioconjug. Chem. 5:647-654; Le Bolc'h et al. (1995) Tetrahedron Lett. 36:6681-6684; Loeffler et al. (1993) Meth. Enzymol. 217:599-618]. Lipofection [see, e.g., Strauss (1996) Meth. Mol. Biol. 54:307-327] may also be used to introduce DNA into cells. This method is particularly well-suited for transfer of exogenous DNA into chicken
25 cells (e.g., chicken blastodermal cells and primary chicken fibroblasts; see Brazolot et al. (1991) Mol. Reprod. Dev. 30:304-312). In particular, DNA of interest can be introduced into chickens in operative linkage with promoters from genes, such as lysozyme and ovalbumin, that are

-54-

expressed in the egg, thereby permitting expression of the heterologous DNA in the egg.

Additional methods useful in the direct transfer of DNA into cells include particle gun electroporation [see, e.g., U.S. Patent Nos. 4,955,378, 5 4,923,814, 4,476,004, 4,906,576 and 4,441,972] and virion-mediated gene transfer.

A commonly used approach for gene transfer in land plants involves the direct introduction of purified DNA into protoplasts. The three basic methods for direct gene transfer into plant cells include: 1) polyethylene glycol (PEG)-mediated DNA uptake, 2) electroporation-mediated DNA 10 uptake and 3) microinjection. In addition, plants may be transformed using ultrasound treatment [see, e.g., International PCT application publication No. WO 91/00358].

b. Electroporation

15 Electroporation involves providing high-voltage electrical pulses to a solution containing a mixture of protoplasts and foreign DNA to create reversible pores in the membranes of plant protoplasts as well as other cells. Electroporation is generally used for prokaryotes or other cells, such as plants that contain substantial cell-wall barriers. Methods for 20 effecting electroporation are well known [see, e.g., U.S. Patent Nos. 4,784,737, 5,501,967, 5,501,662, 5,019,034, 5,503,999; see, also Frommet et al. (1985) Proc. Natl. Acad. Sci. U.S.A. 82:5824-5828].

For example, electroporation is often used for transformation of plants [see, e.g., Ag Biotechnology News 7:3 and 17 25 (September/October 1990)]. In this technique, plant protoplasts are electroporated in the presence of the DNA of interest that also includes a phenotypic marker. Electrical impulses of high field strength reversibly permeabilize biomembranes allowing the introduction of the plasmids. Electroporated plant protoplasts reform the cell wall, divide, and form

-55-

plant callus. Transformed plant cells will be identified by virtue of the expressed phenotypic marker. The exogenous DNA may be added to the protoplasts in any form such as, for example, naked linear, circular or supercoiled DNA, DNA encapsulated in liposomes, DNA in spheroplasts, DNA in other plant protoplasts, DNA complexed with salts, and other methods.

c. Microcells

The chromosomes can be transferred by preparing microcells containing an artificial chromosome and then fusing with selected target cells. Methods for such preparation and fusion of microcells are well known [see the Examples and also see, e.g., U.S. Patent Nos. 5,240,840, 4,806,476, 5,298,429, 5,396,767, Fournier (1981) Proc. Natl. Acad. Sci. U.S.A. 78:6349-6353; and Lambert et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:5907-59]. Microcell fusion, using microcells that contain an artificial chromosome, is a particularly useful method for introduction of MACs into avian cells, such as DT40 chicken pre-B cells [for a description of DT40 cell fusion, see, e.g., Dieken et al. (1996) Nature Genet. 12:174-182].

2. Hosts

Suitable hosts include any host known to be useful for introduction and expression of heterologous DNA. Of particular interest herein, animal and plant cells and tissues, including, but not limited to insect cells and larvae, plants, and animals, particularly transgenic (non-human) animals, and animal cells. Other hosts include, but are not limited to mammals, birds, particularly fowl such as chickens, reptiles, amphibians, insects, fish, arachnids, tobacco, tomato, wheat, monocots, dicots and algae, and any host into which introduction of heterologous DNA is

-56-

desired. Such introduction can be effected using the MACs provided herein, or, if necessary by using the MACs provided herein to identify species-specific centromeres and/or functional chromosomal units and then using the resulting centromeres or chromosomal units as artificial
5 chromosomes, or alternatively, using the methods exemplified herein for production of MACs to produce species-specific artificial chromosomes.

a. Introduction of DNA into embryos for production of transgenic (non-human) animals and introduction of DNA into animal cells

10 Transgenic (non-human) animals can be produced by introducing exogenous genetic material into a pronucleus of a mammalian zygote by microinjection [see, e.g., U.S. Patent Nos. 4,873,191 and 5,354,674; see, also, International PCT application publication No. WO 95/14769, which is based on U.S. application Serial No. 08/159,084]. The zygote
15 is capable of development into a mammal. The embryo or zygote is transplanted into a host female uterus and allowed to develop. Detailed protocols and examples are set forth below.

Nuclear transfer [see, Wilmut et al. (1997) Nature 385:810-813, International PCT application Nos. WO 97/07669 and WO 97/07668].

20 Briefly in this method, the SATAC containing the genes of interest is introduced by any suitable method, into an appropriate donor cell, such as a mammary gland cell, that contains totipotent nuclei. The diploid nucleus of the cell, which is either in G0 or G1 phase, is then introduced, such as by cell fusion or microinjection, into an unactivated oöcyte,
25 preferably enucleated cell, which is arrested in the metaphase of the second meiotic division. Enucleation may be effected by any suitable method, such as actual removal, or by treating with means, such as ultraviolet light, that functionally remove the nucleus. The oöcyte is then activated, preferably after a period of contact, about 6-20 hours for
30 cattle, of the new nucleus with the cytoplasm, while maintaining correct

-57-

ploidy, to produce a reconstituted embryo, which is then introduced into a host. Ploidy is maintained during activation, for example, by incubating the reconstituted cell in the presence of a microtubule inhibitor, such as nocodazole, colchicine, cocemid, and taxol, whereby the DNA replicates
5 once.

Transgenic chickens can be produced by injection of dispersed blastodermal cells from Stage X chicken embryos into recipient embryos at a similar stage of development [see e.g., Etches et al. (1993) Poultry Sci. 72:882-889; Petitte et al. (1990) Development 108:185-189].

10 Heterologous DNA is first introduced into the donor blastodermal cells using methods such as, for example, lipofection [see, e.g., Brazolot et al. (1991) Mol. Repro. Dev. 30:304-312] or microcell fusion [see, e.g., Dieken et al. (1996) Nature Genet. 12:174-182]. The transfected donor cells are then injected into recipient chicken embryos [see e.g., Carsience
15 et al. (1993) Development 117: 669-675]. The recipient chicken embryos within the shell are candled and allowed to hatch to yield a germline chimeric chicken.

DNA can be introduced into animal cells using any known procedure, including, but not limited to: direct uptake, incubation with
20 polyethylene glycol [PEG], microinjection, electroporation, lipofection, cell fusion, microcell fusion, particle bombardment, including microprojectile bombardment [see, e.g., U.S. Patent No. 5,470,708, which provides a method for transforming unattached mammalian cells via particle
25 bombardment], and any other such method. For example, the transfer of plasmid DNA in liposomes directly to human cells *in situ* has been approved by the FDA for use in humans [see, e.g., Nabel, et al. (1990) Science 249:1285-1288 and U.S. Patent No. 5,461,032].

-58-

b. Introduction of heterologous DNA into plants

Numerous methods for producing or developing transgenic plants are available to those of skill in the art. The method used is primarily a function of the species of plant. These methods include, but are not
5 limited to: direct transfer of DNA by processes, such as PEG-induced DNA uptake, protoplast fusion, microinjection, electroporation, and microprojectile bombardment [see, e.g., Uchimiya et al. (1989) J. of Biotech. 12: 1-20 for a review of such procedures, see, also, e.g., U.S. Patent Nos. 5,436,392 and 5,489,520 and many others]. For purposes
10 herein, when introducing a MAC, microinjection, protoplast fusion and particle gun bombardment are preferred.

Plant species, including tobacco, rice, maize, rye, soybean, Brassica napus, cotton, lettuce, potato and tomato, have been used to produce transgenic plants. Tobacco and other species, such as petunias,
15 often serve as experimental models in which the methods have been developed and the genes first introduced and expressed.

DNA uptake can be accomplished by DNA alone or in the presence of PEG, which is a fusion agent, with plant protoplasts or by any variations of such methods known to those of skill in the art [see, e.g.,
20 U.S. Patent No. 4,684,611 to Schilperroot et al.]. Electroporation, which involves high-voltage electrical pulses to a solution containing a mixture of protoplasts and foreign DNA to create reversible pores, has been used, for example, to successfully introduce foreign genes into rice and Brassica napus. Microinjection of DNA into plant cells, including cultured
25 cells and cells in intact plant organs and embryoids in tissue culture and microprojectile bombardment [acceleration of small high density particles, which contain the DNA, to high velocity with a particle gun apparatus, which forces the particles to penetrate plant cell walls and membranes] have also been used. All plant cells into which DNA can be introduced

-59-

and that can be regenerated from the transformed cells can be used to produce transformed whole plants which contain the transferred artificial chromosome. The particular protocol and means for introduction of the DNA into the plant host may need to be adapted or refined to suit the particular plant species or cultivar.

c. Insect cells

Insects are useful hosts for introduction of artificial chromosomes for numerous reasons, including, but not limited to: (a) amplification of genes encoding useful proteins can be accomplished in the artificial chromosome to obtain higher protein yields in insect cells; (b) insect cells support required post-translational modifications, such as glycosylation and phosphorylation, that can be required for protein biological functioning; (c) insect cells do not support mammalian viruses, and, thus, eliminate the problem of cross-contamination of products with such infectious agents; (d) this technology circumvents traditional recombinant baculovirus systems for production of nutritional, industrial or medicinal proteins in insect cell systems; (e) the low temperature optimum for insect cell growth (28° C) permits reduced energy cost of production; (f) serum-free growth medium for insect cells permits lower production costs; (g) artificial chromosome-containing cells can be stored indefinitely at low temperature; and (h) insect larvae will be biological factories for production of nutritional, medicinal or industrial proteins by microinjection of fertilized insect eggs [see, e.g., Joy et al. (1991) Current Science 66:145-150, which provides a method for microinjecting heterologous DNA into *Bombyx mori* eggs].

Either MACs or insect-specific artificial chromosomes [BUGACs] will be used to introduce genes into insects. As described in the Examples, it appears that MACs will function in insects to direct expression of heterologous DNA contained thereon. For example, as

-60-

described in the Examples, a MAC containing the *B. mori* actin gene promoter fused to the lacZ gene has been generated by transfection of EC3/7C5 cells with a plasmid containing the fusion gene. Subsequent fusion of the *B. mori* cells with the transfected EC3/7C5 cells that
5 survived selection yielded a MAC-containing insect-mouse hybrid cell line in which β -galactosidase expression was detectable.

Insect host cells include, but are not limited to, hosts such as *Spodoptera frugiperda* [caterpillar], *Aedes aegypti* [mosquito], *Aedes albopictus* [mosquito], *Drosophila melanogaster* [fruitfly], *Bombyx mori*
10 [silkworm], *Manduca sexta* [tomato horn worm] and *Trichoplusia ni* [cabbage looper]. Efforts have been directed toward propagation of insect cells in culture. Such efforts have focused on the fall armyworm, *Spodoptera frugiperda*. Cell lines have been developed also from other insects such as the cabbage looper, *Trichoplusia ni* and the silkworm,
15 *Bombyx mori*. It has also been suggested that analogous cell lines can be created using the tomato hornworm, *Manduca sexta*. To introduce DNA into an insect, it should be introduced into the larvae, and allowed to proliferate, and then the hemolymph recovered from the larvae so that the proteins can be isolated therefrom.

20 The preferred method herein for introduction of artificial chromosomes into insect cells is microinjection [see, e.g., Tamura et al. (1991) Bio Ind. 8:26-31; Nikolaev et al. (1989) Mol. Biol. (Moscow) 23:1177-87; and methods exemplified and discussed herein].

E. Applications for and Uses of Artificial chromosomes

25 Artificial chromosomes provide convenient and useful vectors, and in some instances [e.g., in the case of very large heterologous genes] the only vectors, for introduction of heterologous genes into hosts. Virtually any gene of interest is amenable to introduction into a host via artificial chromosomes. Such genes include, but are not limited to, genes that

-61-

encode receptors, cytokines, enzymes, proteases, hormones, growth factors, antibodies, tumor suppressor genes, therapeutic products and multigene pathways.

The artificial chromosomes provided herein will be used in methods of protein and gene product production, particularly using insects as host cells for production of such products, and in cellular (e.g., mammalian cell) production systems in which the artificial chromosomes (particularly MACs) provide a reliable, stable and efficient means for optimizing the biomanufacturing of important compounds for medicine and industry. They are also intended for use in methods of gene therapy, and for production of transgenic plants and animals [discussed above, below and in the EXAMPLES].

1. Gene Therapy

Any nucleic acid encoding a therapeutic gene product or product of a multigene pathway may be introduced into a host animal, such as a human, or into a target cell line for introduction into an animal, for therapeutic purposes. Such therapeutic purposes include, genetic therapy to cure or to provide gene products that are missing or defective, to deliver agents, such as anti-tumor agents, to targeted cells or to an animal, and to provide gene products that will confer resistance or reduce susceptibility to a pathogen or ameliorate symptoms of a disease or disorder. The following are some exemplary genes and gene products. Such exemplification is not intended to be limiting.

a. Anti-HIV ribozymes

As exemplified below, DNA encoding anti-HIV ribozymes can be introduced and expressed in cells using MACs, including the euchromatin-based minichromosomes and the SATACs. These MACs can be used to make a transgenic mouse that expresses a ribozyme and, thus, serves as a model for testing the activity of such ribozymes or from

-62-

which ribozyme-producing cell lines can be made. Also, introduction of a MAC that encodes an anti-HIV ribozyme into human cells will serve as treatment for HIV infection. Such systems further demonstrate the viability of using any disease-specific ribozyme to treat or ameliorate a particular disease.

b. Tumor Suppressor Genes

Tumor suppressor genes are genes that, in their wild-type alleles, express proteins that suppress abnormal cellular proliferation. When the gene coding for a tumor suppressor protein is mutated or deleted, the resulting mutant protein or the complete lack of tumor suppressor protein expression may result in a failure to correctly regulate cellular proliferation. Consequently, abnormal cellular proliferation may take place, particularly if there is already existing damage to the cellular regulatory mechanism. A number of well-studied human tumors and tumor cell lines have been shown to have missing or nonfunctional tumor suppressor genes.

Examples of tumor suppression genes include, but are not limited to, the retinoblastoma susceptibility gene or RB gene, the p53 gene, the gene that is deleted in colon carcinoma [*i.e.*, the DCC gene] and the neurofibromatosis type 1 [NF-1] tumor suppressor gene [see, *e.g.*, U.S. Patent No. 5,496,731; Weinberg *et al.* (1991) *Science* 254:1138-1146]. Loss of function or inactivation of tumor suppressor genes may play a central role in the initiation and/or progression of a significant number of human cancers.

25 The p53 Gene

Somatic cell mutations of the p53 gene are said to be the most frequent of the gene mutations associated with human cancer [see, *e.g.*, Weinberg *et al.* (1991) *Science* 254:1138-1146]. The normal or wild-type p53 gene is a negative regulator of cell growth, which, when

-63-

damaged, favors cell transformation. The p53 expression product is found in the nucleus, where it may act in parallel or cooperatively with other gene products. Tumor cell lines in which p53 has been deleted have been successfully treated with wild-type p53 vector to reduce
5 tumorigenicity [see, Baker *et al.* (1990) *Science* 249:912-915].

DNA encoding the p53 gene and plasmids containing this DNA are well known [see, *e.g.*, U.S. Patent No. 5,260,191; see, also Chen *et al.* (1990) *Science* 250:1576; Farrel *et al.* (1991) *EMBO J.* 10:2879-2887; plasmids containing the gene are available from the ATCC, and the
10 sequence is in the GenBank Database, accession nos. X54156, X60020, M14695, M16494, K03199].

c. The CFTR gene

Cystic fibrosis [CF] is an autosomal recessive disease that affects epithelia of the airways, sweat glands, pancreas, and other organs. It is
15 a lethal genetic disease associated with a defect in chloride ion transport, and is caused by mutations in the gene coding for the cystic fibrosis transmembrane conductance regulator [CFTR], a 1480 amino acid protein that has been associated with the expression of chloride conductance in a variety of eukaryotic cell types. Defects in CFTR destroy or reduce the
20 ability of epithelial cells in the airways, sweat glands, pancreas and other tissues to transport chloride ions in response to cAMP-mediated agonists and impair activation of apical membrane channels by cAMP-dependent protein kinase A [PKA]. Given the high incidence and devastating nature of this disease, development of effective CF treatments is imperative.

25 The CFTR gene [~ 250 kb] can be transferred into a MAC for use, for example, in gene therapy as follows. A CF-YAC [see Green *et al.* *Science* 250:94-98] may be modified to include a selectable marker, such as a gene encoding a protein that confers resistance to puromycin or hygromycin, and λ -DNA for use in site-specific integration into a neo-

-64-

minichromosome or a SATAC. Such a modified CF-YAC can be introduced into MAC-containing cells, such as EC3/7C5 or 19C5xHa4 cells, by fusion with yeast protoplasts harboring the modified CF-YAC or microinjection of yeast nuclei harboring the modified CF-YAC into the
5 cells. Stable transformants are then selected on the basis of antibiotic resistance. These transformants will carry the modified CF-YAC within the MAC contained in the cells.

2. Animals, birds, fish and plants that are genetically altered to possess desired traits such as resistance to disease

10 Artificial chromosomes are ideally suited for preparing animals, including vertebrates and invertebrates, including birds and fish as well as mammals, that possess certain desired traits, such as, for example, disease resistance, resistance to harsh environmental conditions, altered growth patterns, and enhanced physical characteristics.

15 One example of the use of artificial chromosomes in generating disease-resistant organisms involves the preparation of multivalent vaccines. Such vaccines include genes encoding multiple antigens that can be carried in a MAC, or species-specific artificial chromosome, and either delivered to a host to induce immunity, or incorporated into
20 embryos to produce transgenic (non-human) animals and plants that are immune or less susceptible to certain diseases.

Disease-resistant animals and plants may also be prepared in which resistance or decreased susceptibility to disease is conferred by introduction into the host organism or embryo of artificial chromosomes
25 containing DNA encoding gene products (e.g., ribozymes and prot ins that are toxic to certain pathogens) that destroy or attenuate pathogens or limit access of pathogens to the host.

Animals and plants possessing desired traits that might, for example, enhance utility, processibility and commercial value of the

-65-

organisms in areas such as the agricultural and ornamental plant industries may also be generated using artificial chromosomes in the same manner as described above for production of disease-resistant animals and plants. In such instances, the artificial chromosomes that
5 are introduced into the organism or embryo contain DNA encoding gene products that serve to confer the desired trait in the organism.

Birds, particularly fowl such as chickens, fish and crustaceans will serve as model hosts for production of genetically altered organisms using artificial chromosomes.

10 3. **Use of MACs and other artificial chromosomes for preparation and screening of libraries**

Since large fragments of DNA can be incorporated into each artificial chromosome, the chromosomes are well-suited for use as cloning vehicles that can accommodate entire genomes in the preparation
15 of genomic DNA libraries, which then can be readily screened. For example, MACs may be used to prepare a genomic DNA library useful in the identification and isolation of functional centromeric DNA from different species of organisms. In such applications, the MAC used to prepare a genomic DNA library from a particular organism is one that is
20 not functional in cells of that organism. That is, the MAC does not stably replicate, segregate or provide for expression of genes contained within it in cells of the organism. Preferably, the MACs contain an indicator gene (e.g., the lacZ gene encoding β -galactosidase or genes encoding products that confer resistance to antibiotics such as
25 neomycin, puromycin, hygromycin) linked to a promoter that is capable of promoting transcription of the indicator gene in cells of the organism. Fragments of genomic DNA from the organism are incorporated into the MACs, and the MACs are transferred to cells from the organism. Cells that contain MACs that have incorporated functional centromeres

-66-

contained within the genomic DNA fragments are identified by detection of expression of the marker gene.

4. Use of MACs and other artificial chromosomes for stable, high-level protein production

5 Cells containing the MACs and/or other artificial chromosomes provided herein are advantageously used for production of proteins, particularly several proteins from one cell line, such as multiple proteins involved in a biochemical pathway or multivalent vaccines. The genes encoding the proteins are introduced into the artificial chromosomes
10 which are then introduced into cells. Alternatively, the heterologous gene(s) of interest are transferred into a production cell line that already contains artificial chromosomes in a manner that targets the gene(s) to the artificial chromosomes. The cells are cultured under conditions whereby the heterologous proteins are expressed. Because the proteins
15 will be expressed at high levels in a stable permanent extra-genomic chromosomal system, selective conditions are not required.

Any transfectable cells capable of serving as recombinant hosts adaptable to continuous propagation in a cell culture system [see, e.g., McLean (1993) Trends In Biotech. 11:232-238] are suitable for use in an
20 artificial chromosome-based protein production system. Exemplary host cell lines include, but are not limited to, the following: Chinese hamster ovary (CHO) cells [see, e.g., Zang et al. (1995) Biotechnology 13:389-392], HEK 293, Ltk⁻, COS-7, DG44, and BHK cells. CHO cells are particularly preferred host cells. Selection of host cell lines for use in
25 artificial chromosome-based protein production systems is within the skill of the art, but often will depend on a variety of factors, including the properties of the heterologous protein to be produced, potential toxicity of the protein in the host cell, any requirements for post-translational modification (e.g., glycosylation, amination, phosphorylation) of the

-67-

protein, transcription factors available in the cells, the type of promoter element(s) being used to drive expression of the heterologous gene, whether production will be completely intracellular or the heterologous protein will preferably be secreted from the cell, and the types of
5 processing enzymes in the cell.

The artificial chromosome-based system for heterologous protein production has many advantageous features. For example, as described above, because the heterologous DNA is located in an independent, extra-genomic artificial chromosome (as opposed to randomly inserted in
10 an unknown area of the host cell genome or located as extrachromosomal element(s) providing only transient expression) it is stably maintained in an active transcription unit and is not subject to ejection via recombination or elimination during cell division. Accordingly, it is unnecessary to include a selection gene in the host
15 cells and thus growth under selective conditions is also unnecessary. Furthermore, because the artificial chromosomes are capable of incorporating large segments of DNA, multiple copies of the heterologous gene and linked promoter element(s) can be retained in the chromosomes, thereby providing for high-level expression of the foreign
20 protein(s). Alternatively, multiple copies of the gene can be linked to a single promoter element and several different genes may be linked in a fused polygene complex to a single promoter for expression of, for example, all the key proteins constituting a complete metabolic pathway [see, e.g., Beck von Bodman et al. (1995) Biotechnology 13:587-591].
25 Alternatively, multiple copies of a single gene can be operatively linked to a single promoter, or each or one or several copies may be linked to different promoters or multiple copies of the same promoter. Additionally, because artificial chromosomes have an almost unlimited capacity for integration and expression of foreign genes, they can be

-68-

used not only for the expression of genes encoding end-products of interest, but also for the expression of genes associated with optimal maintenance and metabolic management of the host cell, e.g., genes encoding growth factors, as well as genes that may facilitate rapid
5 synthesis of correct form of the desired heterologous protein product, e.g., genes encoding processing enzymes and transcription factors. The MACS are suitable for expression of any proteins or peptides, including proteins and peptides that require in vivo posttranslational modification for their biological activity. Such proteins include, but are
10 not limited to antibody fragments, full-length antibodies, and multimeric antibodies, tumor suppressor proteins, naturally occurring or artificial antibodies and enzymes, heat shock proteins, and others.

Thus, such cell-based "protein factories" employing MACs can generated using MACs constructed with multiple copies [theoretically an
15 unlimited number or at least up to a number such that the resulting MAC is about up to the size of a genomic chromosome (i.e., endogenous)] of protein-encoding genes with appropriate promoters, or multiple genes driven by a single promoter, i.e., a fused gene complex [such as a complete metabolic pathway in plant expression system; see, e.g., Beck
20 von Bodman (1995) Biotechnology 13:587-591]. Once such MAC is constructed, it can be transferred to a suitable cell culture system, such as a CHO cell line in protein-free culture medium [see, e.g., (1995) Biotechnology 13:389-39] or other immortalized cell lines [see, e.g., (1993) TIBTECH 11:232-238] where continuous production can be
25 established.

The ability of MACs to provide for high-level expression of heterologous proteins in host cells is demonstrated, for example, by analysis of the H1D3 and G3D5 cell lines described herein and deposited with the ECACC. Northern blot analysis of mRNA obtained from these

-69-

cells reveals that expression of the hygromycin-resistance and β -galactosidase genes in the cells correlates with the amplicon number of the megachromosome(s) contained therein.

5 F. Methods for the synthesis of DNA sequences containing repeated DNA units

Generally, assembly of tandemly repeated DNA poses difficulties such as unambiguous annealing of the complementary oligos. For example, separately annealed products may ligate in an inverted
10 orientation. Additionally, tandem or inverted repeats are particularly susceptible to recombination and deletion events that may disrupt the sequence. Selection of appropriate host organisms (e.g., rec⁻ strains) for use in the cloning steps of the synthesis of sequences of tandemly repeated DNA units may aid in reduction and elimination of such events.

15 Methods are provided herein for the synthesis of extended DNA sequences containing repeated DNA units. These methods are particularly applicable to the synthesis of arrays of tandemly repeated DNA units, which are generally difficult or not possible to construct utilizing other known gene assembly strategies. A specific use of these
20 methods is in the synthesis of sequences of any length containing simple (e.g., ranging from 2-6 nucleotides) tandem repeats (such as telomeres and satellite DNA repeats and trinucleotide repeats of possible clinical significance) as well as complex repeated DNA sequences. An particular example of the synthesis of a telomere sequence containing over 150
25 successive repeated hexamers utilizing these methods is provided herein.

The methods provided herein for synthesis of arrays of tandem DNA repeats are based in a series of extension steps in which successive doublings of a sequence of repeats results in an exponential expansion of the array of tandem repeats. These methods provide several advantages
30 over previously known methods of gene assembly. For instance, the

-70-

starting oligonucleotides are used only once. The intermediates in, as well as the final product of, the construction of the DNA arrays described herein may be obtained in cloned form in a microbial organism (e.g., *E. coli* and yeast). Of particular significance, with regard to these methods

5 is the fact that sequence length increases exponentially, as opposed to linearly, in each extension step of the procedure even though only two oligonucleotides are required in the methods. The construction process does not depend on the compatibility of restriction enzyme recognition sequences and the sequence of the repeated DNA because restriction

10 sites are used only temporarily during the assembly procedure. No adaptor is necessary, though a region of similar function is located between two of the restriction sites employed in the process. The only limitation with respect to restriction site use is that the two sites employed in the method must not be present elsewhere in the vector

15 utilized in any cloning steps. These procedures can also be used to construct complex repeats with perfectly identical repeat units, such as the variable number tandem repeat (VNTR) 3' of the human apolipoprotein B100 gene (a repeat unit of 30 bp, 100% AT) or alphoid satellite DNA.

20 The method of synthesizing DNA sequences containing tandem repeats may generally be described as follows.

1. Starting materials

Two oligonucleotides are utilized as starting materials. Oligonucleotide 1 is of length k of repeated sequence (the flanks of which are not relevant) and contains a relatively short stretch (60-90 nucleotides) of the repeated sequence, flanked with appropriately chosen restriction sites:

[illegible]

-71-

wherein S1 is restriction site 1 cleaved by E1 [preferably an enzyme producing a 3'-overhang (e.g., PacI, PstI, SphI, NsiI, etc.) or blunt-end], S2 is a second restriction site cleaved by E2 (preferably an enzyme producing a 3'-overhang or one that cleaves outside the recognition sequence, such as TspRI), > represents a simple repeat unit, and '___' denotes a short (8-10) nucleotide flanking sequence complementary to oligonucleotide 2:

3'-_____S3-5'

wherein S3 is a third restriction site for enzyme E3 and which is present in the vector to be used during the construction.

Because there is a large variety of restriction enzymes that recognize many different DNA sequences as cleavage sites, it should always be possible to select sites and enzymes (preferably those that yield a 3'-protruding end) suitable for these methods in connection with the synthesis of any one particular repeat array. In most cases, only 1 (or perhaps 2) nucleotide(s) has of a restriction site is required to be present in the repeat sequence, and the remaining nucleotides of the restriction site can be removed, for example:

PacI: TTAAT/TAA-- (Klenow/dNTP) TAA--
PstI: CTGCA/G-- (Klenow/dNTP) G--
NsiI: ATGCA/T-- (Klenow/dNTP) T--
KpnI: GGTAC/C-- (Klenow/dNTP) C--

Though there is no known restriction enzyme leaving a single A behind, this problem can be solved with enzymes leaving behind none at all, for example:

TaiI: ACGT/ (Klenow/dNTP) --
NlaIII: CATG/ (Klenow/dNTP) --

Additionally, if mung bean nuclease is used instead of Klenow, then the following

-72-

XbaI: T/CTAGA Mung bean nuclease A--

Furthermore, there are a number of restriction enzymes that cut outside of the recognition sequence, and in this case, there is no limitation at all:

5 TspRI NNCAGTGNN/-- (Klenow/dNTP) --

BsmI GAATG CN/-- (Klenow/dNTP) --

CTTAC/GN -- (Klenow/dNTP) --

2. Step 1 - Annealing

Oligonucleotides 1 and 2 are annealed at a temperature selected
10 depending on the length of overlap (typically in the range of 30-65 °C).

3. Step 2 - Generating a double-stranded molecule

The annealed oligonucleotides are filled-in with Klenow polymerase in the presence of dNTP to produce a double-stranded (ds) sequence:

[illegible][illegible]

4. Step 3 - Incorporation of double-stranded DNA into a vector

The double-stranded DNA is cleaved with restriction enzymes E1 and E3 and subsequently ligated into a vector (e.g., pUC19 or a yeast vector) that has been cleaved with the same enzymes E1 and E3. The ligation product is used to transform competent host cells compatible with the vector being used (e.g., when pUC19 is used, bacterial cells such as E. coli DH5 α are suitable hosts) which are then plated onto selection plates. Recombinants can be identified either by color (e.g., by X-gal staining for β -galactosidase expression) or by colony hybridization using ³²P-labeled oligonucleotide 2 (detection by hybridization to oligonucleotide 2 is preferred because its sequence is removed in each of the subsequent extension steps and thus is present only in recombinants that contain DNA that has undergone successful extension of the repeated sequence).

-73-

5. Step 4 - Isolation of insert from the plasmid

An aliquot of the recombinant plasmid containing k nucleotides of the repeat sequence is digested with restriction enzymes E1 and E3, and the insert is isolated on a gel (native polyacrylamide while the insert is short, but agarose can be used for isolation of longer inserts in subsequent steps). A second aliquot of the recombinant plasmid is cut with enzymes E2 (treated with Klenow and dNTP to remove the 3'-overhang) and E3, and the large fragment (plasmid DNA plus the insert) is isolated.

10 6. Step 5 - Extension of the DNA sequence of k repeats

The two DNAs (the S1-S3 insert fragment and the vector plus insert) are ligated, plated to selective plates, and screened for extended recombinants as in Step 3. Now the length of the repeat sequence between restriction sites is twice that of the repeat sequence in the previous step, i.e., 2xk.

15 7. Step 6 - Extension of the DNA sequence of 2xk repeats

Steps 4 and 5 are repeated as many times as needed to achieve the desired repeat sequence size. In each extension cycle, the repeat sequence size doubles, i.e., if m is the number of extension cycles, the size of the repeat sequence will be $k \times 2^m$ nucleotides.

The following examples are included for illustrative purposes only and are not intended to limit the scope of the invention.

EXAMPLE 1**General Materials and Methods**

25 The following materials and methods are exemplary of methods that are used in the following Examples and that can be used to prepare cell lines containing artificial chromosomes. Other suitable materials and methods known to those of skill in the art may be used. Modifications of

-74-

these materials and methods known to those of skill in the art may also be employed.

A. Culture of cell lines, cell fusion, and transfection of cells

1. Chinese hamster K-20 cells and mouse A9 fibroblast
5 cells were cultured in F-12 medium. EC3/7 [see, U.S. Patent No. 5,288,625, and deposited at the European Collection of Animal cell Culture (ECACC) under accession no. 90051001; see, also Hadlaczky et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110 and U.S. application Serial No. 08/375,271] and EC3/7C5 [see, U.S. Patent No.
10 5,288,625 and Praznovszky et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:11042-11046] mouse cell lines, and the KE1-2/4 hybrid cell line were maintained in F-12 medium containing 400 μ g/ml G418 [SIGMA, St. Louis, MO].
2. TF1004G19 and TF1004G-19C5 mouse cells,
15 described below, and the 19C5xHa4 hybrid, described below, and its sublines were cultured in F-12 medium containing up to 400 μ g/ml Hygromycin B [Calbiochem]. LP11 cells were maintained in F-12 medium containing 3-15 μ g/ml Puromycin [SIGMA, St. Louis, MO].
3. Cotransfection of EC3/7C5 cells with plasmids
20 [pH132, pCH110 available from Pharmacia, see, also Hall et al. (1983) J. Mol. Appl. Gen. 2:101-109] and with λ DNA was conducted using the calcium phosphate DNA precipitation method [see, e.g., Chen et al. (1987) Mol. Cell. Biol. 7:2745-2752], using 2-5 μ g plasmid DNA and 20 μ g λ phage DNA per 5×10^6 recipient cells.

25 **4. Cell fusion**

Mouse and hamster cells were fused using polyethylene glycol [Davidson et al. (1976) Som. Cell Genet. 2:165-176]. Hybrid cells were selected in HAT medium containing 400 μ g/ml Hygromycin B.

-75-

Approximately 2×10^7 recipient and 2×10^6 donor cells were fused using polyethylene glycol [Davidson et al. (1976) Som. Cell Genet. 2:165-176]. Hybrids were selected and maintained in F-12/HAT medium [Szybalsky et al. (1962) Natl. Cancer Inst. Monogr. 7:75-89] containing
5 10% FCS and 400 $\mu\text{g/ml}$ G418. The presence of "parental" chromosomes in the hybrid cell lines was verified by *in situ* hybridization with species-specific probes using biotin-labeled human and hamster genomic DNA, and a mouse long interspersed repetitive DNA [pMCPE1.51].

10 5. Microcell fusion

Microcell-mediated transfer of artificial chromosomes from EC3/7C5 cells to recipient cells was done according to Saxon et al. [(1985) Mol. Cell. Biol. 1:140-146] with the modifications of Goodfellow et al. [(1989) Techniques for mammalian genome transfer. In *Genome Analysis a Practical Approach*. K.E. Davies, ed., IRL Press, Oxford,
15 Washington DC. pp.1-17] and Yamada et al. [(1990) Oncogene 5:1141-1147]. Briefly, 5×10^6 EC3/7C5 cells in a T25 flask were treated first with 0.05 $\mu\text{g/ml}$ colcemid for 48 hr and then with 10 $\mu\text{g/ml}$ cytochalasin B for 30 min. The T25 flasks were centrifuged on edge and the pelleted
20 microcells were suspended in serum free DME medium. The microcells were filtered through first a 5 micron and then a 3 micron polycarbonate filter, treated with 50 $\mu\text{g/ml}$ of phytohemagglutinin, and used for polyethylene glycol mediated fusion with recipient cells. Selection of cells containing the MMCneo was started 48 hours after fusion in
25 medium containing 400-800 $\mu\text{g/ml}$ G418.

Microcells were also prepared from 1B3 and GHB42 donor cells as follows in order to be fused with E2D6K cells [a CHO K-20 cell line carrying the puromycin N-acetyltransferase gene, i.e., the puromycin resistance gene, under the control of the SV40 early promoter]. The

-76-

donor cells were seeded to achieve 60-75% confluency within 24-36 hours. After that time, the cells were arrested in mitosis by exposure to colchicine (10 μ g/ml) for 12 or 24 hours to induce micronucleation. To promote micronucleation of GHB42 cells, the cells were exposed to
5 hypotonic treatment (10 min at 37°C). After colchicine treatment, or after colchicine and hypotonic treatment, the cells were grown in colchicine-free medium.

The donor cells were trypsinized and centrifuged and the pellets were suspended in a 1:1 Percoll medium and incubated for 30-40 min at
10 37°C. After the incubation, $1-3 \times 10^7$ cells (60-70% micronucleation index) were loaded onto each Percoll gradient (each fusion was distributed on 1-2 gradients). The gradients were centrifuged at 19,000 rpm for 80 min in a Sorvall SS-34 rotor at 34-37°C. After centrifugation, two visible bands of cells were removed, centrifuged at
15 2000 rpm, 10 min at 4°C, resuspended and filtered through 8 μ m pore size nucleopore filters.

The microcells prepared from the 1B3 and GHB42 cells were fused with E2D6K. The E2D6K cells were generated by CaPO_4 transfection of CHO K-20 cells with pCHTV2. Plasmid pCHTV2 contains the puromycin-
20 resistance gene linked to the SV40 promoter and polyadenylation signal, the Saccharomyces cerevisiae URA3 gene, 2.4- and 3.2-kb fragments of a Chinese hamster chromosome 2-specific satellite DNA (HC-2 satellite; see Fatyol *et al.* (1994) Nuc. Acids Res. 22:3728-3736), two copies of the diphtheria toxin-A chain gene (one linked to the herpes simplex virus
25 thymidine kinase (HSV-TK) gene promoter and SV40 polyadenylation signal and the other linked to the HSV-TK promoter without a polyadenylation signal), the ampicillin-resistance gene and the ColE1 origin of replication. Following transfection, puromycin-resistant colonies were isolated. The presence of the pCHTV2 plasmid in the E2D6K cell

-77-

line was confirmed by nucleic acid amplification of DNA isolated from the cells.

The purified microcells were centrifuged as described above and resuspended in 2 ml of phytohemagglutinin-P (PHA-P, 100 μ g/ml). The
5 microcell suspension was then added to a 60-70% confluent recipient culture of E2D6K cells. The preparation was incubated at room temperature for 30-40 min to agglutinate the microcells. After the PHA-P was removed, the cells were incubated with 1 ml of 50% polyethylene-glycol (PEG) for one min. The PEG was removed and the culture was
10 washed three times with F-12 medium without serum. The cells were incubated in non-selective medium for 48-60 hours. After this time, the cell culture was trypsinized and plated in F-12 medium containing 400 μ g/ml hygromycin B and 10 g/ml puromycin to select against the parental cell lines.

15 Hybrid clones were isolated from the cells that had been cultured in selective medium. These clones were then analyzed for expression of β -galactosidase by the X-gal staining method. Four of five hybrid clones analyzed that had been generated by fusion of GHB42 microcells with E2D6K cells yielded positive staining results indicating expression of β -
20 galactosidase from the lacZ gene contained in the megachromosome contributed by the GHB42 cells. Similarly, a hybrid clone that had been generated by fusion of 1B3 microcells with E2D6K cells yielded positive staining results indicating expression of β -galactosidase from the lacZ gene contained in the megachromosome contributed by the 1B3 cells. *In*
25 *situ* hybridization analysis of the hybrid clones is also performed to analyze the mouse chromosome content of the mouse-hamster hybrid cells.

-78-

B. Chromosome banding

Trypsin G-banding of chromosomes was performed using the method of Wang & Fedoroff [(1972) Nature 235:52-54], and the detection of constitutive heterochromatin with the BSG. C-banding method was done according to Sumner [(1972) Exp. Cell Res. 75:304-306]. For the detection of chromosome replication by bromodeoxyuridine [BrdU] incorporation, the Fluorescein Plus Giemsa [FPG] staining method of Perry & Wolff [(1974) Nature 251:156-158] was used.

10 C. Immunolabelling of chromosomes and *in situ* hybridization

Indirect immunofluorescence labelling with human anti-centromere serum LU851 [Hadlaczky et al. (1986) Exp. Cell Res. 167:1-15], and indirect immunofluorescence and *in situ* hybridization on the same preparation were performed as described previously [see, Hadlaczky et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110, see, also U.S. application Serial No. 08/375,271]. Immunolabelling with fluorescein-conjugated anti-BrdU monoclonal antibody [Boehringer] was performed according to the procedure recommended by the manufacturer, except that for treatment of mouse A9 chromosomes, 2 M hydrochloric acid was used at 37° C for 25 min, and for chromosomes of hybrid cells, 1 M hydrochloric acid was used at 37° C for 30 min.

D. Scanning electron microscopy

Preparation of mitotic chromosomes for scanning electron microscopy using osmium impregnation was performed as described previously [Sumner (1991) Chromosoma 100:410-418]. The chromosomes were observed with a Hitachi S-800 field emission scanning electron microscope operated with an accelerating voltage of 25 kV.

-79-

E. DNA manipulations, plasmids and probes

1. General methods

- All general DNA manipulations were performed by standard procedures [see, e.g., Sambrook et al. (1989) *Molecular cloning: A Laboratory Manual* Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY]. The mouse major satellite probe was provided by Dr. J. B. Rattner [University of Calgary, Alberta, Canada]. Cloned mouse satellite DNA probes [see Wong et al. (1988) *Nucl. Acids Res.* 16:11645-11661], including the mouse major satellite probe, were gifts from Dr. J. B. Rattner, University of Calgary. Hamster chromosome painting was done with total hamster genomic DNA, and a cloned repetitive sequence specific to the centromeric region of chromosome 2 [Fátyol et al. (1994) *Nucl. Acids Res.* 22:3728-3736] was also used. Mouse chromosome painting was done with a cloned long interspersed repetitive sequence [pMCP1.51] specific for the mouse euchromatin.

For cotransfection and for *in situ* hybridization, the pCH110 β -galactosidase construct [Pharmacia or Invitrogen], and λ cl 875 Sam7 phage DNA [New England Biolabs] were used.

2. Construction of Plasmid pPuroTel

- Plasmid pPuroTel, which carries a Puromycin-resistance gene and a cloned 2.5 kb human telomeric sequence [see SEQ ID No. 3], was constructed from the pBabe-puro retroviral vector [Morgenstern et al. (1990) *Nucl. Acids Res.* 18:3587-3596; provided by Dr. L. Székely (Microbiology and Tumorbiology Center, Karolinska Institutet, Stockholm); see, also Tonghua et al. (1995) *Chin. Med. J.* (Beijing, Engl. Ed.) 108:653-659; Couto et al. (1994) *Infect. Immun.* 62:2375-2378; Dunckley et al. (1992) *FEBS Lett.* 296:128-34; French et al. (1995) *Anal. Biochem.* 228:354-355; Liu et al. (1995) *Blood* 85:1095-1103;

-80-

International PCT application Nos. WO 9520044; WO 9500178, and WO 9419456].

F. Deposited cell lines

Cell lines KE1-2/4, EC3/7C5, TF1004G19C5, 19C5xHa4, G3D5
5 and H1D3 have been deposited in accord with the Budapest Treaty at the European Collection of Animal Cell Culture (ECACC) under Accession Nos. 96040924, 96040925, 96040926, 96040927, 96040928 and 96040929, respectively. The cell lines were deposited on April 9, 1996, at the European Collection of Animal Cell Cultures (ECACC) Vaccine
10 Research and Production Laboratory, Public Health Laboratory Service, Centre for Applied Microbiology and Research, Porton Down, Salisbury, Wiltshire SP4 0JG, United Kingdom. The deposits were made in the name of Gyula Hadlaczky of H. 6723, SZEGED, SZAMOS U.1.A. IX. 36. HUNGARY, who has authorized reference to the deposited cell lines in
15 this application and who has provided unreserved and irrevocable consent to the deposited cell lines being made available to the public in accordance with Rule 28(1)(d) of the European Patent Convention.

EXAMPLE 2

Preparation of EC3/7, EC3/7C5 and related cell lines

20 The EC3/7 cell line is an LMTK⁻ mouse cell line that contains the neo-centromere. The EC3/7C5 cell line is a single-cell subclone of EC3/7 that contains the neo-minichromosome.

A. EC3/7 Cell line

As described in U.S. Patent No. 5,288,625 [see, also Praznovszky
25 et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:11042-11046 and Hadlaczky et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110] *de novo* centromere formation occurs in a transformed mouse LMTK⁻ fibroblast cell line [EC3/7] after cointegration of λ constructs [λ CM8 and λ gtWESneo] carrying human and bacterial DNA.

-81-

By cotransfection of a 14 kb human DNA fragment cloned in λ [ACM8] and a dominant marker gene [λ gtWESneo], a selectable centromere linked to a dominant marker gene [neo-centromere] was formed in mouse LMTK cell line EC3/7 [Hadlaczky *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110, see Figure 1]. Integration of the heterologous DNA [the λ DNA and marker gene-encoding DNA] occurred into the short arm of an acrocentric chromosome [chromosome 7 (see, Figure 1B)], where an amplification process resulted in the formation of the new centromere [neo-centromere (see Figure 1C)]. On the dicentric chromosome (Figure 1C), the newly formed centromere region contains all the heterologous DNA (human, λ , and bacterial) introduced into the cell and an active centromere.

Having two functionally active centromeres on the same chromosome causes regular breakages between the centromeres [see, Figure 1E]. The distance between the two centromeres on the dicentric chromosome is estimated to be ~10-15 Mb, and the breakage that separates the minichromosome occurred between the two centromeres. Such specific chromosome breakages result in the appearance [in approximately 10% of the cells] of a chromosome fragment that carries the neo-centromere [Figure 1F]. This chromosome fragment is principally composed of human, λ , plasmid, and neomycin-resistance gene DNA, but it also has some mouse chromosomal DNA. Cytological evidence suggests that during the stabilization of the MMCneo, there was an inverted duplication of the chromosome fragment bearing the neo-centromere. The size of minichromosomes in cell lines containing the MMCneo is approximately 20-30 Mb; this finding indicates a two-fold increase in size.

From the EC3/7 cell line, which contains the dicentric chromosome [Figure 1E], two sublines [EC3/7C5 and EC3/7C6] were selected by

-82-

repeated single-cell cloning. In these cell lines, the neo-centromere was found exclusively on a small chromosome [neo-minichromosome], while the formerly dicentric chromosome carried detectable amounts of the exogenously-derived DNA sequences but not an active neo-centromere
5 [Figure 1F and 1G].

The minichromosomes of cell lines EC3/7C5 and EC3/7C6 are similar. No differences are detected in their architectures at either the cytological or molecular level. The minichromosomes were indistinguishable by conventional restriction endonuclease mapping or by
10 long-range mapping using pulsed field electrophoresis and Southern hybridization. The cytoskeleton of cells of the EC3/7C6 line showed an increased sensitivity to colchicine, so the EC3/7C5 line was used for further detailed analysis.

B. Preparation of the EC3/7C5 and EC3/7C6 cell lines

15 The EC3/7C5 cells, which contain the neo-minichromosome, were produced by subcloning the EC3/7 cell line in high concentrations of G418 [40-fold the lethal dose] for 350 generations. Two single cell-derived stable cell lines [EC3/7C5 and EC3/7C6] were established. These cell lines carry the neo-centromere on minichromosomes and also
20 contain the remaining fragment of the dicentric chromosome. Indirect immunofluorescence with anti-centromere antibodies and subsequent in situ hybridization experiments demonstrated that the minichromosomes derived from the dicentric chromosome. In EC3/7C5 and EC3/7C6 cell lines (140 and 128 metaphases, respectively) no intact dicentric
25 chromosomes were found, and minichromosomes were detected in 97.2% and 98.1% of the cells, respectively. The minichromosomes have been maintained for over 150 cell generations. They do contain the remaining portion of the formerly dicentric chromosome.

-83-

Multiple copies of telomeric DNA sequences were detected in the marker centromeric region of the remaining portion of the formerly dicentric chromosome by in situ hybridization. This indicates that mouse telomeric sequences were coamplified with the foreign DNA sequences.

- 5 These stable minichromosome-carrying cell lines provide direct evidence that the extra centromere is functioning and is capable of maintaining the minichromosomes [see, U.S. Patent No. 5,288,625].

- The chromosome breakage in the EC3/7 cells, which separates the neo-centromere from the mouse chromosome, occurred in the G-band
10 positive "foreign" DNA region. This is supported by the observation of traces of λ and human DNA sequences at the broken end of the formerly dicentric chromosome. Comparing the G-band pattern of the chromosome fragment carrying the neo-centromere with that of the stable neo-minichromosome, reveals that the neo-minichromosome is an
15 inverted duplicate of the chromosome fragment that bears the neo-centromere. This is also evidenced by the observation that although the neo-minichromosome carries only one functional centromere, both ends of the minichromosome are heterochromatic, and mouse satellite DNA sequences were found in these heterochromatic regions by *in situ*
20 hybridization.

- These two cell lines, EC3/7C5 and EC3/7C6, thus carry a selectable mammalian minichromosome [MMCneo] with a centromere linked to a dominant marker gene [Hadlaczký et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110]. MMCneo is intended to be used as a
25 vector for minichromosome-mediated gene transfer and has been used as mod 1 of a minichromosome-based vector system.

Long range mapping studies of the MMCneo indicated that human DNA and the neomycin-resistance gene constructs integrated into the mouse chromosome separately, followed by the amplification of the

-84-

chromosome region that contains the exogenous DNA. The MMCneo contains about 30-50 copies of the λ CM8 and λ gtWESneo DNA in the form of approximately 160 kb repeated blocks, which together cover at least a 3.5 Mb region. In addition to these, there are mouse telomeric sequences [Praznovszky et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:11042-11046] and any DNA of mouse origin necessary for the correct higher-ordered structural organization of chromatids.

Using a chromosome painting probe mCPE1.51 [mouse long interspersed repeated DNA], which recognizes exclusively euchromatic mouse DNA, detectable amounts of interspersed repeat sequences were found on the MMCneo by *in situ* hybridization. The neo-centromere is associated with a small but detectable amount of satellite DNA. The chromosome breakage that separates the neo-centromere from the mouse chromosome occurs in the "foreign" DNA region. This is demonstrated by the presence of λ and human DNA at the broken end of the formerly dicentric chromosome. At both ends of the MMCneo, however, there are traces of mouse major satellite DNA as evidenced by *in situ* hybridization. This observation suggests that the doubling in size of the chromosome fragment carrying the neo-centromere during the stabilization of the MMCneo is a result of an inverted duplication. Although mouse telomere sequences, which coamplified with the exogenous DNA sequences during the neo-centromere formation, may provide sufficient telomeres for the MMCneo, the duplication could have supplied the functional telomeres for the minichromosome.

The nucleotide sequence of portions of the neo-minichromosomes was determined as follows. Total DNA was isolated from EC3/7C5 cells according to standard procedures. The DNA was subjected to nucleic acid amplification using the Expand Long Template PCR system [Boehringer Mannheim] according to the manufacturer's procedures. The

-85-

amplification procedure required only a single 33-mer oligonucleotide primer corresponding to sequence in a region of the phage λ right arm, which is contained in the neo-minichromosome. The sequence of this oligonucleotide is set forth as the first 33 nucleotides of SEQ ID No. 13.

- 5 Because the neo-minichromosome contains a series of inverted repeats of this sequence, the single oligonucleotide was used as a forward and reverse primer resulting in amplification of DNA positioned between sets of inverted repeats of the phage λ DNA. Three products were obtained from the single amplification reaction, which suggests that the sequence
- 10 of the DNA located between different sets of inverted repeats may differ. In a repeating nucleic acid unit within an artificial chromosome, minor differences may be present and may occur during culturing of cells containing the artificial chromosome. For example, base pair changes may occur as well as integration of mobile genetic elements and
- 15 deletions of repeated sequences.

- Each of the three products was subjected to DNA sequence analysis. The sequences of the three products are set forth in SEQ ID Nos. 13, 14, and 15, respectively. To be certain that the sequenced products were amplified from the neo-minichromosome, control
- 20 amplifications were conducted using the same primers on DNA isolated from negative control cell lines (mouse Ltk⁺ cells) lacking minichromosomes and the formerly dicentric chromosome, and positive control cell lines [the mouse-hamster hybrid cell line GB43 generated by treating 19C5xHa4 cells (see Figure 4) with BrdU followed by growth in G418-containing selective medium and retreatment with BrdU] containing a neo-minichromosome only. Only the positive control cell line yielded three amplification products; no amplification product was detected in the negative control reaction. The results obtained in the positive amplification also demonstrate that the neo-minichromosome

-86-

DNA, and not the fragment of the formerly dicentric mouse chromosome, was amplified.

The sequences of the three amplification products were compared to those contained in the Genbank/EMBL database. SEQ ID Nos. 13 and 14 showed high (~96%) homology to portions of DNA from intracisternal A-particles from mouse. SEQ ID No. 15 showed no significant homology with sequences available in the database. All three of these sequences may be used for generating gene targeting vectors as homologous DNAs to the neo-minichromosome.

10 **C. Isolation and partial purification of minichromosomes**

Mitotic chromosomes of EC3/7C5 cells were isolated as described by Hadlaczky et al. [(1981) Chromosoma 81:537-555], using a glycine-hexylene glycol buffer system [Hadlaczky et al. (1982) Chromosoma 86:643-659]. Chromosome suspensions were centrifuged at 1,200 x g for 30 minutes. The supernatant containing minichromosomes was centrifuged at 5,000 x g for 30 minutes and the pellet was resuspended in the appropriate buffer. Partially purified minichromosomes were stored in 50% glycerol at -20° C.

20 **D. Stability of the MMCneo maintenance and *neo* expression**

EC3/7C5 cells grown in non-selective medium for 284 days and then transferred to selective medium containing 400 µg/ml G418 showed a 96% plating efficiency (colony formation) compared to control cells cultured permanently in the presence of G418. Cytogenetic analysis indicated that the MMCneo is stably maintained at one copy per cell under selective and non-selective culture conditions. Only two metaphases with two MMCneo were found in 2,270 metaphases analyzed.

Southern hybridization analysis showed no detectable changes in DNA restriction patterns, and similar hybridization intensities were

-87-

observed with a *neo* probe when DNA from cells grown under selective or non-selective culture conditions were compared.

Northern analysis of RNA transcripts from the *neo* gene isolated from cells grown under selective and non-selective conditions showed only minor and not significant differences. Expression of the *neo* gene persisted in EC3/7C5 cells maintained in F-12 medium free of G418 for 290 days under non-selective culture conditions. The long-term expression of the *neo* gene(s) from the minichromosome may be influenced by the nuclear location of the MMCneo. *In situ* hybridization experiments revealed a preferential peripheral location of the MMCneo in the interphase nucleus. In more than 60% of the 2,500 nuclei analyses, the minichromosome was observed at the perimeter of the nucleus near the nuclear envelope.

EXAMPLE 3

15 Minichromosome transfer and production of the λ -neo-chromosome

A. Minichromosome transfer

The neo-minichromosome [referred to as MMCneo, FIG. 2C] has been used for gene transfer by fusion of minichromosome-containing cells [EC3/7C5 or EC3/7C6] with different mammalian cells, including hamster and human. Thirty-seven stable hybrid cell lines have been produced. All established hybrid cell lines proved to be true hybrids as evidenced by *in situ* hybridization using biotinylated human, and hamster genomic, or pMCPE1.51 mouse long interspersed repeated DNA probes for "chromosome painting". The MMCneo has also been successfully transferred into mouse A9, L929 and pluripotent F9 teratocarcinoma cells by fusion of microcells derived from EC3/7C5 cells. Transfer was confirmed by PCR, Southern blotting and *in situ* hybridization with minichromosome-specific probes. The cytogenetic analysis confirmed

-88-

that, as expected for microcell fusion, a few cells [1-5%] received [or retained] the MMCneo.

These results demonstrate that the MMCneo is tolerated by a wide range of cells. The prokaryotic genes and the extra dosage for the
5 human and λ sequences carried on the minichromosome seem to be not disadvantageous for tissue culture cells.

The MMCneo is the smallest chromosome of the EC3/7C5 genome and is estimated to be approximately 20-30 Mb, which is significantly smaller than the majority of the host cell (mouse) chromosomes. By
10 virtue of the smaller size, minichromosomes can be partially purified from a suspension of isolated chromosomes by a simple differential centrifugation. In this way, minichromosome suspensions of 15-20% purity have been prepared. These enriched minichromosome
15 preparations can be used to introduce, such as by microinjection or lipofection, the minichromosome into selected target cells. Target cells include therapeutic cells that can be use in methods of gene therapy, and also embryonic cells for the preparation of transgenic (non-human) animals.

The MMCneo is capable of autonomous replication, is stably
20 maintained in cells, and permits persistent expression of the *neo* gene(s), even after long-term culturing under non-selective conditions. It is a non-integrative vector that appears to occupy a territory near the nuclear envelope. Its peripheral localization in the nucleus may have an important role in maintaining the functional integrity and stability of the
25 MMCneo. Functional compartmentalization of the host nucleus may have an effect on the function of foreign sequences. In addition, MMCneo contains megabases of λ DNA sequences that should serve as a target site for homologous recombination and thus integration of desired gene(s) into the MMCneo. It can be transferred by cell and microcell

-89-

fusion, microinjection, electroporation, lipid-mediated carrier systems or chromosome uptake. The neo-centromere of the MMCneo is capable of maintaining and supporting the normal segregation of a larger 150-200 Mb λ neo-chromosome. This result demonstrates that the MMCneo chromosome should be useful for carrying large fragments of heterologous DNA.

B. Production of the λ neo-chromosome

In the hybrid cell line KE1-2/4 made by fusion of EC3/7 and Chinese hamster ovary cells [FIG 2], the separation of the neo-centromere from the dicentric chromosome was associated with a further amplification process. This amplification resulted in the formation of a stable chromosome of average size [i.e., the λ neo-chromosome; see, Praznovszky *et al.* (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88:11042-11046]. The λ neo-chromosome carries a terminally located functional centromere and is composed of seven large amplicons containing multiple copies of λ , human, bacterial, and mouse DNA sequences [see FIG 2]. The amplicons are separated by mouse major satellite DNA [Praznovszky *et al.* (1991) *Proc. Natl. Acad. Sci. U.S.A.* 88:11042-11046] which forms narrow bands of constitutive heterochromatin between the amplicons.

EXAMPLE 4

Formation of the "sausage chromosome" [SC]

The findings set forth in the above EXAMPLES demonstrate that the centromeric region of the mouse chromosome 7 has the capacity for large-scale amplification [other results indicate that this capacity is not unique to chromosome 7]. This conclusion is further supported by results from cotransfection experiments, in which a second dominant selectable marker gene and a non-selected marker gene were introduced into EC3/7C5 cells carrying the formerly dicentric chromosome 7 and the

-90-

neo-minichromosome. The EC3/7C5 cell line was transformed with λ phage DNA, a hygromycin-resistance gene construct [pH132], and a β -galactosidase gene construct [pCH110]. Stable transformants were selected in the presence of high concentrations [400 μ g/ml] Hygromycin B, and analyzed by Southern hybridization. Established transformant cell lines showing multiple copies of integrated exogenous DNA were studied by *in situ* hybridization to localize the integration site(s), and by LacZ staining to detect β -galactosidase expression.

A. Materials and methods

10 1. Construction of pH132

The pH132 plasmid carries the hygromycin B resistance gene and the anti-HIV-1 *gag* ribozyme [see, SEQ ID NO. 6 for DNA sequence that corresponds to the sequence of the ribozyme] under control of the β -actin promoter. This plasmid was constructed from pHyg plasmid [Sugden *et al.* (1985) Mol. Cell. Biol. 5:410-413; a gift from Dr. A. D. Riggs, Beckman Research Institute, Duarte; see, also, *e.g.*, U.S. Patent No. 4,997,764], and from pPC-RAG12 plasmid [see, Chang *et al.* (1990) Clin Biotech 2:23-31; provided by Dr. J. J. Rossi, Beckman Research Institute, Duarte; see, also U.S. Patent Nos. 5,272,262, 5,149,796 and 5,144,019, which describes the anti-HIV *gag* ribozyme and construction of a mammalian expression vector containing the ribozyme insert linked to the β -actin promoter and SV40 late gene transcriptional termination and polyA signals]. Construction of pPC-RAG12 involved insertion of the ribozyme insert flanked by BamHI linkers was into BamHI-digested pH β -Apr-1gpt [see, Gunning *et al.* (1987) Proc. Natl. Acad. Sci. U.S.A. 84:4831-4835, see, also U.S. Patent No. 5,144,019].

Plasmid pH132 was constructed as follows. First, pPC-RAG12 [described by Chang *et al.* (1990) Clin. Biotech. 2:23-31] was digested with BamHI to excise a fragment containing an anti-HIV ribozyme gene

-91-

[referred to as ribozyme D by Chang et al. [(1990) Clin. Biotech. 2:23-31]; see also U.S. Patent No. 5,144,019 to Rossi et al., particularly Figure 4 of the patent] flanked by the human β -actin promoter at the 5' end of the gene and the SV40 late transcriptional termination and polyadenylation signals at the 3' end of the gene. As described by Chang et al. [(1990) Clin. Biotech. 2:23-31], ribozyme D is targeted for cleavage of the translational initiation region of the HIV gag gene. This fragment of pPC-RAG12 was subcloned into pBluescript-KS(+) [Stratagene, La Jolla, CA] to produce plasmid 132. Plasmid 132 was then digested with XhoI and EcoRI to yield a fragment containing the ribozyme D gene flanked by the β -actin promoter at the 5' end and the SV40 termination and polyadenylation signals at the 3' end of the gene. This fragment was ligated to the largest fragment generated by digestion of pHyg [Sugden et al. (1985) Mol. Cell. Biol. 5:410-413] with EcoRI and SalI to yield pH132. Thus, pH132 is an ~9.3 kb plasmid containing the following elements: the β -actin promoter linked to an anti-HIV ribozyme gene followed by the SV40 termination and polyadenylation signals, the thymidine kinase gene promoter linked to the hygromycin-resistance gene followed by the thymidine kinase gene polyadenylation signal, and the E. coli ColE1 origin of replication and the ampicillin-resistance gene.

The plasmid pHyg [see, e.g., U.S. Patent Nos. 4,997,764, 4,686,186 and 5,162,215], which confers resistance to hygromycin B using transcriptional controls from the HSV-1 tk gene, was originally constructed from pKan2 [Yates et al. (1984) Proc. Natl. Acad. Sci. U.S.A. 81:3806-3810] and pLG89 [see, Gritz et al. (1983) Gene 25:179-188]. Briefly pKan2 was digested with SmaI and BglII to remove the sequences derived from transposon Tn5. The hygromycin-resistance hph gene was inserted into the digested pKan2 using blunt-end ligation at the SnaI site and "sticky-end" ligation [using 1 Weiss unit of T4 DNA

-92-

ligase (BRL) in 20 microliter volume] at the BglIII site. The SmaI and BglII sites of pKan2 were lost during ligation.

The resulting plasmid pH132, produced from introduction of the anti-HIV ribozyme construct with promoter and polyA site into pHyg,

- 5 includes the anti-HIV ribozyme under control of the β -actin promoter as well as the hygromycin-resistance gene under control of the TK promoter.

2. Chromosome banding

Trypsin G-banding of chromosomes was performed as described in

10 EXAMPLE 1.

3. Cell cultures

TF1004G19 and TF1004G-19C5 mouse cells and the 19C5xHa4 hybrid, described below, and its sublines were cultured in F-12 medium containing 400 μ g/ml Hygromycin B [Calbiochem].

15 B. Cotransfection of EC3/7C5 to produce TF1004G19

Cotransfection of EC3/7C5 cells with plasmids [pH132, pCH110 available from Pharmacia, see, also Hall et al. (1983) J. Mol. Appl. Gen. 2:101-109] and with λ DNA [λ cl 875 Sam 7(New England Biolabs)] was conducted using the calcium phosphate DNA precipitation method [see, 20 e.g., Chen et al. (1987) Mol. Cell. Biol. 7:2745-2752], using 2-5 μ g plasmid DNA and 20 μ g λ phage DNA per 5×10^6 recipient cells.

C. Cell lines containing the sausage chromosome

Analysis of one of the transformants, designated TF1004G19, revealed that it has a high copy number of integrated pH132 and 25 pCH110 sequences, and a high level of β -galactosidase expression. G-banding and *in situ* hybridization with a human probe [CM8; see, e.g., U.S. application S rial No. 08/375,271] revealed unexpectedly that integration had occurred in the formerly dicentric chromosome 7 of the EC3/7C5 c ll line. Furthermore, this chromosome carried a newly formed

-93-

heterochromatic chromosome arm. The size of this heterochromatic arm varied between ~ 150 and ~ 800 Mb in individual metaphases.

By single cell cloning from the TF1004G19 cell line, a subclone TF1004G-19C5 [FIG 2D], which carries a stable chromosome 7 with a
5 ~ 100-150 Mb heterochromatic arm [the sausage chromosome] was obtained. This cell line has been deposited in the ECACC under Accession No. 96040926. This chromosome arm is composed of four to five satellite segments rich in satellite DNA, and evenly spaced integrated heterologous "foreign" DNA sequences. At the end of the compact
10 heterochromatic arm of the sausage chromosome, a less condensed euchromatic terminal segment is regularly observed. This subclone was used for further analyses.

D. Demonstration that the sausage chromosome is derived from the formerly dicentric chromosome

15 *In situ* hybridization with λ phage and pH132 DNA on the TF1004G-19C5 cell line showed positive hybridization only on the minichromosome and on the heterochromatic arm of the "sausage" chromosome [Fig. 2D]. It appears that the "sausage" chromosome [herein also referred to as the SC] developed from the formerly dicentric
20 chromosome (FD) of the EC3/7C5 cell line.

To establish this, the integration sites of pCH110 and pH132 plasmids were determined. This was accomplished by *in situ* hybridization on these cells with biotin-labeled subfragments of the hygromycin-resistance gene and the β -galactosidase gene. Both
25 experiments resulted in narrow hybridizing bands on the heterochromatic arm of the sausage chromosome. The same hybridization pattern was detected on the sausage chromosome using a mixture of biotin-labeled λ probe and pH132 plasmid, proving the cointegration of λ phages, pH132 and pCH110 plasmids.

-94-

To examine this further, the cells were cultured in the presence of the DNA-binding dye Hoechst 33258. Culturing of mouse cells in the presence of this dye results in under-condensation of the pericentric heterochromatin of metaphase chromosomes, thereby permitting better observation of the hybridization pattern. Using this technique, the heterochromatic arm of the sausage chromosome of TF1004G-19C5 cells showed regular under-condensation revealing the details of the structure of the "sausage" chromosome by *in situ* hybridization. Results of *in situ* hybridization on Hoechst-treated TF1004G-19C5 cells with biotin-labeled subfragments of hygromycin-resistance and β -galactosidase genes shows that these genes are localized only in the heterochromatic arm of the sausage chromosome. In addition, an equal banding hybridization pattern was observed. This pattern of repeating units [amplicons] clearly indicates that the sausage chromosome was formed by an amplification process and that the λ phage, pH132 and pCH110 plasmid DNA sequences border the amplicons.

In another series of experiments using fluorescence *in situ* hybridization [FISH] carried out with mouse major satellite DNA, the main component of the mouse pericentric heterochromatin, the results confirmed that the amplicons of the sausage chromosome are primarily composed of satellite DNA.

E. The sausage chromosome has one centromere

To determine whether mouse centromeric sequences had participated in the amplification process forming the "sausage" chromosome and whether or not the amplicons carry inactive centromeres, *in situ* hybridization was carried out with mouse minor satellite DNA. Mouse minor satellite DNA is localized specifically near the centromeres of all mouse chromosomes. Positive hybridization was detected in all mouse centromeres including the sausage chromosome,

-95-

which, however, only showed a positive signal at the beginning of the heterochromatic arm.

Indirect immunofluorescence with a human anti-centromere antibody [LU 851] which recognizes only functional centromeres [see, 5 e.g., Hadlaczky et al. (1989) Chromosoma 97:282-288] proved that the sausage chromosome has only one active centromere. The centromere comes from the formerly dicentric part of the chromosome and co-localizes with the *in situ* hybridization signal of the mouse minor DNA probe.

10 F. The selected and non-selected heterologous DNA in the heterochromatin of the sausage chromosome is expressed

1. High levels of the heterologous genes are expressed

The TF1004G-19C5 cell line thus carries multiple copies of hygromycin-resistance and β -galactosidase genes localized only in the 15 heterochromatic arm of the sausage chromosome. The TF1004G-19C5 cells can grow very well in the presence of 200 μ g/ml or even 400 μ g/ml hygromycin B. [The level of expression was determined by Northern hybridization with a subfragment of the hygromycin-resistance gene and single copy gene.]

20 The expression of the non-selected β -galactosidase gene in the TF1004G-19C5 transformant was detected with LacZ staining of the cells. By this method one hundred percent of the cells stained dark blue, showing that there is a high level of β -galactosidase expression in all of TF1004G-19C5 cells.

25 2. The heterologous genes that are expressed are in the heterochromatin of the sausage chromosome

To demonstrate that the genes localized in the constitutive heterochromatin of the sausage chromosome provide the hygromycin resistance and the LacZ staining capability of TF1004G-19C5 30 transformants [i.e. β -gal expression], PEG-induced cell fusion between

-96-

TF1004G-19C5 mouse cells and Chinese hamster ovary cells was performed. The hybrids were selected and maintained in HAT medium containing G418 [400 μ g/ml] and hygromycin [200 μ g/ml]. Two hybrid clones designated 19C5xHa3 and 19C5xHa4, which have been

5 deposited in the ECACC under Accession No. 96040927, were selected. Both carry the sausage chromosome and the minichromosome.

Twenty-seven single cell derived colonies of the 19C5xHa4 hybrid were maintained and analyzed as individual subclones. *In situ* hybridization with hamster and mouse chromosome painting probes and

10 hamster chromosome 2-specific probes verified that the 19C5xHa4 clone contains the complete Chinese hamster genome and a partial mouse genome. All 19C5xHa4 subclones retained the hamster genome, but different subclones showed different numbers of mouse chromosomes indicating the preferential elimination of mouse chromosomes.

15 To promote further elimination of mouse chromosomes, hybrid cells were repeatedly treated with BrdU. The BrdU treatments, which destabilize the genome, result in significant loss of mouse chromosomes. The BrdU-treated 19C5xHa4 hybrid cells were divided to three groups. One group of the hybrid cells (GH) were maintained in the presence of

20 hygromycin (200 μ g/ml) and G418 (400 μ g/ml), and the other two groups of the cells were cultured under G418 (G) or hygromycin (H) selection conditions to promote the elimination of the sausage chromosome or minichromosome.

One month later, single cell derived subclones were established

25 from these three subcultures of the 19C5xHa4 hybrid line. The subclones were monitored by *in situ* hybridization with biotin-labeled λ phage and hamster chromosome painting probes. Four individual clones [G2B5, G3C5, G4D6, G2B4] selected in the presence of G418 that had lost the sausage chromosome but retained the minichromosome were

-97-

found. Under hygromycin selection only one subclone (H17D1) lost the minichromosome. In this case the minichromosome [see Example 5] was present.

Since hygromycin-resistance and β -galactosidase genes were
5 thought to be expressed from the sausage chromosome, the expression of these genes was analyzed in the four subclones that had lost the sausage chromosome. In the presence of 200 μ g/ml hygromycin, one hundred percent of the cells of four individual subclones died. In order to detect the β -galactosidase expression hybrid, subclones were analyzed
10 by LacZ staining. One hundred percent of the cells of the four subclones that lost the sausage chromosome also lost the LacZ staining capability. All of the other hybrid subclones that had not lost the sausage chromosome under the non-selective culture conditions showed positive LacZ staining.

15 These findings demonstrate that the expression of hygromycin-resistance and β -galactosidase genes is linked to the presence of the sausage chromosome. Results of *in situ* hybridizations show that the heterologous DNA is expressed from the constitutive heterochromatin of the sausage chromosome.

20 *In situ* hybridization studies of three other hybrid subclones [G2C6, G2D1, and G4D5] did not detect the presence of the sausage chromosome. By the LacZ staining method, some stained cells were detected in these hybrid lines, and when these subclones were transferred to hygromycin selection some colonies survived. Cytological
25 analysis and *in situ* hybridization of these hygromycin-resistant colonies revealed the presence of the sausage chromosome, suggesting that only the cells of G2C6, G2D1 and G4D5 hybrids that had not lost the sausage chromosome were able to preserve the hygromycin resistance and β -galactosidase expression. These results confirmed that the expression of

-98-

these genes is linked to the presence of the sausage chromosome. The level of β -galactosidase expression was determined by the immunoblot technique using a monoclonal antibody.

Hygromycin resistance and β -galactosidase expression of the cells which contained the sausage chromosome were provided by the genes localized in the mouse pericentric heterochromatin. This was demonstrated by performing Southern DNA hybridizations on the hybrid cells that lack the sausage chromosome using PCR-amplified subfragments of hygromycin-resistance and β -galactosidase genes as probes. None of the subclones showed hybridization with these probes; however, all of the analyzed clones contained the minichromosome. Other hybrid clones that contain the sausage chromosome showed intense hybridization with these DNA probes. These results lead to the conclusion that hygromycin resistance and β -galactosidase expression of the cells that contain the sausage chromosome were provided by the genes localized in the mouse pericentric heterochromatin.

EXAMPLE 5

The gigachromosome

As described in Example 4, the sausage chromosome was transferred into Chinese hamster cells by cell fusion. Using Hygromycin B/HAT and G418 selection, two hybrid clones 19C5xHa3 and 19C5xHa4 were produced that carry the sausage chromosome. *In situ* hybridization, using hamster and mouse chromosome-painting probes and a hamster chromosome 2-specific probe, verified that clone 19C5xHa4 contains a complete Chinese hamster genome as well as partial mouse genomes. Twenty-seven separate colonies of 19C5xHa4 cells were maintained and analyzed as individual subclones. Twenty-six out of 27 subclones contained a morphologically unchanged sausage chromosome.

-99-

In one subclone of the 19C5xHa3 cell line, 19C5xHa47 [see Fig. 2E], the heterochromatic arm of the sausage chromosome became unstable and showed continuous intrachromosomal growth. In extreme cases, the amplified chromosome arm exceeded 1000 Mb in size (gigachromosome).

EXAMPLE 6

The stable megachromosome

A. Generation of cell lines containing the megachromosome

All 19C5xHa4 subclones retained a complete hamster genome, but different subclones showed different numbers of mouse chromosomes, indicating the preferential elimination of mouse chromosomes. As described in Example 4, to promote further elimination of mouse chromosomes, hybrid cells were treated with BrdU, cultured under G418 (G) or hygromycin (H) selection conditions followed by repeated treatment with 10^{-4} M BrdU for 16 hours and single cell subclones were established. The BrdU treatments appeared to destabilize the genome, resulting in a change in the sausage chromosome as well. A gradual increase in a cell population in which a further amplification had occurred was observed. In addition to the ~100-150 Mb heterochromatic arm of the sausage chromosome, an extra centromere and a ~150-250 Mb heterochromatic chromosome arm were formed, which differed from those of mouse chromosome 7. By the acquisition of another euchromatic terminal segment, a new submetacentric chromosome (megachromosome) was formed. Seventy-nine individual subclones were established from these BrdU-treated cultures by single-cell cloning: 42 subclones carried the intact megachromosome, 5 subclones carried the sausage chromosome, and in 32 subclones fragments or translocated segments of the megachromosome were observed. Twenty-six subclones that carried the megachromosome were cultured under non-

-100-

selective conditions over a two-month period. In 19 out of 26 subclones, the megachromosome was retained. Those subclones which lost the megachromosomes all became sensitive to Hygromycin B and had no β -galactosidase expression, indicating that both markers were
5 linked to the megachromosome.

Two sublines (G3D5 and H1D3), which were chosen for further experiments, showed no changes in the morphology of the megachromosome during more than 100 generations under selective conditions. The G3D5 cells had been obtained by growth of 19C5xHa4
10 cells in G418-containing medium followed by repeated BrdU treatment, whereas H1D3 cells had been obtained by culturing 19C5xHa4 cells in hygromycin-containing medium followed by repeated BrdU treatment.

B. Structure of the megachromosome

The following results demonstrate that, apart from the euchromatic
15 terminal segments, the integrated foreign DNA (and as in the exemplified embodiments, rDNA sequence), the whole megachromosome is constitutive heterochromatin, containing a tandem array of at least 40 [~ 7.5 Mb] blocks of mouse major satellite DNA [see Figures 2 and 3]. Four satellite DNA blocks are organized into a giant palindrome
20 [amplicon] carrying integrated exogenous DNA sequences at each end. The long and short arms of the submetacentric megachromosome contains 6 and 4 amplicons, respectively. It is of course understood that the specific organization and size of each component can vary among species, and also the chromosome in which the amplification
25 event initiates.

1. The megachromosome is composed primarily of heterochromatin

Except for the terminal regions and the integrated foreign DNA, the megachromosome is composed primarily of heterochromatin. This was

-101-

- demonstrated by C-banding of the megachromosome, which resulted in positive staining characteristic of constitutive heterochromatin. Apart from the terminal regions and the integrated foreign DNA, the whole megachromosome appears to be heterochromatic. Mouse major satellite
- 5 DNA is the main component of the pericentric, constitutive heterochromatin of mouse chromosomes and represents ~10% of the total DNA [Waring *et al.* (1966) *Science* 154:791-794]. Using a mouse major satellite DNA probe for *in situ* hybridization, strong hybridization was observed throughout the megachromosome, except for its terminal
- 10 regions. The hybridization showed a segmented pattern: four large blocks appeared on the short arm and usually 4-7 blocks were seen on the long arm. By comparing these segments with the pericentric regions of normal mouse chromosomes that carry ~15 Mb of major satellite DNA, the size of the blocks of major satellite DNA on the
- 15 megachromosome was estimated to be ~30 Mb.
- Using a mouse probe specific to euchromatin [pMCPE1.51; a mouse long interspersed repeated DNA probe], positive hybridization was detected only on the terminal segments of the megachromosome of the H1D3 hybrid subline. In the G3D5 hybrids, hybridization with a hamster-
- 20 specific probe revealed that several megachromosomes contained terminal segments of hamster origin on the long arm. This observation indicated that the acquisition of the terminal segments on these chromosomes happened in the hybrid cells, and that the long arm of the megachromosome was the recently formed one arm. When a mouse
- 25 minor satellite probe was used, specific to the centromeres of mouse chromosomes [Wong *et al.* (1988) *Nucl. Acids Res.* 16:11645-11661], a strong hybridization signal was detected only at the primary constriction of the megachromosome, which colocalized with the positive

-102-

immunofluorescence signal produced with human anti-centromere serum [LU 851].

- In situ* hybridization experiments with pH132, pCH110, and λ DNA probes revealed that all heterologous DNA was located in the gaps
- 5 between the mouse major satellite DNA segments. Each segment of mouse major satellite DNA was bordered by a narrow band of integrated heterologous DNA, except at the second segment of the long arm where a double band of heterologous DNA existed, indicating that the major satellite DNA segment was missing or considerably reduced in size here.
- 10 This chromosome region served as a useful cytological marker in identifying the long arm of the megachromosome. At a frequency of 10^{-4} , "restoration" of these missing satellite DNA blocks was observed in one chromatid, when the formation of a whole segment on one chromatid occurred.
- 15 After Hoechst 33258 treatment (50 μ g/ml for 16 hours), the megachromosome showed undercondensation throughout its length except for the terminal segments. This made it possible to study the architecture of the megachromosome at higher resolution. *In situ* hybridization with the mouse major satellite probe on undercondensed
- 20 megachromosomes demonstrated that the ~ 30 Mb major satellite segments were composed of four blocks of ~ 7.5 Mb separated from each other by a narrow band of non-hybridizing sequences [Figure 3]. Similar segmentation can be observed in the large block of pericentric heterochromatin in metacentric mouse chromosomes from the LMTK and
- 25 A9 cell lines.

2. The megachromosome is composed of segments containing two tandem ~ 7.5 Mb blocks followed by two inverted blocks

- Because of the asymmetry in thymidine content between the two
- 30 strands of the DNA of the mouse major satellite, when mouse cells are

-103-

grown in the presence of BrdU for a single S phase, the constitutive heterochromatin shows lateral asymmetry after FPG staining. Also, in the 19C5xHa4 hybrids, the thymidine-kinase [*Tk*] deficiency of the mouse fibroblast cells was complemented by the hamster *Tk* gene, 5 permitting BrdU incorporation experiments.

A striking structural regularity in the megachromosome was detected using the FPG technique. In both chromatids, alternating dark and light staining that produced a checkered appearance of the megachromosome was observed. A similar picture was obtained by 10 labelling with fluorescein-conjugated anti-BrdU antibody. Comparing these pictures to the segmented appearance of the megachromosome showed that one dark and one light FPG band corresponded to one ~30 Mb segment of the megachromosome. These results suggest that the two halves of the ~30 Mb segment have an inverted orientation. This 15 was verified by combining *in situ* hybridization and immunolabelling of the incorporated BrdU with fluorescein-conjugated anti-BrdU antibody on the same chromosome. Since the ~30 Mb segments [or amplicons] of the megachromosome are composed of four blocks of mouse major satellite DNA, it can be concluded that two tandem ~7.5 Mb blocks are 20 followed by two inverted blocks within one segment.

Large-scale mapping of megachromosome DNA by pulsed-field electrophoresis and Southern hybridization with "foreign" DNA probes revealed a simple pattern of restriction fragments. Using endonucleases with none, or only a single cleavage site in the integrated foreign DNA 25 sequences, followed by hybridization with a *hyg* probe, 1-4 predominant fragments were detected. Since the megachromosome contains 10-12 amplicons with an estimated 3-8 copies of *hyg* sequences per amplicon (30-90 copies per megachromosome), the small number of hybridizing fragments indicates the homogeneity of DNA in the amplified segments.

-104-

3. Scanning electron microscopy of the megachromosome confirmed the above findings

The homogeneous architecture of the heterochromatic arms of the megachromosome was confirmed by high resolution scanning electron
5 microscopy. Extended arms of megachromosomes, and the pericentric heterochromatic region of mouse chromosomes, treated with Hoechst 33258, showed similar structure. The constitutive heterochromatic regions appeared more compact than the euchromatic segments. Apart from the terminal regions, both arms of the megachromosome were
10 completely extended, and showed faint grooves, which should correspond to the border of the satellite DNA blocks in the non-amplified chromosomes and in the megachromosome. Without Hoechst treatment, the grooves seemed to correspond to the amplicon borders on the megachromosome arms. In addition, centromeres showed a more
15 compact, finely fibrous appearance than the surrounding heterochromatin.

4. The megachromosome of 1B3 cells contains rRNA gene sequence

The sequence of the megachromosome in the region of the sites of
20 integration of the heterologous DNA was investigated by isolation of these regions through using cloning methods and sequence analysis of the resulting clones. The results of this analysis revealed that the heterologous DNA was located near mouse ribosomal RNA gene (i.e., rDNA) sequences contained in the megachromosome.

25 a. Cloning of regions of the megachromosomes in which heterologous DNA had integrated

Megachromosomes were isolated from 1B3 cells (which were generated by repeated BrdU treatment and single cell cloning of H1xHE41 cells (see Figure 4) and which contain a truncated
30 megachromosome) using fluorescence-activated cell sorting methods as

-105-

described herein (see Example 10). Following separation of the SATACs (megachromosomes) from the endogenous chromosomes, the isolated megachromosomes were stored in GH buffer (100 mM glycine, 1% hexylene glycol, pH 8.4-8.6 adjusted with saturated calcium hydroxide solution; see Example 10) and centrifuged into an agarose bed in 0.5 M EDTA.

Large-scale mapping of the megachromosome around the area of the site of integration of the heterologous DNA revealed that it is enriched in sequence containing rare-cutting enzyme sites, such as the recognition site for NotI. Additionally, mouse major satellite DNA (which makes up the majority of the megachromosome) does not contain NotI recognition sites. Therefore, to facilitate isolation of regions of the megachromosome associated with the site of integration of the heterologous DNA, the isolated megachromosomes were cleaved with NotI, a rare cutting restriction endonuclease with an 8-bp GC recognition site. Fragments of the megachromosome were inserted into plasmid pWE15 (Stratagene, La Jolla, California) as follows. Half of a 100- μ l low melting point agarose block (mega-plug) containing the isolated SATACs was digested with NotI overnight at 37°C. Plasmid pWE15 was similarly digested with NotI overnight. The mega-plug was then melted and mixed with the digested plasmid, ligation buffer and T4 ligase. Ligation was conducted at 16°C overnight. Bacterial DH5 α cells were transformed with the ligation product and transformed cells were plated onto LB/Amp plates. Fifteen to twenty colonies were grown on each plate for a total of 189 colonies. Plasmid DNA was isolated from colonies that survived growth on LB/Amp medium and was analyzed by Southern blot hybridization for the presence of DNA that hybridized to a pUC19 probe. This screening methodology assured that all clones, even clones lacking an insert but yet containing the pWE15 plasmid, would be detected. Any

-106-

clones containing insert DNA would be expected to contain non-satellite, GC-rich megachromosome DNA sequences located at the site of integration of the heterologous DNA. All colonies were positive for hybridizing DNA.

- 5 Liquid cultures of all 189 transformants were used to generate cosmid minipreps for analysis of restriction sites within the insert DNA. Six of the original 189 cosmid clones contained an insert. These clones were designated as follows: 28 (~9-kb insert), 30 (~9-kb insert), 60 (~4-kb insert), 113 (~9-kb insert), 157 (~9-kb insert) and 161 (~9-kb insert). Restriction enzyme analysis indicated that three of the clones
- 10 (113, 157 and 161) contained the same insert.

b. *In situ* hybridization experiments using isolated segments of the megachromosome as probes

- Insert DNA from clones 30, 113, 157 and 161 was purified,
- 15 labeled and used as probes in *in situ* hybridization studies of several cell lines. Counterstaining of the cells with propidium iodide facilitated identification of the cytological sites of the hybridization signals. The locations of the signals detected within the cells are summarized in the following table:

CELL TYPE	PROBE	LOCATION OF SIGNAL
Human Lymphocyte (male)	No. 161	4-5 pairs of acrocentric chromosomes at centromeric regions.
Mouse Spleen	No. 161	Acrocentric ends of 4 pairs of chromosomes.
EC3/7C5 Cells	No. 161	Minichromosome and the end of the formerly dicentric chromosome. Pericentric heterochromatin of one of the metacentric mouse chromosomes. Centromeric region of some of the other mouse chromosomes.
K20 Chinese Hamster Cells	No. 30	Ends of at least 6 pairs of chromosomes. An interstitial signal on a short chromosome.

-107-

CELL TYPE	PROBE	LOCATION OF SIGNAL
HB31 Cells (mouse-hamster hybrid cells derived from H1D3 cells by repeated BrdU treatment and single cell cloning which carries the megachromosome)	No. 30	Acrocentric ends of at least 12 pairs of chromosomes. Centromeres of certain chromosomes and the megachromosome. Borders of the amplicons of the megachromosome.
Mouse Spleen Cells	No. 30	Similar to signal observed for probe no. 161. Centromeres of 5 pairs of chromosomes. Weak cross-hybridization to pericentric heterochromatin.
HB31 Cells	No. 113	Similar to signal observed for probe no. 30.
Mouse Spleen Cells	No. 113	Centromeric region of 5 pairs of chromosomes.
K20 Cells	No. 113	At least 6 pairs of chromosomes. Weak signal at some telomeres and several interspersed signals.
Human Lymphocyte Cells (male)	No. 157	Similar to signal observed for probe no. 161.

c. Southern blot hybridization using isolated segments of the megachromosome as probes

DNA was isolated from mouse spleen tissue, mouse LMTK⁺ cells, K20 Chinese hamster ovary cells, EJ30 human fibroblast cells and H1D3 cells. The isolated DNA and lambda phage DNA, was subjected to Southern blot hybridization using inserts isolated from megachromosome clone nos. 30, 113, 157 and 161 as probes. Plasmid pWE15 was used as a negative control probe. Each of the four megachromosome clone inserts hybridized in a multi-copy manner (as demonstrated by the intensity of hybridization and the number of hybridizing bands) to all of the DNA samples, except the lambda phage DNA. Plasmid pWE15 hybridized to lambda DNA only.

-108-

d. Sequence analysis of megachromosome clone no. 161

Megachromosome clone no. 161 appeared to show the strongest hybridization in the *in situ* and Southern hybridization experiments and
5 was chosen for analysis of the insert sequence. The sequence analysis was approached by first subcloning the insert of cosmid clone no. 161 to obtain five subclones as follows.

To obtain the end fragments of the insert of clone no. 161, the clone was digested with NotI and BamHI and ligated with NotI/BamHI-
10 digested pBluescript KS (Stratagene, La Jolla, California). Two fragments of the insert of clone no. 161 were obtained: a 0.2-kb and a 0.7-kb insert fragment. To subclone the internal fragment of the insert of clone no. 161, the same digest was ligated with BamHI-digested pUC19. Three fragments of the insert of clone no. 161 were obtained: a 0.6-kb,
15 a 1.8-kb and a 4.8-kb insert fragment.

The ends of all the subcloned insert fragments were first sequenced manually. However, due to their extremely high GC content, autoradiographs were difficult to interpret and sequencing was repeated using an ABI sequencer and the dye-terminator cycle protocol. A
20 comparison of the sequence data to sequences in the GENBANK database revealed that the insert of clone no. 161 corresponds to an internal section of the mouse ribosomal RNA gene (rDNA) repeat unit between positions 7551-15670 as set forth in GENBANK accession no. X82564, which is provided as SEQ ID NO. 16 herein. The sequence
25 data obtained for the insert of clone no. 161 is set forth in SEQ ID NOS. 18-24. Specifically, the individual subclones corresponded to the following positions in GENBANK accession no. X82564 (i.e., SEQ ID NO. 16) and in SEQ ID NOS. 18-24:

-109-

5	Subclone	Start	End	Site	SEQ ID No.
		in X82564			
	161k1	7579	7755	<u>NotI</u> , <u>BamHI</u>	18
	161m5	7756	8494	<u>BamHI</u>	19
	161m7	8495	10231	<u>BamHI</u>	20 (shows only sequence corresponding to nt. 8495-8950), 21 (shows only sequence corresponding to nt. 9851- 10231)
	161m12	10232	15000	<u>BamHI</u>	22 (shows only sequence corresponding to nt. 10232-10600), 23 (shows only sequence corresponding to nt. 14267-15000).
	161k2	15001	15676	<u>NotI</u> , <u>BamHI</u>	24

The sequence set forth in SEQ ID NOs. 18-24 diverges in some positions from the sequence presented in positions 7551-15670 of GENBANK accession no. X82564. Such divergence may be attributable to random mutations between repeat units of rDNA. The results of the sequence analysis of clone no. 161, which reveal that it corresponds to rDNA, correlate with the appearance of the *in situ* hybridization signal it generated in human lymphocytes and mouse spleen cells. The hybridization signal was clearly observed on acrocentric chromosomes in these cells, and such types of chromosomes are known to include rDNA adjacent to the pericentric satellite DNA on the short arm of the chromosome. Furthermore, rRNA genes are highly conserved in mammals as supported by the cross-species hybridization of clone no. 161 to human chromosomal DNA.

To isolate amplification-replication control regions such as those found in rDNA, it may be possible to subject DNA isolated from megachromosome-containing cells, such as H1D3 cells, to nucleic acid amplification using, e.g., the polymerase chain reaction (PCR) with the following primers:

-110-

amplification control element forward primer (1-30)

5'-GAGGAATTCCCCATCCCTAATCCAGATTGGTG-3' (SEQ ID NO. 25)

amplification control element reverse primer (2142-2111)

5'-AAACTGCAGGCCGAGCCACCTCTCTTCTGTGTTTG-3' (SEQ ID NO. 26)

5 origin of replication region forward primer (2116-2141)

5'-AGGAATTCACAGAAGAGAGGTGGCTCGGCCTGC-3' (SEQ ID NO. 27)

origin of replication region reverse primer (5546-5521)

5'-AGCCTGCAGGAAGTCATACCTGGGGAGGTGGCCC-3' (SEQ ID NO. 28)

C. Summary of the formation of the megachromosome

- 10 Figure 2 schematically sets forth events leading to the formation of a stable megachromosome beginning with the generation of a dicentric chromosome in a mouse LMTK⁻ cell line: (A) A single E-type amplification in the centromeric region of the mouse chromosome 7 following transfection of LMTK⁻ cells with λ CM8 and λ gtWESneo generates the
- 15 neo-centromere linked to the integrated foreign DNA, and forms a dicentric chromosome. Multiple E-type amplification forms the λ neo-chromosome, which was derived from chromosome 7 and stabilized in a mouse-hamster hybrid cell line; (B) Specific breakage between the centromeres of a dicentric chromosome 7 generates a chromosome
- 20 fragment with the neo-centromere, and a chromosome 7 with traces of foreign DNA at the end; (C) Inverted duplication of the fragment bearing the neo-centromere results in the formation of a stable neo-minichromosome; (D) Integration of exogenous DNA into the foreign DNA region of the formerly dicentric chromosome 7 initiates H-type
- 25 amplification, and the formation of a heterochromatic arm. By capturing a euchromatic terminal segment, this new chromosome arm is stabilized in the form of the "sausage" chromosome; (E) BrdU treatment and/or drug selection appears to induce further H-type amplification, which results in the formation of an unstable gigachromosome: (F) Repated

-111-

BrdU treatments and/or drug selection induce further H-type amplification including a centromere duplication, which leads to the formation of another heterochromatic chromosome arm. It is split off from the chromosome 7 by chromosome breakage and acquires a terminal
5 segment to form the stable megachromosome.

D. Expression of β -galactosidase and hygromycin transferase genes in cell lines carrying the megachromosome or derivatives thereof

The level of heterologous gene (i.e., β -galactosidase and
hygromycin transferase genes) expression in cell lines containing the
10 megachromosome or a derivative thereof was quantitatively measured.
The relationship between the copy-number of the heterologous genes
and the level of protein expressed therefrom was also determined.

1. Materials and methods

a. Cell lines

15 Heterologous gene expression levels of H1D3 cells, carrying a
250-400 Mb megachromosome as described above, and mM2C1 cells,
carrying a 50-60 Mb micro-megachromosome, were quantitatively
evaluated. mM2C1 cells were generated by repeated BrdU treatment and
single cell cloning of the H1xHe41 cell line (mouse-hamster-human hybrid
20 cell line carrying the megachromosome and a single human chromosome
with CD4 and neo' genes; see Figure 4). The cell lines were grown under
standard conditions in F12 medium under selective (120 μ g/ml
hygromycin) or non-selective conditions.

b. Preparation of cell extract for β -galactosidase assays

25 Monolayers of mM2C1 or H1D3 cell cultures were washed three
times with phosphate-buffered saline (PBS). Cells were scraped by
rubber policemen and suspended and washed again in PBS. Washed
cells were resuspended into 0.25 M Tris-HCl, pH 7.8, and disrupted by

-112-

three cycles of freezing in liquid nitrogen and thawing at 37°C. The extract was clarified by centrifugation at 12,000 rpm for 5 min. at 4°C.

c. β -galactosidase assay

The β -galactosidase assay mixture contained 1 mM $MgCl_2$,
5 45 mM β -mercaptoethanol, 0.8 mg/ml o-nitrophenyl- β -D-galactopyranoside and 66 mM sodium phosphate, pH 7.5. After incubating the reaction mixture with the cell extract at 37°C for increasing time, the reaction was terminated by the addition of three volumes of 1M Na_2CO_3 , and the optical density was measured at 420 nm. Assay mixture incubated
10 without cell extract was used as a control. The linear range of the reaction was determined to be between 0.1-0.8 OD₄₂₀. One unit of β -galactosidase activity is defined as the amount of enzyme that will hydrolyse 3 nmoles of o-nitrophenyl- β -D-galactopyranoside in 1 minute at 37 C.

d. Preparation of cell extract for hygromycin phosphotransferase assay

15 Cells were washed as described above and resuspended into 20 mM Hepes buffer, pH 7.3, 100 mM potassium acetate, 5 mM Mg acetate and 2 mM dithiothreitol). Cells were disrupted at 0°C by six 10 sec bursts in an MSE ultrasonic disintegrator using a microtip probe. Cells
20 were allowed to cool for 1 min after each ultrasonic burst. The extracts were clarified by centrifuging for 1 min at 2000 rpm in a microcentrifug .

e. Hygromycin phosphotransferase assay

Enzyme activity was measured by means of the phosphocellulose paper binding assay as described by Haas and Dowding [(1975). Meth.
25 Enzymol. 43:611-628]. The cell extract was supplemented with 0.1 M ammonium chloride and 1 mM adenosine- γ -³²P-triphosphate (specific activity: 300 Ci/mmol). The reaction was initiated by the addition of 0.1 mg/ml hygromycin and incubated for increasing time at 37°C. The reaction was terminated by heating the samples for 5 min at 75 C in a

-113-

water bath, and after removing the precipitated proteins by centrifugation for 5 min in a microcentrifuge, an aliquot of the supernatant was spotted on a piece of Whatman P-81 phosphocellulose paper (2 cm²). After 30 sec at room temperature the papers are placed into 500 ml of hot (75°C) distilled water for 3 min. While the radioactive ATP remains in solution under these conditions, hygromycin phosphate binds strongly and quantitatively to phosphocellulose. The papers are rinsed 3 times in 500 ml of distilled water and the bound radioactivity was measured in toluene scintillation cocktail in a Beckman liquid scintillation counter. Reaction mixture incubated without added hygromycin served as a control.

f. Determination of the copy-number of the heter log us genes

DNA was prepared from the H1D3 and mM2C1 cells using standard purification protocols involving SDS lysis of the cells followed by Proteinase K treatment and phenol/chloroform extractions. The isolated DNA was digested with an appropriate restriction endonuclease, fractionated on agarose gels, blotted to nylon filters and hybridized with a radioactive probe derived either from the β -galactosidase or the hygromycin phosphotransferase genes. The level of hybridization was quantified in a Molecular Dynamics PhosphorImage Analyzer. To control the total amount of DNA loaded from the different cells lines, the filters were reprobated with a single copy gene, and the hybridization of β -galactosidase and hygromycin phosphotransferase genes was normalized to the single copy gene hybridization.

g. Determination of protein concentration

The total protein content of the cell extracts was measured by the Bradford colorimetric assay using bovine serum albumin as standard.

-114-

2. Characterization of the β -galactosidase and hygromycin phosphotransferase activity expressed in H1D3 and mM2C1 cells

In order to establish quantitative conditions, the most important kinetic parameters of β -galactosidase and hygromycin phosphotransferase activity have been studied. The β -galactosidase activity measured with this colorimetric assay was linear between the 0.1-0.8 OD₄₂₀ range both for the mM2C1 and H1D3 cell lines. The β -galactosidase activity was also proportional in both cell lines with the amount of protein added to the reaction mixture within 5-100 μ g total protein concentration range. The hygromycin phosphotransferase activity of mM2C1 and H1D3 cell lines was also proportional with the reaction time or the total amount of added cell extract under the conditions described for the β -galactosidase.

a. Comparison of β -galactosidase activity of mM2C1 and H1D3 cell lines

Cell extracts prepared from logarithmically growing mM2C1 and H1D3 cell lines were tested for β -galactosidase activity, and the specific activities were compared in 10 independent experiments. The β -galactosidase activity of H1D3 cell extracts was 440 ± 25 U/mg total protein. Under identical conditions the β -galactosidase activity of the mM2C1 cell extracts was 4.8 times lower: 92 ± 13 U/mg total protein.

β -galactosidase activities of highly subconfluent, subconfluent and nearly confluent cultures of H1D3 and mM2C1 cell lines were also compared. In these experiments different numbers of logarithmic H1D3 and mM2C1 cells were seeded in constant volume of culture medium and grown for 3 days under standard conditions. No significant difference was found in the β -galactosidase specific activities of cell cultures grown at different cell densities, and the ratio of H1D3/mM2C1 β -galactosidase specific activities was also similar for all three cell densities. In

-115-

confluent, stationary cell cultures of H1D3 or mM2C1 cells, however, the expression of β -galactosidase significantly decreased due likely to cessation of cell division as a result of contact inhibition.

5 **b. Comparison of hygromycin phosphotransferase activity of H1D3 and mM2C1 cell lines**

The bacterial hygromycin phosphotransferase is present in a membrane-bound form in H1D3 or mM2C1 cell lines. This follows from the observation that the hygromycin phosphotransferase activity can be completely removed by high speed centrifugation of these cell extracts,
10 and the enzyme activity can be recovered by resuspending the high speed pellet.

The ratio of the enzyme's specific activity in H1D3 and mM2C1 cell lines was similar to that of β -galactosidase activity, i.e., H1D3 cells have 4.1 times higher specific activity compared with mM2C1 cells.

15 **c. Hygromycin phosphotransferase activity in H1D3 and mM2C1 cells grown under non-selective conditions**

The level of expression of the hygromycin phosphotransferase gene was measured on the basis of quantitation of the specific enzyme activities in H1D3 and mM2C1 cell lines grown under non-selective
20 conditions for 30 generations. The absence of hygromycin in the medium did not influence the expression of the hygromycin phosphotransferase gene.

25 **3. Quantitation of the number of β -galactosidase and hygromycin phosphotransferase gene copies in H1D3 and mM2C1 cell lines**

As described above, the β -galactosidase and hygromycin phosphotransferase genes are located only within the megachromosome, or micro-megachromosome in H1D3 and mM2C1 cells. Quantitative analysis of genomic Southern blots of DNA isolated from H1D3 and
30 mM2C1 cell lines with the PhosphorImage Analyzer revealed that the

-116-

copy number of β -galactosidase genes integrated into the megachromosome is approximately 10 times higher in H1D3 cells than in mM2C1 cells. The copy-number of hygromycin phosphotransferase genes is approximately 7 times higher in H1D3 cells than in mM2C1 cells.

5 4. **Summary and conclusions of results of quantitation of heterologous gene expression in cells containing megachromosomes or derivatives thereof**

Quantitative determination of β -galactosidase activity of higher eukaryotic cells (e.g., H1D3 cells) carrying the bacterial β -galactosidase gene in heterochromatic megachromosomes confirmed the observed
10 high-level expression of the integrated bacterial gene detected by cytological staining methods. It has generally been established in reports of studies of the expression of foreign genes in transgenic animals that although transgene expression shows correct tissue and developmental
15 specificity, the level of expression is typically low and shows extensive position-dependent variability (i.e., the level of transgene expression depends on the site of chromosomal integration). It is generally assumed that the low-level transgene expression may be due to the absence of special DNA sequences which can insulate the transgene from the
20 inhibitory effect of the surrounding chromatin and promote the formation of active chromatin structure required for efficient gene expression. Several cis-activating DNA sequence elements have been identified which can abolish this position-dependent variability, and can ensure high-level expression of the transgene locus activating region (LAR) sequences in
25 higher eukaryotes and specific chromatin structure (scs) elements in lower eukaryotes (see, e.g., Eissenberg and Elgin (1991) Trends in Genet. 7:335-340). If these cis-acting DNA sequences are absent, the level of transgene expression is low and copy-number independent.

Although the bacterial β -galactosidase reporter gene contained in
30 the heterochromatic megachromosomes of H1D3 and mM2C1 cells is

-117-

driven by a potent eukaryotic promoter-enhancer element, no specific cis-acting DNA sequence element was designed and incorporated into the bacterial DNA construct which could function as a boundary element. Thus, the high-level β -galactosidase expression measured in these cells is of significance, particularly because the β -galactosidase gene in the megachromosome is located in a long, compact heterochromatic environment, which is known to be able to block gene expression. The megachromosome appears to contain DNA sequence element(s) in association with the bacterial DNA sequences that function to override the inhibitory effect of heterochromatin on gene expression.

The specificity of the heterologous gene expression in the megachromosome is further supported by the observation that the level of β -galactosidase expression is copy-number dependent. In the H1D3 cell line, which carries a full-size megachromosome, the specific activity of β -galactosidase is about 5-fold higher than in mM2C1 cells, which carry only a smaller, truncated version of the megachromosome. A comparison of the number of β -galactosidase gene copies in H1D3 and mM2C1 cell lines by quantitative hybridization techniques confirmed that the expression of β -galactosidase is copy-number dependent. The number of integrated β -galactosidase gene copies is approximately 10-fold higher in the H1D3 cells than in mM2C1 cells. Thus, the cell line containing the greater number of copies of the β -galactosidase gene also yields higher levels of β -galactosidase activity, which supports the copy-number dependency of expression. The copy number dependency of the β -galactosidase and hygromycin phosphotransferase enzyme levels in cell lines carrying different derivatives of the megachromosome indicates that neither the chromatin organization surrounding the site of integration of the bacterial genes, nor the heterochromatic environment of the megachromosome suppresses the expression of the genes.

-118-

- The relative amount of β -galactosidase protein expressed in H1D3 cells can be estimated based on the V_{\max} of this enzyme [500 for homogeneous, crystallized bacterial β -galactosidase (Naider *et al.* (1972) Biochemistry 11:3202-3210)] and the specific activity of H1D3 cell
- 5 protein. A V_{\max} of 500 means that the homogeneous β -galactosidase protein hydrolyzes 500 μ moles of substrate per minute per mg of enzyme protein at 37°C. One mg of total H1D3 cell protein extract can hydrolyze 1.4 μ moles of substrate per minute at 37°C, which means that 0.28% of the protein present in the H1D3 cell extract is β -galactosidase.
- 10 The hygromycin phosphotransferase is present in a membrane-bound form in H1D3 and mM2C1 cells. The tendency of the enzyme to integrate into membranes in higher eukaryotic cells may be related to its periplasmic localization in prokaryotic cells. The bacterial hygromycin phosphotransferase has not been purified to homogeneity; thus, its V_{\max}
- 15 has not been determined. Therefore, no estimate can be made on the total amount of hygromycin phosphotransferase protein expressed in these cell lines. The 4-fold higher specific activity of hygromycin phosphotransferase in H1D3 cells as compared to mM2C1 cells, however, indicates that its expression is also copy number dependent.
- 20 The constant and high level expression of the β -galactosidase gene in H1D3 and mM2C1 cells, particularly in the absence of any selective pressure for the expression of this gene, clearly indicates the stability of the expression of genes carried in the heterochromatic megachromosomes. This conclusion is further supported by the observation that the
- 25 level of hygromycin phosphotransferase expression did not change when H1D3 and mM2C1 cells were grown under non-selective conditions. The consistent high-level, stable, and copy-number dependent expression of bacterial marker genes clearly indicates that the megachromosome is an ideal vector system for expression of foreign genes.

-119-

EXAMPLE 7

Summary of some of the cell lines with SATACS and minichromosomes that have been constructed

5 1. EC3/7-Derived cell lines

The LMTK⁻-derived cell line, which is a mouse fibroblast cell line, was transfected with λ CM8 and λ gtWESneo DNA [see, EXAMPLE 2] to produce transformed cell lines. Among these cell lines was EC3/7, deposited at the European Collection of Animal cell Culture (ECACC) under Accession No. 90051001 [see, U.S. Patent No. 5,288,625; see, also Hadlaczky *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110 and U.S. application Serial No. 08/375,271]. This cell line contains the dicentric chromosome with the neo-centromere. Recloning and selection produced cell lines such as EC3/7C5, which are cell lines with the stable neo-minichromosome and the formerly dicentric chromosome [see, Fig. 2C].

2. KE1-2/4 Cells

Fusion of EC3/7 with CHO-K20 cells and selection with G418/HAT produced hybrid cell lines, among these was KE1-2/4, which has been deposited with the ECACC under Accession No. 96040924. KE1-2/4 is a stable cell line that contains the λ neo-chromosome [see, Fig. 2D; see, also U.S. Patent No. 5,288,625], produced by E-type amplifications. KE1-2/4 has been transfected with vectors containing λ DNA, selectable markers, such as the puromycin-resistance gene, and genes of interest, such as p53 and the anti-HIV ribozyme gene. These vectors target the gene of interest into the λ neo-chromosome by virtue of homologous recombination with the heterologous DNA in the chromosome.

-120-

3. C5pMCT53 Cells

The EC3/7C5 cell line has been co-transfected with pH132, pCH110 and λ DNA [see, EXAMPLE 2] as well as other constructs. Various clones and subclones have been selected. For example

- 5 transformation with a construct that includes p53 encoding DNA, produced cells designated C5pMCT53.

4. TF1004G24 Cells

- As discussed above, cotransfection of EC3/7C5 cells with plasmids [pH132, pCH110 available from Pharmacia, see, also Hall *et al.* (1983) *J. Mol. Appl. Gen.* 2:101-109] and with λ DNA [λ cl 875 Sam 7 (New England Biolabs)] produced transformed cells. Among these is TF1004G24, which contains the DNA encoding the anti-HIV ribozyme in the neo-minichromosome. Recloning of TF1004G24 produced numerous cell lines. Among these is the NHHL24 cell line. This cell line also has
- 10 the anti-HIV ribozyme in the neo-minichromosome and expresses high levels of β -gal. It has been fused with CHO-K20 cells to produce various hybrids.

5. TF1004G19-Derived cells

- Recloning and selection of the TF1004G transformants produced
- 20 the cell line TF1004G19, discussed above in EXAMPLE 4, which contains the unstable sausage chromosome and the neo-minichromosome. Single cell cloning produced the TF1004G-19C5 [see Figure 4] cell line, which has a stable sausage chromosome and the neo-minichromosome. TF1004G-19C5 has been fused with CHO cells and
- 25 the hybrids grown under selective conditions to produce the 19C5xHa4 and 19C5xHa3 cell lines [see, EXAMPLE 4] and others. Recloning of the 19C5xHa3 cell line yielded a cell line containing a gigachromosome, i.e., cell line 19C5xHa47, see Figure 2E. BrdU treatment of 19C5xHa4 cells and growth under selective conditions [neomycin (G) and/or hygromycin

-121-

(H)] has produced hybrid cell lines such as the G3D5 and G4D6 cell lines and others. G3D5 has the neo-minichromosome and the megachromosome. G4D6 has only the neo-minichromosome.

- Recloning of 19C5xHa4 cells in H medium produced numerous clones. Among these is H1D3 [see Figure 4], which has the stable megachromosome. Repeated BrdU treatment and recloning of H1D3 cells has produced the HB31 cell line, which has been used for transformations with the pTEMPUD, pTEMPU, pTEMPU3, and pCEPUR-132 vectors [see, Examples 12 and 14, below].
- 10 H1D3 has been fused with a CD4⁺ Hela cell line that carries DNA encoding CD4 and neomycin resistance on a plasmid [see, e.g., U.S. Patent Nos. 5,413,914, 5,409,810, 5,266,600, 5,223,263, 5,215,914 and 5,144,019, which describe these Hela cells]. Selection with GH has produced hybrids, including H1xHE41 [see Figure 4], which carries the
- 15 megachromosome and also a single human chromosome that includes the CD4neo construct. Repeated BrdU treatment and single cell cloning has produced cell lines with the megachromosome [cell line 1B3, see Figure 4]. About 25% of the 1B3 cells have a truncated megachromosome [~ 90-120 Mb]. Another of these subclones,
- 20 designated 2C5, was cultured on hygromycin-containing medium and megachromosome-free cell lines were obtained and grown in G418-containing medium. Recloning of these cells yielded cell lines such as 1B4 and others that have a dwarf megachromosome [~ 150-200 Mb], and cell lines, such as 11C3 and mM2C1, which have a micro-
- 25 megachromosome [~ 50-90 Mb]. The micro-megachromosome of cell

-122-

line mM2C1 has no telomeres; however, if desired, synthetic telomeres, such as those described and generated herein, may be added to the mM2C1 cell micro-megachromosomes. Cell lines containing smaller truncated megachromosomes, such as the mM2C1 cell line containing
5 the micro-megachromosome, can be used to generate even smaller megachromosomes, e.g., ~10-30 Mb in size. This may be accomplished, for example, by breakage and fragmentation of the micro-megachromosome in these cells through exposing the cells to X-ray irradiation, BrdU or telomere-directed in vivo chromosome fragmentation.

10

EXAMPLE 8

Replication of the megachromosome

The homogeneous architecture of the megachromosomes provides a unique opportunity to perform a detailed analysis of the replication of the
15 constitutive heterochromatin.

A. Materials and methods

1. Culture of cell lines

H1D3 mouse-hamster hybrid cells carrying the megachromosome [see, EXAMPLE 4] were cultured in F-12 medium containing 10% fetal
20 calf serum [FCS] and 400 μ g/ml Hygromycin B [Calbiochem]. G3D5 hybrid cells [see, Example 4] were maintained in F-12 medium containing 10% FCS, 400 μ g/ml Hygromycin B (Calbiochem), and 400 μ g/ml G418 [SIGMA]. Mouse A9 fibroblast cells were cultured in F-12 medium supplemented with 10% FCS.

25 2. BrdU labelling

In typical experiments, 20-24 parallel semi-confluent cell cultures were set up in 10 cm Petri dishes. Bromodeoxyuridine (BrdU) (Fluka) was dissolved in distilled water alkalized with a drop of NaOH, to make a 10^{-2} M stock solution. Aliquots of 10-50 μ l of this BrdU stock solution

-123-

were added to each 10 ml culture, to give a final BrdU concentration of 10-50 μ M. The cells were cultured in the presence of BrdU for 30 min, and then washed with warm complete medium, and incubated without BrdU until required. At this point, 5 μ g/ml colchicine was added to a sample culture every 1 or 2 h. After 1-2 h colchicine treatment, mitotic cells were collected by "shake-off" and regular chromosome preparations were made for immunolabelling.

3. Immunolabelling of chromosomes and *in situ* hybridization

Immunolabelling with fluorescein-conjugated anti-BrdU monoclonal antibody (Boehringer) was done according to the manufacturer's recommendations, except that for mouse A9 chromosomes, 2 M hydrochloric acid was used at 37° C for 25 min, while for chromosomes of hybrid cells, 1 M hydrochloric acid was used at 37° C for 30 min. *In situ* hybridization with biotin-labelled probes, and indirect immunofluorescence and *in situ* hybridization on the same preparation, were performed as described previously [Hadlaczký *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110, see, also U.S. Patent No. 5,288,625].

4. Microscopy

All observations and microphotography were made by using a Vanox AHBS (Olympus) microscope. Fujicolor 400 Super G or Fujicolor 1600 Super HG high-speed colour negatives were used for photographs.

B. Results

The replication of the megachromosome was analyzed by BrdU pulse labelling followed by immunolabelling. The basic parameters for DNA labelling *in vivo* were first established. Using a 30-min pulse of 50 μ M BrdU in parallel cultures, samples were taken and fixed at 5 min intervals from the beginning of the pulse, and every 15 min up to 1 h after the removal of BrdU. Incorporated BrdU was detected by

-124-

immunolabelling with fluorescein-conjugated anti-BrdU monoclonal antibody. At the first time point (5 min) 38% of the nuclei were labelled, and a gradual increase in the number of labelled nuclei was observed during incubation in the presence of BrdU, culminating in 46% in the 30-
5 min sample, at the time of the removal of BrdU. At further time points (60, 75, and 90 min) no significant changes were observed, and the fraction of labelled nuclei remained constant [44.5-46%].

These results indicate that (i) the incorporation of the BrdU is a rapid process, (ii) the 30 min pulse-time is sufficient for reliable labelling
10 of S-phase nuclei, and (iii) the BrdU can be effectively removed from the cultures by washing.

The length of the cell cycle of the H1D3 and G3D5 cells was estimated by measuring the time between the appearance of the earliest BrdU signals on the extreme late replicating chromosome segments and
15 the appearance of the same pattern only on one of the chromatids of the chromosomes after one completed cell cycle. The length of G2 period was determined by the time of the first detectable BrdU signal on prophase chromosomes and by the labelled mitoses method [Qastler et al. (1959) Exp. Cell Res. 17:420-438]. The length of the S-phase was
20 determined in three ways: (i) on the basis of the length of cell cycle and the fraction of nuclei labelled during the 30-120 min pulse; (ii) by measuring the time between the very end of the replication of the extreme late replicating chromosomes and the detection of the first signal on the chromosomes at the beginning of S phase; (iii) by the labelled
25 mitoses method. In repeated experiments, the duration of the cell cycle was found to be 22-26 h, the S phase 10-14 h, and the G2 phase 3.5-4.5 h.

-125-

Analyses of the replication of the megachromosome were made in parallel cultures by collecting mitotic cells at two hour intervals following two hours of colchicine treatment. In a repeat experiment, the same analysis was performed using one hour sample intervals and one hour colchicine treatment. Although the two procedures gave comparable results, the two hour sample intervals were viewed as more appropriate since approximately 30% of the cells were found to have a considerably shorter or longer cell cycle than the average. The characteristic replication patterns of the individual chromosomes, especially some of the late replicating hamster chromosomes, served as useful internal markers for the different stages of S-phase. To minimize the error caused by the different lengths of cell cycles in the different experiments, samples were taken and analyzed throughout the whole cell cycle until the appearance of the first signals on one chromatid at the beginning of the second S-phase.

The sequence of replication in the megachromosome is as follows. At the very beginning of the S-phase, the replication of the megachromosome starts at the ends of the chromosomes. The first initiation of replication in an interstitial position can usually be detected at the centromeric region. Soon after, but still in the first quarter of the S-phase, when the terminal region of the short arm has almost completed its replication, discrete initiation signals appear along the chromosome arms. In the second quarter of the S-phase, as replication proceeds, the BrdU-labelled zones gradually widen, and the checkered pattern of the megachromosome becomes clear [see, e.g., Fig. 2F]. At the same time, pericentric regions of mouse chromosomes also show intense incorporation of BrdU. The replication of the megachromosome peaks at the end of the second quarter and in the third quarter of the S-phase. At the end of the third quarter, and at the very beginning of the last quarter

-126-

of the S-phase, the megachromosome and the pericentric heterochromatin of the mouse chromosomes complete their replication. By the end of S-phase, only the very late replicating segments of mouse and hamster chromosomes are still incorporating BrdU.

- 5 The replication of the whole genome occurs in distinct phases. The signal of incorporated BrdU increased continuously until the end of the first half of the S-phase, but at the beginning of the third quarter of the S-phase chromosome segments other than the heterochromatic regions hardly incorporated BrdU. In the last quarter of the S-phase, the
- 10 BrdU signals increased again when the extreme late replicating segments showed very intense incorporation.

- Similar analyses of the replication in mouse A9 cells were performed as controls. To increase the resolution of the immunolabelling pattern, pericentric regions of A9 chromosomes were decondensed by
- 15 treatment with Hoechst 33258. Because of the intense replication of the surrounding euchromatic sequences, precise localization of the initial BrdU signal in the heterochromatin was normally difficult, even on undercondensed mouse chromosomes. On those chromosomes where the initiation signal(s) were localized unambiguously, the replication of
- 20 the pericentric heterochromatin of A9 chromosomes was similar to that of the megachromosome. Chromosomes of A9 cells also exhibited replication patterns and sequences similar to those of the mouse chromosomes in the hybrid cells. These results indicate that the replicators of the megachromosome and mouse chromosomes retained
- 25 their original timing and specificity in the hybrid cells.

By comparing the pattern of the initiation sites obtained after BrdU incorporation with the location of the integration sites of the "foreign" DNA in a detailed analysis of the first quarter of the S-phase, an attempt was made to identify origins of replication (initiation sites) in relation to

-127-

the amplicon structure of the megachromosome. The double band of integrated DNA on the long arm of the megachromosome served as a cytological marker. The results showed a colocalization of the BrdU and *in situ* hybridization signals found at the cytological level, indicating that the "foreign" DNA sequences are in close proximity to the origins of replication, presumably integrated into the non-satellite sequences between the replicator and the satellite sequences [see, Figure 3]. As described in Example 6.B.4, the rDNA sequences detected in the megachromosome are also localized at the amplicon borders at the site of integration of the "foreign" DNA sequences, suggesting that the origins of replication responsible for initiation of replication of the megachromosome involve rDNA sequences. In the pericentric region of several other chromosomes, dot-like BrdU signals can also be observed that are comparable to the initiation signals on the megachromosome. These signals may represent similar initiation sites in the heterochromatic regions of normal chromosomes.

At a frequency of 10^{-4} , "uncontrolled" amplification of the integrated DNA sequences was observed in the megachromosome. Consistent with the assumption (above) that "foreign" sequences are in proximity of the replicators, this spatially restricted amplification is likely to be a consequence of uncontrolled repeated firings of the replication origin(s) without completing the replication of the whole segment.

C. Discussion

It has generally been thought that the constitutive heterochromatin of the pericentric regions of chromosomes is late replicating [see, e.g., Miller (1976) Chromosoma 55:165-170]. On the contrary, these experiments evidence that the replication of the heterochromatic blocks starts at a discrete initiation site in the first half of the S-phase and continues through approximately three-quarters of S-phase. This

-128-

difference can be explained in the following ways: (i) in normal chromosomes, actively replicating euchromatic sequences that surround the satellite DNA obscure the initiation signals, and thus the precise localization of initiation sites is obscured; (ii) replication of the

5 heterochromatin can only be detected unambiguously in a period during the second half of the S-phase, when the bulk of the heterochromatin replicates and most other chromosomal regions have already completed their replication, or have not yet started it. Thus, low resolution

10 cytological techniques, such as analysis of incorporation of radioactively labelled precursors by autoradiography, only detect prominent replication signals in the heterochromatin in the second half of S-phase, when adjacent euchromatic segments are no longer replicating.

In the megachromosome, the primary initiation sites of replication colocalize with the sites where the "foreign" DNA sequences and rDNA

15 sequences are integrated at the amplicon borders. Similar initiation signals were observed at the same time in the pericentric heterochromatin of some of the mouse chromosomes that do not have "foreign" DNA, indicating that the replication initiation sites at the borders of amplicons may reside in the non-satellite flanking sequences

20 of the satellite DNA blocks. The presence of a primary initiation site at each satellite DNA doublet implies that this large chromosome segment is a single huge unit of replication [megareplicon] delimited by the primary initiation site and the termination point at each end of the unit. Several lines of evidence indicate that, within this higher-order replication unit,

25 "secondary" origins and replicons contribute to the complete replication of the megareplicon:

1. The total replication time of the heterochromatic regions of the megachromosome was ~9-11 h. At the rate of movement of replication forks, 0.5-5 kb per minute, that is typical of eukaryotic

-129-

chromosomes [Kornberg et al. (1992) *DNA Replication*. 2nd. ed., New York: W.H. Freeman and Co, p. 474], replication of a ~ 15 Mb replicon would require 50-500 h. Alternatively, if only a single replication origin was used, the average replication speed would have to be 25 kb per
5 minute to complete replication within 10 h. By comparing the intensity of the BrdU signals on the euchromatic and the heterochromatic chromosome segments, no evidence for a 5- to 50-fold difference in their replication speed was found.

2. Using short BrdU pulse labelling, a single origin of replication
10 would produce a replication band that moves along the replicon, reflecting the movement of the replication fork. In contrast, a widening of the replication zone that finally gave rise to the checkered pattern of the megachromosome was observed, and within the replication period, the most intensive BrdU incorporation occurred in the second half of the
15 S-phase. This suggests that once the megareplicator has been activated, it permits the activation and firing of "secondary" origins, and that the replication of the bulk of the satellite DNA takes place from these "secondary" origins during the second half of the S-phase. This is supported by the observation that in certain stages of the replication of
20 the megachromosome, the whole amplicon can apparently be labelled by a short BrdU pulse.

Megareplicators and secondary replication origins seem to be under strict temporal and spatial control. The first initiation within the megachromosomes usually occurred at the centromere, and shortly
25 afterward all the megareplicators become active. The last segment of the megachromosome to complete replication was usually the second segment of the long arm. Results of control experiments with mouse A9 chromosomes indicate that replication of the heterochromatin of mouse chromosomes corresponds to the replication of the megachromosome

-130-

amplicons. Therefore, the pre-existing temporal control of replication in the heterochromatic blocks is preserved in the megachromosome.

Positive [Hassan *et al.* (1994) *J. Cell. Sci.* 107:425-434] and negative [Haase *et al.* (1994) *Mol. Cell. Biol.* 14:2516-2524] correlations

- 5 between transcriptional activity and initiation of replication have been proposed. In the megachromosome, transcription of the integrated genes seems to have no effect on the original timing of the replication origins. The concerted, precise timing of the megareplicator initiations in the different amplicons suggests the presence of specific, cis-acting
- 10 sequences, origins of replication.

- Considering that pericentric heterochromatin of mouse chromosomes contains thousands of short, simple repeats spanning 7-15 Mb, and the centromere itself may also contain hundreds of kilobases, the existence of a higher-order unit of replication seems probable. The
- 15 observed uncontrolled intrachromosomal amplification restricted to a replication initiation region of the megachromosome is highly suggestive of a rolling-circle type amplification, and provides additional evidence for the presence of a replication origin in this region.

- The finding that a specific replication initiation site occurs at the
- 20 boundaries of amplicons suggests that replication might play a role in the amplification process. These results suggest that each amplicon of the megachromosome can be regarded as a huge megareplicon defined by a primary initiation site [megareplicator] containing "secondary" origins of replication. Fusion of replication bubbles from different origins of bi-
- 25 directional replication [DePamphilis (1993) *Ann. Rev. Biochem.* 62:29-63] within the megareplicon could form a giant replication bubble, which would correspond to the whole megareplicon. In the light of this, the formation of megabase-size amplicons can be accommodated by a replication-directed amplification mechanism. In H and E-type

-131-

amplifications, intrachromosomal multiplication of the amplicons was observed [see, above EXAMPLES], which is consistent with the unequal sister chromatid exchange model. Induced or spontaneous unscheduled replication of a megareplicon in the constitutive heterochromatin may
5 also form new amplicon(s) leading to the expansion of the amplification or to the heterochromatic polymorphism of "normal" chromosomes. The "restoration" of the missing segment on the long arm of the megachromosome may well be the result of the re-replication of one amplicon limited to one strand.

10 Taken together, without being bound by any theory, a replication-directed mechanism is a plausible explanation for the initiation of large-scale amplifications in the centromeric regions of mouse chromosomes, as well as for the *de novo* chromosome formations. If specific [amplifier, i.e., sequences controlling amplification] sequences play a role in
15 promoting the amplification process, sequences at the primary replication initiation site [megareplicator] of the megareplicon are possible candidates.

The presence of rRNA gene sequence at the amplicon borders near the foreign DNA in the megachromosome suggests that this sequence
20 contributes to the primary replication initiation site and participates in large-scale amplification of the pericentric heterochromatin in *de novo* formation of SATACs. Ribosomal RNA genes have an intrinsic amplification mechanism that provides for multiple copies of tandem genes. Thus, for purposes herein, in the construction of SATACs in
25 cells, rDNA will serve as a region for targeted integration, and as components of SATACs constructed in vitro.

-132-

EXAMPLE 9**Generation of chromosomes with amplified regions derived from mouse chromosome 1**

To show that the events described in EXAMPLES 2-7 are not
5 unique to mouse chromosome 7 and to show that the EC7/3 cell line is not required for formation of the artificial chromosomes, the experiments have been repeated using different initial cell lines and DNA fragments. Any cell or cell line should be amenable to use or can readily be determined that it is not.

10 A. Materials

The LP11 cell line was produced by the "scrape-loading" transfection method [Fechheimer et al. (1987) Proc. Natl. Acad. Sci. U.S.A. 84:8463-8467] using 25 μ g plasmid DNA for 5×10^6 recipient cells. LP11 cells were maintained in F-12 medium containing 3-15 μ g/ml
15 Puromycin [SIGMA].

B. Amplification in LP11 cells

The large-scale amplification described in the above Examples is not restricted to the transformed EC3/7 cell line or to the chromosome 7 of mouse. In an independent transformation experiment, LMTK⁻ cells
20 were transfected using the calcium phosphate precipitation procedure with a selectable puromycin-resistance gene-containing construct designated pPuroTel [see Example 1.E.2. for a description of this plasmid], to establish cell line LP11. Cell line LP11 carries chromosome(s) with amplified chromosome segments of different lengths (~150-600 Mb).
25 Cytological analysis of the LP11 cells indicated that the amplification occurred in the pericentric region of the long arm of a submetacentric chromosome formed by Robertsonian translocation. This chromosome arm was identified by G-banding as chromosome 1. C-banding and *in situ* hybridization with mouse major satellite DNA probe showed that an

-133-

E-type amplification had occurred: the newly formed region was composed of an array of euchromatic chromosome segments containing different amounts of heterochromatin. The size and C-band pattern of the amplified segments were heterogeneous. In several cells, the number
5 of these amplified units exceeded 50; single-cell subclones of LP11 cell lines, however, carry stable marker chromosomes with 10-15 segments and constant C-band patterns.

Sublines of the thymidine kinase-deficient LP11 cells (e.g., LP11-15P 1C5/7 cell line) established by single-cell cloning of LP11 cells were
10 transfected with a thymidine kinase gene construct. Stable TK⁺ transfectants were established.

EXAMPLE 10

Isolation of SATACS and other chromosomes with atypical base content and/or size

15 I. Isolation of artificial chromosomes from endogenous chromosomes

Artificial chromosomes, such as SATACs, may be sorted from endogenous chromosomes using any suitable procedures, and typically involve isolating metaphase chromosomes, distinguishing the artificial
20 chromosomes from the endogenous chromosomes, and separating the artificial chromosomes from endogenous chromosomes. Such procedures will generally include the following basic steps: (1) culture of a sufficient number of cells (typically about 2×10^7 mitotic cells) to yield, preferably on the order of 1×10^6 artificial chromosomes, (2) arrest of
25 the cell cycle of the cells in a stage of mitosis, preferably metaphase, using a mitotic arrest agent such as colchicine, (3) treatment of the cells, particularly by swelling of the cells in hypotonic buffer, to increase susceptibility of the cells to disruption, (4) by application of physical force to disrupt the cells in the presence of isolation buffers for
30 stabilization of the released chromosomes, (5) dispersal of chromosomes

-134-

- in the presence of isolation buffers for stabilization of free chromosomes, (6) separation of artificial from endogenous chromosomes and (7) storage (and shipping if desired) of the isolated artificial chromosomes in appropriate buffers. Modifications and variations of the general
- 5 procedure for isolation of artificial chromosomes, for example to accommodate different cell types with differing growth characteristics and requirements and to optimize the duration of mitotic block with arresting agents to obtain the desired balance of chromosome yield and level of debris, may be empirically determined.
- 10 Steps 1-5 relate to isolation of metaphase chromosomes. The separation of artificial from endogenous chromosomes (step 6) may be accomplished in a variety of ways. For example, the chromosomes may be stained with DNA-specific dyes such as Hoeschst 33258 and chromomycin A₃ and sorted into artificial and endogenous chromosomes
- 15 on the basis of dye content by employing fluorescence-activated cell sorting (FACS). To facilitate larger scale isolation of the artificial chromosomes, different separation techniques may be employed such as swinging bucket centrifugation (to effect separation based on chromosome size and density) [see, e.g., Mendelsohn *et al.* (1968) J. Mol. Biol. 32:101-108], zonal rotor centrifugation (to effect separation on
- 20 the basis of chromosome size and density) [see, e.g., Burki *et al.* (1973) Prep. Biochem. 3:157-182; Stubblefield *et al.* (1978) Biochem. Biophys. Res. Commun. 83:1404-1414, velocity sedimentation (to effect separation on the basis of chromosome size and shape) [see e.g., Collard
- 25 *et al.* (1984) Cytometry 5:9-19]. Immuno-affinity purification may also be employed in larger scale artificial chromosome isolation procedures. In this process, large populations of artificial chromosome-containing cells (asynchronous or mitotically enriched) are harvested en masse and the mitotic chromosomes (which can be released from the cells using

-135-

standard procedures such as by incubation of the cells in hypotonic buffer and/or detergent treatment of the cells in conjunction with physical disruption of the treated cells) are enriched by binding to antibodies that are bound to solid state matrices (e.g. column resins or magnetic beads). Antibodies suitable for use in this procedure bind to condensed centromeric proteins or condensed and DNA-bound histone proteins. For example, autoantibody LU851 (see Hadlaczky *et al.* (1989) Chromosoma 97:282-288), which recognizes mammalian centromeres may be used for large-scale isolation of chromosomes prior to subsequent separation of artificial from endogenous chromosomes using methods such as FACS. The bound chromosomes would be washed and eventually eluted for sorting. Immunoaffinity purification may also be used directly to separate artificial chromosomes from endogenous chromosomes. For example, SATACs may be generated in or transferred to (e.g., by microinjection or microcell fusion as described herein) a cell line that has chromosomes that contain relatively small amounts of heterochromatin, such as hamster cells (e.g., V79 cells or CHO-K1 cells). The SATACs, which are predominantly heterochromatin, are then separated from the endogenous chromosomes by utilizing anti-heterochromatin binding protein (*Drosophila* HP-1) antibody conjugated to a solid matrix. Such matrix preferentially binds SATACs relative to hamster chromosomes. Unbound hamster chromosomes are washed away from the matrix and the SATACs are eluted by standard techniques.

A. Cell lines and cell culturing procedures

In one isolation procedure, 1B3 mouse-hamster-human hybrid cells [see, Figure 4] carrying the megachromosome or the truncated megachromosome were grown in F-12 medium supplemented with 10% fetal calf serum, 150 μ g/ml hygromycin B and 400 μ g/ml G418. GHB42 [a cell line recloned from G3D5 cells] mouse-hamster hybrid cells carrying

-136-

the megachromosome and the minichromosome were also cultured in F-12 medium containing 10% fetal calf serum, 150 μ g/ml hygromycin B and 400 μ g/ml G418. The doubling time of both cell lines was about 24-40 hours, typically about 32 hours.

- 5 Typically, cell monolayers are passaged when they reach about 60-80% confluence and are split every 48-72 hours. Cells that reach greater than 80% confluence senesce in culture and are not preferred for chromosome harvesting. Cells may be plated in 100-200 100-mm dishes at about 50-70% confluency 12-30 hours before mitotic arrest (see,
10 below).

- Other cell lines that may be used as hosts for artificial chromosomes and from which the artificial chromosomes may be isolated include, but are not limited to, PtK1 (NBL-3) marsupial kidney cells (ATCC accession no. CCL35), CHO-K1 Chinese hamster ovary cells (ATCC accession
15 no. CCL61), V79-4 Chinese hamster lung cells (ATCC accession no. CCL93), Indian muntjac skin cells (ATCC accession no. CCL157), LMTK(-) thymidine kinase deficient murine L cells (ATCC accession no. CCL1.3), Sf9 fall armyworm (*Spodoptera frugiperda*) ovary cells (ATCC accession no. CRL 1711) and any generated heterokaryon (hybrid) cell
20 lines, such as, for example, the hamster-murine hybrid cells described herein, that may be used to construct MACs, particularly SATACs.

- Cell lines may be selected, for example, to enhance efficiency of artificial chromosome production and isolation as may be desired in large-scale production processes. For instance, one consideration in selecting
25 host cells may be the artificial chromosome-to-total chromosome ratio of the cells. To facilitate separation of artificial chromosomes from endogenous chromosomes, a higher artificial chromosome-to-total chromosome ratio might be desirable. For example, for H1D3 cells (a murine/hamster heterokaryon; see Figure 4), this ratio is 1:50, i.e., one

-137-

artificial chromosome (the megachromosome) to 50 total chromosomes. In contrast, Indian muntjac skin cells (ATCC accession no. CCL157) contain a smaller total number of chromosomes (a diploid number of chromosomes of 7), as do kangaroo rat cells (a diploid number of
5 chromosomes of 12) which would provide for a higher artificial chromosome-to-total chromosome ratio upon introduction of, or generation of, artificial chromosomes in the cells.

Another consideration in selecting host cells for production and isolation of artificial chromosomes may be size of the endogenous
10 chromosomes as compared to that of the artificial chromosomes. Size differences of the chromosomes may be exploited to facilitate separation of artificial chromosomes from endogenous chromosomes. For example, because Indian muntjac skin cell chromosomes are considerably larger than minichromosomes and truncated megachromosomes, separation of
15 the artificial chromosome from the muntjac chromosomes may possibly be accomplished using univariate (one dye, either Hoechst 33258 or Chromomycin A3) FACS separation procedures.

Another consideration in selecting host cells for production and isolation of artificial chromosomes may be the doubling time of the cells.
20 For example, the amount of time required to generate a sufficient number of artificial chromosome-containing cells for use in procedures to isolate artificial chromosomes may be of significance for large-scale production. Thus, host cells with shorter doubling times may be desirable. For instance, the doubling time of V79 hamster lung cells is about 9-10 hours
25 in comparison to the approximately 32-hour doubling time of H1D3 cells.

Accordingly, several considerations may go into the selection of host cells for the production and isolation of artificial chromosomes. It may be that the host cell selected as the most desirable for de novo formation of artificial chromosomes is not optimized for large-scale

-138-

production of the artificial chromosomes generated in the cell line. In such cases, it may be possible, once the artificial chromosome has been generated in the initial host cell line, to transfer it to a production cell line more well suited to efficient, high-level production and isolation of the artificial chromosome. Such transfer may be accomplished through several methods, for example through microcell fusion, as described herein, or microinjection into the production cell line of artificial chromosomes purified from the generating cell line using procedures such as described herein. Production cell lines preferably contain two or more copies of the artificial artificial chromosome per cell.

B. Chromosome isolation

In general, cells are typically cultured for two generations at exponential growth prior to mitotic arrest. To accumulate mitotic 1B3 and GHB42 cells in one particular isolation procedure, 5 μ g/ml colchicine was added for 12 hours to the cultures. The mitotic index obtained was 60-80%. The mitotic cells were harvested by selective detachment by gentle pipetting of the medium on the monolayer cells. It is also possible to utilize mechanical shake-off as a means of releasing the rounded-up (mitotic) cells from the plate. The cells were sedimented by centrifugation at 200 x g for 10 minutes.

Cells (grown on plastic or in suspension) may be arrested in different stages of the cell cycle with chemical agents other than colchicine, such as hydroxyurea, vinblastine, colcemid or aphidicolin. Chemical agents that arrest the cells in stages other than mitosis, such as hydroxyurea and aphidicolin, are used to synchronize the cycles of all cells in the population and then are removed from the cell medium to allow the cells to proceed, more or less simultaneously, to mitosis at which time they may be harvested to disperse the chromosomes. Mitotic cells could be enriched for a mechanical shake-off (adherent cells). The

-139-

cell cycles of cells within a population of MAC-containing cells may also be synchronized by nutrient, growth factor or hormone deprivation which leads to an accumulation of cells in the G₁ or G₀ stage; readdition of nutrients or growth factors then allows the quiescent cells to re-enter the

5 the cell cycle in synchrony for about one generation. Cell lines that are known to respond to hormone deprivation in this manner, and which are suitable as hosts for artificial chromosomes, include the Nb2 rat lymphoma cell line which is absolutely dependent on prolactin for stimulation of proliferation (see Gout et al. (1980) Cancer Res. 40:2433-

10 2436). Culturing the cells in prolactin-deficient medium for 18-24 hours leads to arrest of proliferation, with cells accumulating early in the G₁ phase of the cell cycle. Upon addition of prolactin, all the cells progress through the cell cycle until M phase at which point greater than 90% of the cells would be in mitosis (addition of colchicine could increase the

15 amount of the mitotic cells to greater than 95%). The time between reestablishing proliferation by prolactin addition and harvesting mitotic cells for chromosome separation may be empirically determined.

Alternatively, adherent cells, such as V79 cells, may be grown in roller bottles and mitotic cells released from the plastic surface by

20 rotating the roller bottles at 200 rpm or greater (Shwarchuk et al. (1993) Int. J. Radiat. Biol. 64:601-612). At any given time, approximately 1% of the cells in an exponentially growing asynchronous population is in M-phase. Even without the addition of colchicine, 2×10^7 mitotic cells have been harvested from four 1750-cm² roller bottles after a 5-min spin

25 at 200 rpm. Addition of colchicine for 2 hours may increase the yield to 6×10^8 mitotic cells.

Several procedures may be used to isolate metaphase chromosomes from these cells, including, but not limited to, one based on a polyamine buffer system [Cram et al. (1990) Methods in Cell

-140-

- Biology 33:377-382], one on a modified hexylene glycol buffer system [Hadlaczky et al. (1982) Chromosoma 86:643-65], one on a magnesium sulfate buffer system [Van den Engh et al. (1988) Cytometry 9:266-270 and Van den Engh et al. (1984) Cytometry 5:108], one on an acetic acid
- 5 fixation buffer system [Stoehr et al. (1982) Histochemistry 74:57-61], and one on a technique utilizing hypotonic KCl and propidium iodide [Cram et al. (1994) XVII meeting of the International Society for Analytical Cytology, October 16-21, Tutorial IV Chromosome Analysis and Sorting with Commercial Flow Cytometers; Cram et al. (1990)
- 10 Methods in Cell Biology 33:376].

1. Polyamine procedure

- In the polyamine procedure that was used in isolating artificial chromosomes from either 1B3 or GHB42 cells, about 10^7 mitotic cells were incubated in 10 ml hypotonic buffer (75 mM KCl, 0.2 mM
- 15 spermine, 0.5 mM spermidine) for 10 minutes at room temperature to swell the cells. The cells are swollen in hypotonic buffer to loosen the metaphase chromosomes but not to the point of cell lysis. The cells were then centrifuged at 100 x g for 8 minutes, typically at room temperature. The cell pellet was drained carefully and about 10^7 cells
- 20 were resuspended in 1 ml polyamine buffer [15 mM Tris-HCl, 20 mM NaCl, 80 mM KCl, 2 mM EDTA, 0.5 mM EGTA, 14 mM β -mercaptoethanol, 0.1% digitonin, 0.2 mM Spermine, 0.5 mM spermidine] for physical dispersal of the metaphase chromosomes. Chromosomes were then released by gently drawing the cell suspension up and expelling it
- 25 through a 22 G needle attached to a 3 ml plastic syringe. The chromosome concentration was about $1-3 \times 10^8$ chromosomes/ml.

The polyamine buffer isolation protocol is well suited for obtaining high molecular weight chromosomal DNA [Sillar and Young (1981) J. Histochem. Cytochem. 29:74-78; VanDilla et al. (1986) Biotechnology

-141-

4:537-552; Bartholdi et al. (1988) In "Molecular Genetics of Mammalian Cells" (M.Goettsman, ed.), Methods in Enzymology 151:252-267. Academic Press, Orlando]. The chromosome stabilizing buffer uses the polyamines spermine and spermidine to stabilize chromosome structure
5 [Blumenthal et al. (1979)J. Cell Biol. 81:255-259; Lalande et al. (1985) Cancer Genet. Cytogenet. 23:151-157] and heavy metals chelators to reduce nuclease activity.

The polyamine buffer protocol has wide applicability, however, as with other protocols, the following variables must be optimized for each
10 cell type: blocking time, cell concentration, type of hypotonic swelling buffer, swelling time, volume of hypotonic buffer, and vortexing time. Chromosomes prepared using this protocol are typically highly condensed.

There are several hypotonic buffers that may be used to swell the
15 cells, for example buffers such as the following: 75 mM KCl; 75 mM KCl, 0.2 mM spermine, 0.5 mM spermidine; Ohnuki's buffer of 16.2 mM sodium nitrate, 6.5 mM sodium acetate, 32.4 mM KCl [Ohnuki (1965) Nature 208:916-917 and Ohnuki (1968) Chromosoma 25:402-428]; and
20 a variation of Ohnuki's buffer that additionally contains 0.2 mM spermine and 0.5 mM spermidine. The amount and hypotonicity of added buffer vary depending on cell type and cell concentration. Amounts may range from 2.5 - 5.5 ml per 10^7 cells or more. Swelling times may vary from 10-90 minutes depending on cell type and which swelling buffer is used.

The composition of the polyamine isolation buffer may also be
25 varied. For example, one modified buffer contains 15 mM Tris-HCl, pH 7.2, 70 mM NaCl, 80 mM KCl, 2 mM EDTA, 0.5 mM EGTA, 14 mM beta-mercaptoethanol, 0.25% Triton-X, 0.2 mM spermine and 0.5 mM spermidine.

-142-

Chromosomal dispersal may also be accomplished by a variety of physical means. For example, cell suspension may be gently drawn up and expelled in a 3-ml syringe fitted with a 22-gauge needle [Cram et al. (1990) Methods in Cell Biology 33:377-382], cell suspension may be
5 agitated on a bench-top vortex [Cram et al. (1990) Methods in Cell Biology 33:377-382], cell suspension may be disrupted with a homogenizer [Sillar and Young (1981) J. Histochem. Cytochem. 29:74-78; Carrano et al. (1979) Proc. Natl. Acad. Sci. U.S.A. 76:1382-1384] and cell suspension may be disrupted with a bench-top ultrasonic bath
10 [Stoehr et al. (1982) Histochemistry 74:57-61].

2. Hexylene glycol buffer system

In the hexylene glycol buffer procedure that was used in isolating artificial chromosomes from either 1B3 or GHB42 cells, about 8×10^6 mitotic cells were resuspended in 10 ml glycine-hexylene glycol buffer
15 [100 mM glycine, 1% hexylene glycol, pH 8.4-8.6 adjusted with saturated Ca-hydroxide solution] and incubated for 10 minutes at 37°C, followed by centrifugation for 10 minutes to pellet the nuclei. The supernatant was centrifuged again at 200 x g for 20 minutes to pellet the chromosomes. Chromosomes were resuspended in isolation buffer
20 ($1-3 \times 10^8$ chromosomes/ml).

The hexylene glycol buffer composition may also be modified. For example, one modified buffer contains 25 mM Tris-HCl, pH 7.2, 750 mM hexylene glycol, 0.5 mM CaCl_2 , 1.0 mM MgCl_2 [Carrano et al. (1979) Proc. Natl. Acad. Sci. U.S.A. 76:1382-1384].

25 3. Magnesium-sulfate buffer system

This buffer system may be used with any of the methods of cell swelling and chromosomal dispersal, such as described above in connection with the polyamine and hexylene glycol buffer systems. In this procedure, mitotic cells are resuspended in the following buffer: 4.8

-143-

mM HEPES, pH 8.0, 9.8 mM MgSO₄, 48 mM KCl, 2.9 mM dithiothreitol [Van den Engh *et al.* (1985) Cytometry 6:92 and Van den Engh *et al.* (1984) Cytometry 5:108].

4. Acetic acid fixation buffer system

- 5 This buffer system may be used with any of the methods of cell swelling and chromosomal dispersal, such as described above in connection with the polyamine and hexylene glycol buffer systems. In this procedure, mitotic cells are resuspended in the following buffer: 25 mM Tris-HCl, pH 3.2, 750 mM (1,6)-hexandiol, 0.5 mM CaCl₂, 1.0%
10 acetic acid [Stoehr *et al.* (1982) Histochemistry 74:57-61].

5. KCl-propidium iodide buffer system

- This buffer system may be used with any of the methods of cell swelling and chromosomal dispersal, such as described above in connection with the polyamine and hexylene glycol buffer systems. In
15 this procedure, mitotic cells are resuspended in the following buffer: 25 mM KCl, 50 µg/ml propidium iodide, 0.33% Triton X-100, 333 µg/ml RNase [Cram *et al.* (1990) Methods in Cell Biology 33:376].

- The fluorescent dye propidium iodide is used and also serves as a chromosome stabilizing agent. Swelling of the cells in the hypotonic
20 medium (which may also contain propidium iodide) may be monitored by placing a small drop of the suspension on a microscope slide and observing the cells by phase/fluorescent microscopy. The cells should exclude the propidium iodide while swelling, but some may lyse prematurely and show chromosome fluorescence. After the cells have
25 been centrifuged and resuspended in the KCl-propidium iodide buffer system, they will be lysed due to the presence of the detergent in the buffer. The chromosomes may then be dispersed and then incubated at 37°C for up to 30 minutes to permit the RNase to act. The chromosome preparation is then analyzed by flow cytometry. The propidium iodide

-144-

fluorescence can be excited at the 488 nm wavelength of an argon laser and detected through an OG 570 optical filter by a single photomultiplier tube. The single pulse may be integrated and acquired in an univariate histogram. The flow cytometer may be aligned to a CV of 2% or less

- 5 using small (1.5 μ m diameter) microspheres. The chromosome preparation is filtered through 60 μ m nylon mesh before analysis.

C. Staining of chromosomes with DNA-specific dyes

Subsequent to isolation, the chromosome preparation was stained with Hoechst 33258 at 6 μ g/ml and chromomycin A3 at 200 μ g/ml.

- 10 Fifteen minutes prior to analysis, 25 mM Na-sulphite and 10 mM Na-citrate were added to the chromosome suspension.

D. Flow sorting of chromosomes

- Chromosomes obtained from 1B3 and GHB42 cells and maintained were suspended in a polyamine-based sheath buffer (0.5 mM EGTA, 2.0
- 15 mM EDTA, 80 mM KCl, 70 mM NaCl, 15 mM Tris-HCl, pH 7.2, 0.2 mM spermine and 0.5 mM spermidine) [Sillar and Young (1981) J. Histochem. Cytochem. 29:74-78]. The chromosomes were then passed through a dual-laser cell sorter [FACStar Plus or FAXStar Vantage Becton Dickinson Immunocytometry System; other dual-laser sorters may also be
- 20 used, such as those manufactured by Coulter Electronics (Elite ESP) and Cytomation (MoFlo)] in which two lasers were set to excite the dyes separately, allowing a bivariate analysis of the chromosome by size and base-pair composition. Because of the difference between the base composition of the SATACs and the other chromosomes and the
- 25 resulting difference in interaction with the dyes, as well as size differences, the SATACs were separated from the other chromosomes.

-145-

E. Storage of the sorted artificial chromosomes

Sorted chromosomes may be pelleted by centrifugation and resuspended in a variety of buffers, and stored at 4°C. For example, the isolated artificial chromosomes may be stored in GH buffer (100 mM glycine, 1% hexylene glycol pH 8.4–8.6 adjusted with saturated Ca-hydroxide solution) [see, e.g., Hadlaczky et al. (1982) Chromosoma 86:643-659] for one day and embedded by centrifugation into agarose. The sorted chromosomes were centrifuged into an agarose bed and the plugs are stored in 500 mM EDTA at 4° C. Additional storage buffers include CMB-I/polyamine buffer (17.5 mM Tris-HCl, pH 7.4, 1.1 mM EDTA, 50 mM epsilon-amino caproic acid, 5 mM benzamide-HCl, 0.40 mM spermine, 1.0 mM spermidine, 0.25 mM EGTA, 40 mM KCl, 35 mM NaCl) and CMB-II/polyamine buffer (100 mM glycine, pH 7.5, 78 mM hexylene glycol, 0.1 mM EDTA, 50 mM epsilon-amino caproic acid, 5 mM benzamide-HCl, 0.40 mM spermine, 1.0 mM spermidine, 0.25 mM EGTA, 40 mM KCl, 35 mM NaCl).

When microinjection is the intended use, the sorted chromosomes are stored in 30% glycerol at -20° C. Sorted chromosomes may also be stored without glycerol for short periods of time (3-6 days) in storage buffers at 4°C. Exemplary buffers for microinjection include CBM-I (10 mM Tris-HCl, pH 7.5, 0.1 mM EDTA, 50 mM epsilon-amino caproic acid, 5 mM benzamide-HCl, 0.30 mM spermine, 0.75 mM spermidine), CBM-II (100 mM glycine, pH 7.5, 78 mM hexylene glycol, 0.1 mM EDTA, 50 mM epsilon-amino caproic acid, 5 mM benzamide-HCl, 0.30 mM spermine, 0.75 mM spermidine).

For long-term storage of sorted chromosomes, the above buffers are preferably supplemented with 50% glycerol and stored at -20°C.

-146-

F. Quality control**1. Analysis of the purity**

The purity of the sorted chromosomes was checked by fluorescence *in situ* hybridization (FISH) with a biotin-labeled mouse
5 satellite DNA probe [see, Hadlaczký *et al.* (1991) Proc. Natl. Acad. Sci. U.S.A. 88:8106-8110]. Purity of the isolated chromosomes was about 97–99%.

2. Characteristics of the sorted chromosomes

Pulsed field gel electrophoresis and Southern hybridization were
10 carried out to determine the size distribution of the DNA content of the sorted artificial chromosomes.

G. Functioning of the purified artificial chromosomes

To check whether their activity is preserved, the purified artificial chromosomes may be microinjected (using methods such as those
15 described in Example 13) into primary cells, somatic cells and stem cells which are then analyzed for expression of the heterologous genes carried by the artificial chromosomes, e.g., such as analysis for growth on selective medium and assays of β -galactosidase activity.

II. Sorting of mammalian artificial chromosome-containing microcells**20 A. Micronucleation**

Cells were grown to 80–90% confluency in 4 T150 flasks. Colcemid was added to a final concentration of 0.06 μ g/ml, and then incubated with the cells at 37°C for 24 hours.

B. Enucleation

25 Ten μ g/ml cytochalasin B was added and the resulting microcells were centrifuged at 15,000 rpm for 70 minutes at 28–33° C.

C. Purification of microcells by filtration

The microcells were purified using Swinnex filter units and Nucl opore filters [5 μ m and 3 μ m].

-147-

D. Staining and sorting microcells

As above, the cells were stained with Hoechst and chromomycin A3 dyes. The microcells were sorted by cell sorter to isolate the microcells that contain the mammalian artificial chromosomes.

5 E. Fusion

The microcells that contain the artificial chromosome are fused, for example, as described in Example 1.A.5., to selected primary cells, somatic cells, embryonic stem cells to generate transgenic (non-human) animals and for gene therapy purposes, and to other cells to deliver the
10 chromosomes to the cells.

EXAMPLE 11**Introduction of mammalian artificial chromosomes into insect cells**

Insect cells are useful hosts for MACs, particularly for use in the production of gene products, for a number of reasons, including:

- 15 1. A mammalian artificial chromosome provides an extra-genomic specific integration site for introduction of genes encoding proteins of interest [reduced chance of mutation in production system].
2. The large size of an artificial chromosome permits megabase size DNA integration so that genes encoding an entire pathway leading to
20 a protein or nonprotein of therapeutic value, such as an alkaloid [digitalis, morphine, taxol] can be accommodated by the artificial chromosome.
3. Amplification of genes encoding useful proteins can be accomplished in the artificial mammalian chromosome to obtain higher protein yields in insect cells.
- 25 4. Insect cells support required post-translational modifications (glycosylation, phosphorylation) essential for protein biological function.
5. Insect cells do not support mammalian viruses — eliminates cross-contamination of product with human infectious agents.

-148-

6. The ability to introduce chromosomes circumvents traditional recombinant baculovirus systems for production of nutritional, industrial or medicinal proteins in insect cell systems.
7. The low temperature optimum for insect cell growth (28° C) permits reduced energy cost of production.
8. Serum free growth medium for insect cells will result in lower production costs.
9. Artificial chromosome-containing cells can be stored indefinitely at low temperature.
10. Insect larvae will serve as biological factories for the production of nutritional, medicinal or industrial proteins by microinjection of fertilized insect eggs.
 - A. **Demonstration that insect cells recognize mammalian promoters**

Gene constructs containing a mammalian promoter, such as the CMV promoter, linked to a detectable marker gene [*Renilla* luciferase gene (see, e.g., U.S. Patent No. 5,292,658 for a description of DNA encoding the *Renilla* luciferase, and plasmid pTZrLuc-1, which can provide the starting material for construction of such vectors, see also SEQ ID No. 10] and also including the simian virus 40 (SV40) promoter operably linked to the β -galactosidase gene were introduced into the cells of two species *Trichoplusia ni* [cabbage looper] and *Bombyx mori* [silkworm].

After transferring the constructs into the insect cell lines either by electroporation or by microinjection, expression of the marker genes was detected in luciferase assays (see e.g., Example 12.C.3) and in β -galactosidase assays (such as lacZ staining assays) after a 24-h incubation. In each case a positive result was obtained in the samples containing the genes which was absent in samples in which the genes were omitted. In addition, a *B. mori* β -actin promoter-*Renilla* luciferase

-149-

gene fusion was introduced into the *T. ni* and *B. mori* cells which yielded light emission after transfection. Thus, certain mammalian promoters function to direct expression of these marker genes in insect cells.

Therefore, MACs are candidates for expression of heterologous genes in
5 insect cells.

B. Construction of vectors for use in insect cells and fusion with mammalian cells

1. Transform LMTK⁻ cells with expression vector with:
 - a. *B. mori* β -actin promoter— Hyg^r selectable marker
10 gene for insect cells, and
 - b. SV40 or CMV promoters controlling a puromycin^r selectable marker gene for mammalian cells.
2. Detect expression of the mammalian promoter in LMTK cells (puromycin^r LMTK cells)
- 15 3. Use puromycin^r cells in fusion experiments with *Bombyx* and *Trichoplusia* cells, select Hyg^r cells.

C. Insertion of the MACs into insect cells

These experiments are designed to detect expression of a detectable marker gene [such as the β -galactosidase gene expressed
20 under the control of a mammalian promoter, such as pSV40] located on a MAC that has been introduced into an insect cell. Data indicate that β -gal was expressed.

Insect cells are fused with mammalian cells containing mammalian artificial chromosomes, e.g., the minichromosome [EC3/7C5] or the mini
25 and the megachromosome [such as GHB42, which is a cell line recloned from G3D5] or a cell line that carries only the megachromosome [such as H1D3 or a reclone therefrom]. Fusion is carried out as follows:

-150-

1. mammalian + insect cells (50/50%) in log phase growth are mixed;
2. calcium/PEG cell fusion: (10 min — 0.5 h);
3. heterokaryons (+ 72 h) are selected.

5 The following selection conditions to select for insect cells that contain a MAC can be used: [+ = positive selection; - = negative selection]:

1. growth at 28° C (+ insect cells, - mammalian cells);
2. Grace's insect cell medium [SIGMA] (- mammalian cells);
- 10 3. no exogenous CO₂ (- mammalian cells); and/or
4. antibiotic selection (Hyg or G418) (+ transformed insect cells).

Immediately following the fusion protocol, many heterokaryons [fusion events] are observed between the mammalian and each species of insect cells [up to 90% heterokaryons]. After growth [2 + weeks] on
15 insect medium containing G418 and/or hygromycin at selection levels used for selection of transformed mammalian cells, individual colonies are detected growing on the fusion plates. By virtue of selection for the antibiotic resistance conferred by the MAC and selection for insect cells, these colonies should contain MACs.

20 The *B. mori* β -actin gene promoter has been shown to direct expression of the β -galactosidase gene in *B. mori* cells and mammalian cells (e.g., EC3/7C5 cells). The *B. mori* β -actin gene promoter is, thus, particularly useful for inclusion in MACs generated in mammalian cells that will subsequently be transferred into insect cells because the
25 presence of any marker gene linked to the promoter can be determined in the mammalian and resulting insect cell lines.

-151-

EXAMPLE 12**Preparation of chromosome fragmentation vectors and other vectors for targeted integration of DNA into MACs**

- Fragmentation of the megachromosome should ultimately result in smaller stable chromosomes that contain about 15 Mb to 50 Mb that will be easily manipulated for use as vectors. Vectors to effect such fragmentation should also aid in determination and identification of the elements required for preparation of an in vitro-produced artificial chromosome.
- Reduction in the size of the megachromosome can be achieved in a number of different ways including: stress treatment, such as by starvation, or cold or heat treatment; treatment with agents that destabilize the genome or nick DNA, such as BrdU, coumarin, EMS and others; treatment with ionizing radiation [see, e.g., Brown (1992) Curr. Opin. Genes Dev. 2:479-486]; and telomere-directed in vivo chromosome fragmentation [see, e.g., Farr et al. (1995) EMBO J. 14:5444-5454].
- A. Preparation of vectors for fragmentation of the artificial chromosome and also for targeted integration of select d gene products**
- 1. Construction of pTEMPUD**
- Plasmid pTEMPUD [see Figure 5] is a mouse homologous recombination "killer" vector for in vivo chromosome fragmentation, and also for inducing large-scale amplification via site-specific integration. With reference to Figure 5, the ~3,625-bp Sall-PstI fragment was derived from the pBabe-puro retroviral vector [see, Morgenstern et al. (1990) Nucleic Acids Res. 18:3587-3596]. This fragment contains DNA encoding ampicillin resistance, the pUC origin of replication, and the puromycin N-acetyl transferase gene under control of the SV40 early promoter. The URA3 gene portion comes from the pYAC5 cloning vector [SIGMA]. URA3 was cut out of pYAC5 with Sall-XhoI digestion, cloned

-152-

into pNEB193 [New England Biolabs], which was then cut with EcoRI-Sall and ligated to the Sall site of pBabepuro to produce pPU.

A 1293-bp fragment [see SEQ ID No. 1] encoding the mouse major satellite, was isolated as an EcoRI fragment from a DNA library produced
5 from mouse LMTK fibroblast cells and inserted into the EcoRI site of pPU to produce pMPU.

The TK promoter-driven diphtheria toxin gene [DT-A] was derived from pMC1DT-A [see, Maxwell et al. (1986) Cancer Res. 46:4660-4666] by BglII-XhoI digestion and cloned into the pMC1neo poly A expression
10 vector [STRATAGENE, La Jolla, CA] by replacing the neomycin-resistance gene coding sequence. The TK promoter, DT-A gene and poly A sequence were removed from this vector, cohesive ends were filled with Klenow and the resulting fragment blunt end-ligated and ligated into the SnaBI [TACGTA] of pMPU to produce pMPUD.

15 The Hutel 2.5-kb fragment [see SEQ ID No.3] was inserted at the PstI site [see the 6100 PstI - 3625 PstI fragment on pTEMPUD] of pMPUD to produce pTEMPUD. This fragment includes a human telomere. It includes a unique BglII site [see nucleotides 1042-1047 of SEQ ID No.3], which will be used as a site for introduction of a synthetic
20 telomere that includes multiple repeats [80] of TTAGGG with BamHI and BglII ends for insertion into the BglII site which will then remain unique, since the BamHI overhang is compatible with the BglII site. Ligation of a BamHI fragment to a BglII destroys the BglII site, so that only a single BglII site will remain. Selection for the unique BglII site insures that the
25 synthetic telomere will be inserted in the correct orientation. The unique BglII site is the site at which the vector is linearized.

-153-

To generate a synthetic telomere made up of multiple repeats of the sequence TTAGGG, attempts were made to clone or amplify ligation products of 30-mer oligonucleotides containing repeats of the sequence. Two 30-mer oligonucleotides, one containing four repeats of TTAGGG
5 bounded on each end of the complete run of repeats by half of a repeat and the other containing five repeats of the complement AATCCC, were annealed. The resulting double-stranded molecule with 3-bp protruding ends, each representing half of a repeat, was expected to ligate with itself to yield concatamers of $n \times 30$ bp. However, this approach was
10 unsuccessful, likely due to formation of quadruplex DNA from the G-rich strand. Similar difficulty has been encountered in attempts to generate long repeats of the pentameric human satellite II and III units. Thus, it appears that, in general, any oligomer sequence containing periodically spaced consecutive series of guanine nucleotides is likely to form
15 undesired quadruplex formation that hinders construction of long double-stranded DNAs containing the sequence.

Therefore, in another attempt to construct a synthetic telomere for insertion into the BglIII site of pTEMPUD, the starting material was based on the complementary C-rich repeat sequence (i.e., AATCCC) which
20 would not be susceptible to quadruplex structure formation. Two plasmids, designated pTEL280110 and pTel280111, were constructed as follows to serve as the starting materials.

First, a long oligonucleotide containing 9 repeats of the sequence AATCCC (i.e., the complement of telomere sequence TTAGGG) in
25 reverse order bounded on each end of the complete run of repeats by half of a repeat (therefore, in essence, containing 10 repeats), and recognition sites for PstI and PacI restriction enzymes was synthesized using standard methods. The oligonucleotide sequence is as follows:
5'-AAACTGCAGGTTAATTAACCCTAACCCTAACCCTAACCCTAACCCTAAC

-154-

CCTAACCCTAACCCTAACCCTAACCCGGGAT-3' (SEQ ID NO. 29)

A partially complementary short oligonucleotide of sequence

3'-TTGGGCCCTAGGCTTAAGG-5' (SEQ ID NO. 30)

was also synthesized. The oligonucleotides were gel-purified, annealed,
5 repaired with Klenow polymerase and digested with EcoRI and PstI. The
resulting EcoRI/PstI fragment was ligated with EcoRI/PstI-digested
pUC19. The resulting plasmid was used to transform E. coli DH5 α
competent cells and plasmid DNA (pTel102) from one of the
transformants surviving selection on LB/ampicillin was digested with
10 PacI, rendered blunt-ended by Klenow and dNTPs and digested with
HindIII. The resulting 2.7-kb fragment was gel-purified.

Simultaneously, the same plasmid was amplified by the
polymerase chain reaction using extended and more distal 26-mer M13
sequencing primers. The amplification product was digested with SmaI
15 and HindIII, the double-stranded 84-bp fragment containing the 60-bp
telomeric repeat (plus 24 bp of linker sequence) was isolated on a 6%
native polyacrylamide gel, and ligated with the double-digested pTel102
to yield a 120-bp telomeric sequence. This plasmid was used to
transform DH5 α cells. Plasmid DNA from two of the resulting
20 recombinants that survived selection on ampicillin (100 μ g/ml) was
sequenced on an ABI DNA sequencer using the dye-termination method.
One of the plasmids, designated pTel29, contained a sequence of 20
repeats of the sequence TTAGGG (i.e., 19 successive repeats of
TTAGGG bounded on each end of the complete run of repeats with half
25 of a repeat). The other plasmid, designated pTel28, had undergone a
deletion of 2 bp (TA) at the junction where the two sequences, each
containing, in essence, 10 repeats of the TTAGGG sequence, that had
been ligated to yield the plasmid. This resulted in a GGGTGGG motif at
the junction in pTel28. This mutation provides a useful tag in telomere-

-155-

directed chromosome fragmentation experiments. Therefore, the pTel29 insert was amplified by PCR using pUC/M13 sequencing primers based on sequence somewhat longer and farther from the polylinker than usual as follows:

5 5'-GCCAGGGTTTTCCCAGTCACGACGT-3' (SEQ ID NO. 31)

or in some experiments

 5'-GCTGCAAGGCGATTAAGTTGGGTAAC-3' (SEQ ID NO. 32)

as the m13 forward primer, and

 5'-TATGTTGTGTGGAATTGTGAGCGGAT-3' (SEQ ID NO. 33)

10 as the m13 reverse primer.

The amplification product was digested with SmaI and HindIII. The resulting 144-bp fragment was gel-purified on a 6% native polyacrylamide gel and ligated with pTel28 that had been digested with PacI, blunt-ended with Klenow and dNTP and then digested with HindIII

15 to remove linker. The ligation yielded a plasmid designated pTel2801 containing a telomeric sequence of 40 repeats of the sequence TTAGGG in which one of the repeats (i.e., the 30th repeat) lacked two nucleotides (TA), due to the deletion that had occurred in pTel28, to yield a repeat as follows: TGGG.

20 In the next extension step, pTel2801 was digested with SmaI and HindIII and the 264-bp insert fragment was gel-purified and ligated with pTel2801 which had been digested with PacI, blunt-ended and digested with HindIII. The resulting plasmid was transformed into DH5 α cells and plasmid DNA from 12 of the resulting transformants that survived

25 selection on ampicillin was examined by restriction enzyme analysis for the presence of a 0.5-kb EcoRI/PstI insert fragment. Eleven of the recombinants contained the expected 0.5-kb insert. The inserts of two of the recombinants were sequenced and found to be as expected. These plasmids were designated pTel280110 and pTel280111. These

-156-

plasmids, which are identical, both contain 80 repeats of the sequence TTAGGG, in which two of the repeats (i.e., the 30th and 70th repeats) lacked two nucleotides (TA), due to the deletion that had occurred in pTel28, to yield a repeat as follows: TGGG. Thus, in each of the cloning
5 steps (except the first), the length of the synthetic telomere doubled; that is, it was increasing in size exponentially. Its length was 60×2^n bp, wherein n is the number of extension cloning steps undertaken. Therefore, in principle (assuming *E. coli*, or any other microbial host, e.g., yeast, tolerates long tandem repetitive DNA), it is possible to assemble
10 any desirable size of safe telomeric repeats.

In a further extension step, pTel280110 was digested with PacI, blunt-ended with Klenow polymerase in the presence of dNTP, then digested with HindIII. The resulting 0.5-kb fragment was gel purified. Plasmid pTel280111 was cleaved with SmaI and HindIII and the 3.2-kb
15 fragment was gel-purified and ligated to the 0.5-kb fragment from pTel280110. The resulting plasmid was used to transform DH5 α cells. Plasmid DNA was purified from transformants surviving ampicillin selection. Nine of the selected recombinants were examined by restriction enzyme analysis for the presence of a 1.0-kb EcoRI/PstI
20 fragment. Four of the recombinants (designated pTlk2, pTlk6, pTlk7 and pTlk8) were thus found to contain the desired 960 bp telomere DNA insert sequence that included 160 repeats of the sequence TTAGGG in which four of the repeats lacked two nucleotides (TA), due to the deletion that had occurred in pTel28, to yield a repeat as follows: TGGG.
25 Partial DNA sequence analysis of the EcoRI/PstI fragment of two of these plasmids (i.e., pTlk2 and pTlk6), in which approximately 300 bp from both ends of the fragment were elucidated, confirmed that the sequence was composed of successive repeats of the TTAGGG sequence.

-157-

In order to add PmeI and BglII sites to the synthetic telomere sequence, pTlk2 was digested with PacI and PstI and the 3.7-kb fragment (i.e., 2.7-kb pUC19 and 1.0-kb repeat sequence) was gel-purified and ligated at the PstI cohesive end with the following

5 oligonucleotide 5'-GGGTTTAAACAGATCTCTGCA-3' (SEQ ID NO. 34). The ligation product was subsequently repaired with Klenow polymerase and dNTP, ligated to itself and transformed into E. coli strain DH5 α . A total of 14 recombinants surviving selection on ampicillin were obtained. Plasmid DNA from each recombinant was able to be cleaved with BglII

10 indicating that this added unique restriction site had been retained by each recombinant. Four of the 14 recombinants contained the complete 1-kb synthetic telomere insert, whereas the insert of the remaining 10 recombinants had undergone deletions of various lengths. The four plasmids in which the 1-kb synthetic telomere sequence remained intact

15 were designated pTlkV2, pTlkV5, pTlkV8 and pTlkV12. Each of these plasmids could also be digested with PmeI; in addition the presence of both the BglII and PmeI sites was verified by sequence analysis. Any of these four plasmids can be digested with BamHI and BglII to release a fragment containing the 1-kb synthetic telomere sequence which is then

20 ligated with BglII-digested pTEMPUD.

2. Use of pTEMPUD for in vivo chromosome fragmentation

Linearization of pTEMPUD by BglII results in a linear molecule with a human telomere at one end. Integration of this linear fragment into the chromosome, such as the megachromosome in hybrid cells or any mouse

25 chromosome which contains repeats of the mouse major satellite sequence results in integration of the selectable marker puromycin-resistance gene and cleavage of the plasmid by virtue of the telomeric end. The DT gene prevents that entire linear fragment from integrating by random events, since upon integration and expression it is toxic.

-158-

Thus random integration will be toxic, so site-directed integration into the targeted DNA will be selected. Such integration will produce fragmented chromosomes.

The fragmented truncated chromosome with the new telomere will
5 survive, and the other fragment without the centromere will be lost.
Repeated in vivo fragmentations will ultimately result in selection of the smallest functioning artificial chromosome possible. Thus, this vector can be used to produce minichromosomes from mouse chromosomes, or to fragment the megachromosome. In principle, this vector can be used
10 to target any selected DNA sequence in any chromosome to achieve fragmentation.

3. Construction of pTERPUD

A fragmentation/targeting vector analogous to pTEMPUD for in vivo chromosome fragmentation, and also for inducing large-scale
15 amplification via site-specific integration but which is based on mouse rDNA sequence instead of mouse major satellite DNA has been designated pTERPUD. In this vector, the mouse major satellite DNA sequence of pTEMPUD has been replaced with a 4770-bp BamHI fragment of megachromosome clone 161 which contains sequence
20 corresponding to nucleotides 10,232-15,000 in SEQ ID NO. 16.

4. pHASPUD and pTEMPHu3

Vectors that specifically target human chromosomes can be constructed from pTEMPUD. These vectors can be used to fragment specific human chromosomes, depending upon the selected satellite
25 sequence, to produce human minichromosomes, and also to isolate human centromeres.

-159-

a. pHASPUD

To render pTEMPUD suitable for fragmenting human chromosomes, the mouse major satellite sequence is replaced with human satellite sequences. Unlike mouse chromosomes, each human chromosome has a unique satellite sequence. For example, the mouse major satellite has been replaced with a human hexameric α -satellite [or alphoid satellite] DNA sequence. This sequence is an 813-bp fragment [nucleotide 232-1044 of SEQ ID No. 2] from clone pS12, deposited in the EMBL database under Accession number X60716, isolated from a human colon carcinoma cell line Colo320 [deposited under Accession No. ATCC CCL 220.1]. The 813-bp alphoid fragment can be obtained from the pS12 clone by nucleic acid amplification using synthetic primers, each of which contains an EcoRI site, as follows:

GGGGAATTCAT TGGGATGTTT CAGTTGA forward primer [SEQ ID No. 4]
15 CGAAAGTCCCC CTTAGGAGAT CTTAAGGA reverse primer [SEQ ID No. 5].

Digestion of the amplified product with EcoRI results in a fragment with EcoRI ends that includes the human α -satellite sequence. This sequence is inserted into pTEMPUD in place of the EcoRI fragment that contains the mouse major satellite to yield pHASPUD.

20 Vector pHASPUD was linearized with BglII and used to transform EJ30 (human fibroblast) cells by scrape loading. Twenty-seven puromycin-resistant transformant strains were obtained.

b. pTEMPHu3

In pTEMPHu3, the mouse major satellite sequence is replaced by the 3kb human chromosome 3-specific α -satellite from D3Z1 [deposited under ATCC Accession No. 85434; see, also Yrokov (1989) Cytogenet. Cell Genet. 51:1114].

-160-

5. Use of the pTEMPHU3 to induce amplification on human chromosome #3

Each human chromosome contains unique chromosome-specific alphoid sequence. Thus, pTEMPHU3, which is targeted to the
5 chromosome 3-specific α -satellite, can be introduced into human cells under selective conditions, whereby large-scale amplification of the chromosome 3 centromeric region and production of a *de novo* chromosome ensues. Such induced large-scale amplification provides a means for inducing *de novo* chromosome formation and also for in vivo
10 cloning of defined human chromosome fragments up to megabase size.

For example, the break-point in human chromosome 3 is on the short arm near the centromere. This region is involved in renal cell carcinoma formation. By targeting pTEMPHu3 to this region, the induced large-scale amplification may contain this region, which can then be
15 cloned using the bacterial and yeast markers in the pTEMPHu3 vector.

The pTEMPHu3 cloning vector allows not only selection for homologous recombinants, but also direct cloning of the integration site in YACS. This vector can also be used to target human chromosome 3, preferably with a deleted short arm, in a mouse-human mono-
20 chromosomal microcell hybrid line. Homologous recombinants can be screened by nucleic acid amplification (PCR), and amplification can be screened by DNA hybridization, Southern hybridization, and *in situ* hybridization. The amplified region can be cloned into a YAC. This vector and these methods also permit a functional analysis of cloned
25 chromosome regions by reintroducing the cloned amplified region into mammalian cells.

-161-

B. Preparation of libraries in YAC vectors for cloning of centromeres and identification of functional chromosomal units

Another method that may be used to obtain smaller-sized functional mammalian artificial chromosome units and to clone centromeric DNA involves screening of mammalian DNA YAC vector-based libraries and functional analysis of potential positive clones in a transgenic mouse model system. A mammalian DNA library is prepared in a YAC vector, such as YRT2 [see Schedl *et al.* (1993) Nuc. Acids Res. 21:4783-4787], which contains the murine tyrosinase gene. The library is screened for hybridization to mammalian telomere and centromere sequence probes. Positive clones are isolated and microinjected into pronuclei of fertilized oocytes of NMRI/Han mice following standard techniques. The embryos are then transferred into NMRI/Han foster mothers. Expression of the tyrosinase gene in transgenic offspring confers an identifiable phenotype (pigmentation). The clones that give rise to tyrosinase-expressing transgenic mice are thus confirmed as containing functional mammalian artificial chromosome units.

Alternatively, fragments of SATACs may be introduced into the YAC vectors and then introduced into pronuclei of fertilized oocytes of NMRI/Han mice following standard techniques as above. The clones that give rise to tyrosinase-expressing transgenic mice are thus confirmed as containing functional mammalian artificial chromosome units, particularly centromeres.

C. Incorporation of Heterologous Genes into Mammalian Artificial Chromosomes through The Use of Homology Targeting Vectors

As described above, the use of mammalian artificial chromosomes for expression of heterologous genes obviates certain negative effects that may result from random integration of heterologous plasmid DNA into the recipient cell genome. An essential feature of the mammalian artificial chromosome that makes it a useful tool in avoiding the negative

-162-

effects of random integration is its presence as an extra-genomic gene source in recipient cells. Accordingly, methods of specific, targeted incorporation of heterologous genes exclusively into the mammalian artificial chromosome, without extraneous random integration into the
5 genome of recipient cells, are desired for heterologous gene expression from a mammalian artificial chromosome.

One means of achieving site-specific integration of heterologous genes into artificial chromosomes is through the use of homology targeting vectors. The heterologous gene of interest is subcloned into a
10 targeting vector which contains nucleic acid sequences that are homologous to nucleotides present in the artificial chromosome. The vector is then introduced into cells containing the artificial chromosome for specific site-directed integration into the artificial chromosome through a recombination event at sites of homology between the vector
15 and the chromosome. The homology targeting vectors may also contain selectable markers for ease of identifying cells that have incorporated the vector into the artificial chromosome as well as lethal selection genes that are expressed only upon extraneous integration of the vector into the recipient cell genome. Two exemplary homology targeting vectors,
20 λ CF-7 and p λ CF-7-DTA, are described below.

1. Construction of Vector λ CF-7

Vector λ CF-7 contains the cystic fibrosis transmembrane conductance regulator [CFTR] gene as an exemplary therapeutic molecule-encoding nucleic acid that may be incorporated into mammalian
25 artificial chromosomes for use in gene therapy applications. This vector, which also contains the puromycin-resistance gene as a selectable marker, as well as the Saccharomyces cerevisiae ura3 gene [orotidine-5-phosphate decarboxylase], was constructed in a series of steps as follows.

-163-

a. Construction of pURA

Plasmid pURA was prepared by ligating a 2.6-kb Sall/XhoI fragment from the yeast artificial chromosome vector pYAC5 [Sigma; see also Burke et al. (1987) Science 236:806-812 for a description of YAC
5 vectors as well as GenBank Accession no. U01086 for the complete sequence of pYAC5] containing the S. cerevisiae ura3 gene with a 3.3-kb Sall/SmaI fragment of pHyg [see, e.g., U.S. Patent Nos. 4,997,764, 4,686,186 and 5,162,215, and the description above]. Prior to ligation
10 the XhoI end was treated with Klenow polymerase for blunt end ligation to the SmaI end of the 3.3 kb fragment of pHyg. Thus, pURA contains the S. cerevisiae ura3 gene, and the E. coli ColE1 origin of replication and the ampicillin-resistance gene. The uraE gene is included to provide a means to recover the integrated construct from a mammalian cell as a YAC clone.

15 b. Construction of pUP2

Plasmid pURA was digested with Sall and ligated to a 1.5-kb Sall fragment of pCEPUR. Plasmid pCEPUR is produced by ligating the 1.1 kb SnaBI-NheI fragment of pBabe-puro [Morgenstern et al. (1990) Nucl. Acids Res. 18:3587-3596; provided by Dr. L. Székely
20 (Microbiology and Tumorbiology Center, Karolinska Institutet, Stockholm); see, also Tonghua et al. (1995) Chin. Med. J. (Beijing, Engl. Ed.) 108:653-659; Couto et al. (1994) Infect. Immun. 62:2375-2378; Dunckley et al. (1992) FEBS Lett. 296:128-34; French et al. (1995) Anal. Biochem. 228:354-355; Liu et al. (1995) Blood 85:1095-1103;
25 International PCT application Nos. WO 9520044; WO 9500178, and WO 9419456] to the NheI-NruI fragment of pCEP4 [Invitrogen].

-164-

The resulting plasmid, pUP2, contains all the elements of pURA plus the puromycin-resistance gene linked to the SV40 promoter and polyadenylation signal from pCEPUR.

c. Construction of pUP-CFTR

5 The intermediate plasmid pUP-CFTR was generated in order to combine the elements of pUP2 into a plasmid along with the CFTR gene. First, a 4.5-kb Sall fragment of pCMV-CFTR that contains the CFTR-encoding DNA [see, also, Riordan *et al.* (1989) Science 245:1066-1073, U.S. Patent No. 5,240,846, and Genbank Accession no. M28668
10 for the sequence of the CFTR gene] containing the CFTR gene only was ligated to XhoI-digested pCEP4 [Invitrogen and also described herein] in order to insert the CFTR gene in the multiple cloning site of the Epstein Barr virus-based (EBV) vector pCEP4 [Invitrogen, San Diego, CA; see also Yates *et al.* (1985) Nature 313:812-815; see, also U.S. Patent No.
15 5,468,615] between the CMV promoter and SV40 polyadenylation signal. The resulting plasmid was designated pCEP-CFTR. Plasmid pCEP-CFTR was then digested with Sall and the 5.8-kb fragment containing the CFTR gene flanked by the CMV promoter and SV40 polyadenylation signal was ligated to Sall-digested pUP2 to generate
20 pUP-CFTR. Thus, pUP-CFTR contains all elements of pUP2 plus the CFTR gene linked to the CMV promoter and SV40 polyadenylation signal.

d. Construction of λ CF-7

Plasmid pUP-CFTR was then linearized by partial digestion with EcoRI and the 13 kb fragment containing the CFTR gene was ligated
25 with EcoRI-digested Charon 4A λ [see Blattner *et al.* (1977) Science 196:161; Williams and Blattner (1979) J. Virol. 29:555 and Sambrook *et al.* (1989) Molecular Cloning, A Laboratory Manual, Second Ed., Cold Spring Harbor Laboratory Press, Volume 1, Section 2.18, for descriptions of Charon 4A λ]. The resulting vector, λ CF8, contains the Charon 4A λ

-165-

bacteriophage left arm, the CFTR gene linked to the CMV promoter and SV40 polyadenylation signal, the ura3 gene, the puromycin-resistance gene linked to the SV40 promoter and polyadenylation signal, the thymidine kinase promoter [TK], the ColE1 origin of replicaton, the
5 amplicillin resistance gene and the Charon 4A λ bacteriophage right arm. The λ CF8 construct was then digested with XhoI and the resulting 27.1 kb was ligated to the 0.4kb XhoI/EcoRI fragment of pJBP86 [described below], containing the SV40 polyA signal and the EcoRI-digested Charon 4A λ right arm. The resulting vector λ CF-7 contains the Charon 4A λ left
10 arm, the CFTR encoding DNA linked to the CMV promoter and SV40 polyA signal, the ura3 gene, the puromycin resistance gene linked to the SV40 promoter and polyA signal and the Charon 4A λ right arm. The λ DNA fragments provide encode sequences homologous to nucleotides present in the exemplary artificial chromosomes.

15 The vector is then introduced into cells containing the artificial chromosomes exemplified herein. Accordingly, when the linear λ CF-7 vector is introduced into megachromosome-carrying fusion cell lines, such as described herein, it will be specifically integrated into the megachromosome through recombination between the homologous
20 bacteriophage λ sequences of the vector and the artificial chromosome.

2. Construction of Vector λ CF-7-DTA

Vector λ CF-7-DTA also contains all the elements contained in λ CF-7, but additionally contains a lethal selection marker, the diphtheria toxin-A (DT-A) gene as well as the ampicillin-resistance gene and an origin of
25 replication. This vector was constructed in a series of steps as follows.

-166-

a. Construction of pJBP86

Plasmid pJBP86 was used in the construction of λ CF-7, above. A 1.5-kb Sall fragment of pCEPUR containing the puromycin-resistance gene linked to the SV40 promoter and polyadenylation signal was ligated to HindIII-digested pJB8 [see, e.g., Ish-Horowitz *et al.* (1981) Nucleic Acids Res. 9:2989-2998; available from ATCC as Accession No. 37074; commercially available from Amersham, Arlington Heights, IL]. Prior to ligation the Sall ends of the 1.5 kb fragment of pCEPUR and th4 HindIII linearized pJB8 ends were treated with Klenow polymerase. The resulting vector pJBP86 contains the puromycin resistance gene linked to the SV40 promoter and polyA signal, the 1.8 kb COS region of Charon 4A λ , the ColE1 origin of replication and the ampicillin resistance gene.

b. Construction of pMEP-DTA

A 1.1-kb XhoI/Sall fragment of pMC1-DT-A [see, e.g., Maxwell *et al.* (1986) Cancer Res. 46:4660-4666] containing the diphtheria toxin-A gene was ligated to XhoI-digested pMEP4 [Invitrogen, San Diego, CA] to generate pMEP-DTA. To produce pMC1-DT-A, the coding region of the DTA gene was isolated as a 800 bp PstI/HindIII fragment from p2249-1 and inserted into pMC1neopolyA [pMC1 available from Stratagene] in place of the neo gene and under the control of the TK promoter. The resulting construct pMC1DT-A was digested with HindIII, the ends filled by Klenow and Sall linkers were ligated to produce a 1061 bp TK-DTA gene cassette with an XhoI end [5'] and a Sall end containing the 270 bp TK promoter and the ~790 bp DT-A fragment. This fragment was ligated into XhoI-digested pMEP4.

Plasmid pMEP-DTA thus contains the DT-A gene linked to the TK promoter and SV40, ColE1 origin of replication and the ampicillin-resistance gene.

-167-

c. Construction of pJB83-DTA9

Plasmid pJB8 was digested with HindIII and Clal and ligated with an oligonucleotide [see SEQ ID NOs. 7 and 8 for the sense and antisense strands of the oligonucleotide, respectively] to generate pJB83.

- 5 The oligonucleotide that was ligated to Clal/HindIII-digested pJB8 contained the recognition sites of SwaI, PacI and SrfI restriction endonucleases. These sites will permit ready linearization of the p λ CF-7-DTA construct.

- Next, a 1.4-kb XhoI/Sall fragment of pMEP-DTA, containing the DT-A gene was ligated to Sall-digested pJB83 to generate pJB83-DTA9.

d. Construction of λ CF-7-DTA

- The 12-bp overhangs of λ CF-7 were removed by Mung bean nuclease and subsequent T4 polymerase treatments. The resulting 41.1-kb linear λ CF-7 vector was then ligated to pJB83-DTA9 which had been digested with Clal and treated with T4 polymerase. The resulting vector, λ CF-7-DTA, contains all the elements of λ CF-7 as well as the DT-A gene linked to the TK promoter and the SV40 polyadenylation signal, the 1.8 kB Charon 4A λ COS region, the ampicillin-resistance gene [from pJB83-DTA9] and the Col E1 origin of replication [from pJB83-DT9A].

20 D. Targeting vectors using luciferase markers: Plasmid pMCT-RUC

- Plasmid pMCT-RUC [14kbp] was constructed for site-specific targeting of the Renilla luciferase [see, e.g., U.S. Patent Nos. 5,292,658 and 5,418,155 for a description of DNA encoding *Renilla* luciferase, and plasmid pTZrLuc-1, which can provide the starting material for construction of such vectors] gene to a mammalian artificial chromosome. The relevant features of this plasmid are the *Renilla* luciferase gene under transcriptional control of the human cytomegalovirus immediate-early gene enhancer/promoter; the hygromycin-resistance gene a, positive selectable marker, under the

-168-

- transcriptional control of the thymidine kinase promoter. In particular, this plasmid contains plasmid pAG60 [see, e.g., U.S. Patent Nos. 5,118,620, 5,021,344, 5,063,162 and 4,946,952; see, also Colbert-Garapin et al. (1981) J. Mol. Biol. **150**:1-14], which includes DNA (i.e.,
- 5 the neomycin-resistance gene) homologous to the minichromosome, as well as the *Renilla* and hygromycin-resistance genes, the HSV-tk gene under control of the tk promoter as a negative selectable marker for homologous recombination, and a unique HpaI site for linearizing the plasmid.
- 10 This construct was introduced, via calcium phosphate transfection, into EC3/7C5 cells [see, Lorenz et al. (1996) J. Biolum. Chemilum. **11**:31-37]. The EC3/7C5 cells were maintained as a monolayer [see, Gluzman (1981) Cell **23**:175-183]. Cells at 50% confluency in 100 mm Petri dishes were used for calcium phosphate transfection [see, Harper et
- 15 al. (1981) Chromosoma **83**:431-439] using 10 μ g of linearized pMCT-RUC per plate. Colonies originating from single transfected cells were isolated and maintained in F-12 medium containing hygromycin (300 μ g/mL) and 10% fetal bovine serum. Cells were grown in 100 mm Petri dishes prior to the *Renilla* luciferase assay.
- 20 The *Renilla* luciferase assay was performed [see, e.g., Matthys et al. (1977) Biochemistry **16**:85-91]. Hygromycin-resistant cell lines obtained after transfection of EC3/7C5 cells with linearized plasmid pMCT-RUC ["B" cell lines] were grown to 100% confluency for measurements of light emission in vivo and in vitro. Light emission was
- 25 measured in vivo after about 30 generations as follows: growth medium was removed and replaced by 1 mL RPMI 1640 containing coelenterazine [1 mmol/L final concentration]. Light emission from cells was then visualized by placing the Petri dishes in a low light video image analyzer [Hamamatsu Argus-100]. An image was formed after 5 min. of photon

-169-

accumulation using 100% sensitivity of the photon counting tube. For measuring light emission in vitro, cells were trypsinized and harvested from one Petri dish, pelleted, resuspended in 1mL assay buffer [0.5 mol/L NaCl, 1 mmol/L EDTA, 0.1 mol/L potassium phosphate, pH 7.4] and
5 sonicated on ice for 10 s. Lysates were then assayed in a Turner TD-20e luminometer for 10 s after rapid injection of 0.5 mL of 1 mmol/L coelenterazine, and the average value of light emission was recorded as LU [1 LU = 1.6×10^6 hu/s for this instrument].

Independent cell lines of EC3/7C5 cells transfected with linearized
10 plasmid pMCT-RUC showed different levels of *Renilla* luciferase activity. Similar differences in light emission were observed when measurements were performed on lysates of the same cell lines. This variation in light emission was probably due to a position effect resulting from the random integration of plasmid pMCT-RUC into the mouse genome, since
15 enrichment for site targeting of the luciferase gene was not performed in this experiment.

To obtain transfectant populations enriched in cells in which the luciferase gene had integrated into the minichromosome, transfected cells were grown in the presence of ganciclovir. This negative selection
20 medium selects against cells in which the added pMCT-RUC plasmid integrated into the host EC3/7C5 genome. This selection thereby enriches the surviving transfectant population with cells containing pMCT-RUC in the minichromosome. The cells surviving this selection were evaluated in luciferase assays which revealed a more uniform level
25 of luciferase expression. Additionally, the results of in situ hybridization assays indicated that the *Renilla* luciferase gene was contained in the minichromosome in these cells, which further indicates successful targeting of pMCT-RUC into the minichromosome.

-170-

Plasmid pNEM-1, a variant of pMCT-RUC which also contains λ DNA to provide an extended region of homology to the minichromosome [see, other targeting vectors, below], was also used to transfect EC3/7C5 cells. Site-directed targeting of the *Renilla* luciferase gene and the hygromycin-resistance gene in pNEM-1 to the minichromosome in the recipient EC3/7C5 cells was achieved. This was verified by DNA amplification analysis and by *in situ* hybridization. Additionally, luciferase gene expression was confirmed in luciferase assays of the transfectants.

E. Protein secretion targeting vectors

Isolation of heterologous proteins produced intracellularly in mammalian cell expression systems requires cell disruption under potentially harsh conditions and purification of the recombinant protein from cellular contaminants. The process of protein isolation may be greatly facilitated by secretion of the recombinantly produced protein into the extracellular medium where there are fewer contaminants to remove during purification. Therefore, secretion targeting vectors have been constructed for use with the mammalian artificial chromosome system.

A useful model vector for demonstrating production and secretion of heterologous protein in mammalian cells contains DNA encoding a readily detectable reporter protein fused to an efficient secretion signal that directs transport of the protein to the cell membrane and secretion of the protein from the cell. Vectors pLNCX-ILRUC and pLNCX-ILRUC λ , described below, are examples of such vectors. These vectors contain DNA encoding an interleukin-2 (IL2) signal peptide-*Renilla reniformis* luciferase fusion protein. The IL-2 signal peptide [encoded by the sequence set forth in SEQ ID No. 9] directs secretion of the luciferase protein, to which it is linked, from mammalian cells. Upon secretion from the host mammalian cell, the IL-2 signal peptide is cleaved from the fusion protein to deliver mature, active, luciferase protein to the

-171-

extracellular medium. Successful production and secretion of this heterologous protein can be readily detected by performing luciferase assays which measure the light emitted upon exposure of the medium to the bioluminescent luciferin substrate of the luciferase enzyme.

- 5 Thus, this feature will be useful when artificial chromosomes are used for gene therapy. The presence of a functional artificial chromosome carrying an IL-Ruc fusion with the accompanying therapeutic genes will be readily monitored. Body fluids or tissues can be sampled and tested for luciferase expression by adding luciferin and appropriate cofactors
10 and observing the bioluminescence.

1. Construction of Protein Secretion Vector pLNCX-ILRUC

- Vector pLNCX-ILRUC contains a human IL-2 signal peptide-R. reniformis fusion gene linked to the human cytomegalovirus (CMV) immediate early promoter for constitutive expression of the gene in mammalian cells. The
15 construct was prepared as follows.

a. Preparation of the IL-2 signal sequence-encoding DNA

- A 69-bp DNA fragment containing DNA encoding the human IL-2 signal peptide was obtained through nucleic acid amplification, using appropriate primers for IL-2, of an HEK 293 cell line [see, e.g., U.S.
20 Patent No. 4,518,584 for an IL-2 encoding DNA; see, also SEQ ID No. 9; the IL-2 gene and corresponding amino acid sequence is also provided in the Genbank Sequence Database as accession nos. K02056 and J00264]. The signal peptide includes the first 20 amino acids shown in the translations provided in both of these Genbank entries and in SEQ ID
25 NO. 9. The corresponding nucleotide sequence encoding the first 20 amino acids is also provided in these entries [see, e.g., nucleotides 293-52 of accession no. K02056 and nucleotides 478-537 of accession no. J00264), as well as in SEQ ID NO. 9. The amplification primers included an EcoRI site [GAATTC] for subcloning of the DNA fragment after

-172-

ligation into pGEMT [Promega]. The forward primer is set forth in SEQ ID No. 11 and the sequence of the reverse primer is set forth in SEQ ID No. 12.

TTTGAATTCATGTACAGGATGCAACTCCTG forward [SEQ ID No. 11]

5 TTTGAATTCAGTAGGTGCACTGTTTGTGAC reverse [SEQ ID No. 12]

b. Preparation of the R. reniformis luciferase-encoding DNA

The initial source of the R. reniformis luciferase gene was plasmid pLXSN-RUC. Vector pLXSN [see, e.g., U.S. Patent Nos.

10 5,324,655, 5,470,730, 5,468,634, 5,358,866 and Miller et al. (1989) Biotechniques 7:980] is a retroviral vector capable of expressing heterologous DNA under the transcriptional control of the retroviral LTR; it also contains the neomycin-resistance gene operatively linked for expression to the SV40 early region promoter. The R. reniformis
15 luciferase gene was obtained from plasmid pTZrLuc-1 [see, e.g., U.S. Patent No. 5,292,658; see also the Genbank Sequence Database accession no. M63501; and see also Lorenz et al. (1991) Proc. Natl. Acad. Sci. U.S.A. 88:4438-4442] and is shown as SEQ ID NO. 10. The 0.97 kb EcoRI/SmaI fragment of pTZrLuc-1 contains the coding region of
20 the *Renilla* luciferase-encoding DNA. Vector pLXSN was digested with and ligated with the luciferase gene contained on a pLXSN-RUC, which contains the luciferase gene located operably linked to the viral LTR and upstream of the SV40 promoter, which directs expression of the neomycin-resistance gene.

25 c. Fusion of DNA encoding the IL-2 Signal Peptide and the R. reniformis Luciferase Gene to Yi Id pLXSN-ILRUC

The pGEMT vector containing the IL-2 signal peptide-encoding DNA described in 1.a. above was digested with EcoRI, and the resulting
30 fragment encoding the signal peptide was ligated to EcoRI-digested

-173-

pLXSN-RUC. The resulting plasmid, called pLXSN-ILRUC, contains the IL-2 signal peptide-encoding DNA located immediately upstream of the R. reniformis gene in pLXSN-RUC. Plasmid pLXSN-ILRUC was then used as a template for nucleic acid amplification of the fusion gene in order to
5 add a SmaI site at the 3' end of the fusion gene. The amplification product was subcloned into linearized [EcoRI/SmaI-digested] pGEMT [Promega] to generate ILRUC-pGEMT.

10 d. **Introduction of the Fusion Gene into a Vector
Containing Control Elements for Expression in
Mammalian Cells**

Plasmid ILRUC-pGEMT was digested with KspI and SmaI to release a fragment containing the IL-2 signal peptide-luciferase fusion gene which was ligated to HpaI-digested pLNCX. Vector pLNCX [see, e.g., U.S. Patent Nos. 5,324,655 and 5,457,182; see, also Miller and
15 Rosman (1989) Biotechniques 7:980-990] is a retroviral vector for expressing heterologous DNA under the control of the CMV promoter; it also contains the neomycin-resistance gene under the transcriptional control of a viral promoter. The vector resulting from the ligation reaction was designated pLNCX-ILRUC. Vector pLNCX-ILRUC contains
20 the IL-2 signal peptide-luciferase fusion gene located immediately downstream of the CMV promoter and upstream of the viral 3' LTR and polyadenylation signal in pLNCX. This arrangement provides for expression of the fusion gene under the control of the CMV promoter. Placement of the heterologous protein-encoding DNA [i.e., the luciferase
25 gene] in operative linkage with the IL-2 signal peptide-encoding DNA provides for expression of the fusion in mammalian cells transfected with the vector such that the heterologous protein is secreted from the host cell into the extracellular medium.

-174-

2. Construction of Protein Secretion Targeting Vector pLNCX-ILRUC λ

Vector pLNCX-ILRUC may be modified so that it can be used to introduce the IL-2 signal peptide-luciferase fusion gene into a mammalian artificial chromosome in a host cell. To facilitate specific incorporation of the pLNCX-ILRUC expression vector into a mammalian artificial chromosome, nucleic acid sequences that are homologous to nucleotides present in the artificial chromosome are added to the vector to permit site directed recombination.

Exemplary artificial chromosomes described herein contain λ phage DNA. Therefore, protein secretion targeting vector pLNCX-ILRUC λ was prepared by addition of λ phage DNA [from Charon 4A arms] to produce the secretion vector pLNCX-ILRUC.

3. Expression and Secretion of R. reniformis Luciferase from Mammalian Cells

a. Expression of R. reniformis Luciferase Using pLNCX-ILRUC

Mammalian cells [LMTK⁻ from the ATCC] were transiently transfected with vector pLNCX-ILRUC [~ 10 μ g] by electroporation [BIORAD, performed according to the manufacturer's instructions]. Stable transfectants produced by growth in G418 for neo selection have also been prepared.

Transfectants were grown and then analyzed for expression of luciferase. To determine whether active luciferase was secreted from the transfected cells, culture media were assayed for luciferase by addition of coelentrastazine [see, e.g., Matthews et al. (1977) Biochemistry 16:85-91].

The results of these assays establish that vector pLNCX-ILRUC is capable of providing constitutive expression of heterologous DNA in mammalian host cells. Furthermore, the results demonstrate that the

-175-

human IL-2 signal peptide is capable of directing secretion of proteins fused to the C-terminus of the peptide. Additionally, these data demonstrate that the R. reniformis luciferase protein is a highly effective reporter molecule, which is stable in a mammalian cell environment, and
5 forms the basis of a sensitive, facile assay for gene expression.

b. ***Renilla reniformis* luciferase appears to be secreted from LMTK⁻ cells.**

(i) ***Renilla* luciferase assay of cell pellets**

The following cells were tested:

- 10 cells with no vector: LMTK⁻ cells without vector as a negative control;
- cells transfected with pLNCX only;
- cells transfected with RUC-pLNCX [*Renilla* luciferase gene in pLNCX vector];
- 15 cells transfected with pLNCX-ILRUC [vector containing the IL-2 leader sequence + *Renilla* luciferase fusion gene in pLNCX vector].

Forty-eight hours after electroporation, the cells and culture medium were collected. The cell pellet from 4 plates of cells was resuspended in 1 ml assay buffer and was lysed by sonication. Two
20 hundred μ l of the resuspended cell pellet was used for each assay for luciferase activity [see, e.g., Matthews et al. (1977) Biochemistry 16:85-91]. The assay was repeated three times and the average bioluminescence measurement was obtained.

The results showed that there was relatively low background
25 bioluminescence in the cells transformed with pLNCX or the negative control cells; there was a low level observed in the cell pellet from cells containing the vector with the IL-2 leader sequence-luciferase gene fusion and more than 5000 RLU in the sample from cells containing RUC-pLNCX.

-176-

(ii) **Renilla luciferase assay of cell medium**

Forty milliliters of medium from 4 plates of cells were harvested and spun down. Two hundred microliters of medium was used for each luciferase activity assay. The assay was repeated several times and the average bioluminescence measurement was obtained. These results showed that a relatively high level of bioluminescence was detected in the cell medium from cells transformed with pLNCX-ILRUC; about 10-fold lower levels [slightly above the background levels in medium from cells with no vector or transfected with pLNCX only] was detected in the cells transfected with RUC-pLNCX.

(iii) **conclusions**

The results of these experiments demonstrated that Renilla luciferase appears to be secreted from LMTK⁻ cells under the direction of the IL-2 signal peptide. The medium from cells transfected with Renilla luciferase-encoding DNA linked to the DNA encoding the IL-2 secretion signal had substantially higher levels of Renilla luciferase activity than controls or cells containing luciferase-encoding DNA without the signal peptide-encoding DNA. Also, the differences between the controls and cells containing luciferase encoding-DNA demonstrate that the luciferase activity is specifically from luciferase, not from a non-specific reaction. In addition, the results from the medium of RUC-pLNCX transfected cells, which is similar to background, show that the luciferase activity in the medium does not come from cell lysis, but from secreted luciferase.

c. **Expression of R. reniformis Luciferase Using pLNCX-ILRUC λ**

To express the IL-2 signal peptide-R. reniformis fusion gene from an mammalian artificial chromosome, vector pLNCX-ILRUC λ is targeted for site-specific integration into a mammalian artificial chromosome through homologous recombination of the λ DNA sequences contained in

-177-

the chromosome and the vector. This is accomplished by introduction of pLNCX-ILRUC λ into either a fusion cell line harboring mammalian artificial chromosomes or mammalian host cells that contain mammalian artificial chromosomes. If the vector is introduced into a fusion cell line harboring

5 the artificial chromosomes, for example through microinjection of the vector or transfection of the fusion cell line with the vector, the cells are then grown under selective conditions. The artificial chromosomes, which have incorporated vector pLNCX-ILRUC λ , are isolated from the surviving cells, using purification procedures as described above, and

10 then injected into the mammalian host cells.

Alternatively, the mammalian host cells may first be injected with mammalian artificial chromosomes which have been isolated from a fusion cell line. The host cells are then transfected with vector pLNCX-ILRUC λ and grown.

15 The recombinant host cells are then assayed for luciferase expression as described above.

F. Other targeting vectors

These vectors, which are based on vector pMCT-RUC, rely on positive and negative selection to insure insertion and selection for the

20 double recombinants. A single crossover results in incorporation of the DT-A, which kills the cell, double crossover recombinations delete the DT-1 gene.

1. Plasmid pNEM1 contains:

- | | |
|---------|--|
| DT-A: | Diphtheria toxin gene (negative selectable marker) |
| 25 Hyg: | Hygromycin gene (positive selectable marker) |
| ruc: | <u>Renilla</u> luciferase gene (non-selectable marker) |
| 1: | LTR-MMTV promoter |
| 2: | TK promoter |
| 3: | CMV promoter |

-178-

MMR: Homology region (plasmid pAG60)

2. plasmid pNEM-2 and -3 are similar to pNEM 1 except for different negative selectable markers:

pNEM-1: diphtheria toxin gene as "—" selectable marker

5 pNEM-2: hygromycin antisense gene as "—" selectable marker

pNEM-3: thymidine kinase HSV-1 gene as "—" selectable marker

3. Plasmid - λ DNA based homology:

pNEM λ -1: base vector

pNEM λ -2: base vector containing p5 = gene

10 1: LTR MMTV promoter

2: SV40 promoter

3: CMV promoter

4: μ TIIA promoter (metallothionein gene promoter)

— homology region (plasmid pAG60)

15 λ L.A. and λ R.A. homology regions for λ left and right arms (λ gt-WES).

EXAMPLE 13

Microinjection of mammalian cells with plasmid DNA

20 These procedures will be used to microinject MACs into eukaryotic cells, including mammalian and insect cells.

The microinjection technique is based on the use of small glass capillaries as a delivery system into cells and has been used for introduction of DNA fragments into nuclei [see, e.g., Chalfie et al. (1994) Science 263:802-804]. It allows the transfer of almost any type of

25 molecules, e.g., hormones, proteins, DNA and RNA, into either the cytoplasm or nuclei of recipient cells. This technique has no cell type restriction and is more efficient than other methods, including Ca^{2+} —mediated gene transfer and liposome-mediated gene transfer. About 20-30% of the injected cells become successfully transformed.

-179-

Microinjection is performed under a phase-contrast microscope. A glass microcapillary, prefilled with the DNA sample, is directed into a cell to be injected with the aid of a micromanipulator. An appropriate sample volume (1-10 pl) is transferred into the cell by gentle air pressure exerted
5 by a transjector connected to the capillary. Recipient cells are grown on glass slides imprinted with numbered squares for convenient localization of the injected cells.

a. Materials and equipment

Nunclon tissue culture dishes 35 x 10 mm, mouse cell line EC3/7C5
10 Plasmid DNA pCH110 [Pharmacia], Purified Green Florescent Protein (GFP) [GFPs from *Aequorea* and *Renilla* have been purified and also DNA encoding GFPs has been cloned; see, e.g., Prasher et al. (1992) Gene 111:229-233; International PCT Application No. WO 95/07463, which is based on U.S. application Serial No. 08/119,678 and U.S. application
15 Serial No. 08/192,274], ZEISS Axiovert 100 microscope, Eppendorf transjector 5246, Eppendorf micromanipulator 5171, Eppendorf Cellocate coverslips, Eppendorf microloaders, Eppendorf femtotips and other standard equipment

b. Protocol for injecting

20 (1) Fibroblast cells are grown in 35 mm tissue culture dishes (37° C, 5% CO₂) until the cell density reaches 80% confluency. The dishes are removed from the incubator and medium is added to about a 5 mm depth.

(2) The dish is placed onto the dish holder
25 and the cells observed with 10 x objective; the focus is desirably above the cell surface.

(3) Plasmid or chromosomal DNA solution [1 ng/ μ l] and GFP protein solution are further purified by centrifuging the

-180-

DNA sample at a force sufficient to remove any particular debris [typically about 10,000 rpm for 10 minutes in a microcentrifuge].

(4) Two 2 μ l of the DNA solution (1 ng/ μ l) is loaded into a microcapillary with an Eppendorf microloader. During
5 loading, the loader is inserted to the tip end of the microcapillary. GFP (1 mg/ml) is loaded with the same procedure.

(5) The protecting sheath is removed from the microcapillary and the microcapillary is fixed onto the capillary holder connected with the micromanipulator.

10 (6) The capillary tip is lowered to the surface of the medium and is focussed on the cells gradually until the tip of the capillary reaches the surface of a cell. The capillary is lowered further so that the it is inserted into the cell. Various parameters, such as the level
15 of the capillary, the time and pressure, are determined for the particular equipment. For example, using the fibroblast cell line C5 and the above-noted equipment, the best conditions are: injection time 0.4 second, pressure 80 psi. DNA can then be automatically injected into the nuclei of the cells.

(7) After injection, the cells are returned to
20 the incubator, and incubated for about 18-24 hours.

(8) After incubation the number of transformants can be determined by a suitable method, which depends upon the selection marker. For example, if green fluorescent protein is used, the assay can be performed using UV light source and fluorescent
25 filter set at 0-24 hours after injection. If β -gal-containing DNA, such as DNA-derived from pHC110, has been injected, then the transformants can be assayed for β -gal.

-181-

(c) **Detection of β -galactosidase in cells injected with plasmid DNA**

The medium is removed from the culture plate and the cells are fixed by addition of 5 ml of fixation Solution I: (1% glutaraldehyde; 0.1 M sodium phosphate buffer, pH 7.0; 1 mM MgCl_2), and incubated for 15 minutes at 37° C. Fixation Solution I is replaced with 5 ml of X-gal Solution II: [0.2% X-gal, 10 mM sodium phosphate buffer (pH 7.0), 150 mM NaCl, 1 mM MgCl_2 , 3.3 mM $\text{K}_4\text{Fe}(\text{CN})_6\text{H}_2\text{O}$, 3.3 mM $\text{K}_3\text{Fe}(\text{CN})_6$], and the plates are incubated for 30-60 minutes at 37° C. The X-gal solution is removed and 2 ml of 70% glycerol is added to each dish. Blue stained cells are identified under a light microscope.

This method will be used to introduce a MAC, particularly the MAC with the anti-HIV megachromosome, to produce a mouse model for anti-HIV activity.

15

EXAMPLE 14

Transgenic (non-human) animals

Transgenic (non-human) animals can be generated that express heterologous genes which confer desired traits, e.g., disease resistance, in the animals. A transgenic mouse is prepared to serve as a model of a disease-resistant animal. Genes that encode vaccines or that encode therapeutic molecules can be introduced into embryos or ES cells to produce animals that express the gene product and thereby are resistant to or less susceptible to a particular disorder.

The mammalian artificial megachromosome and others of the artificial chromosomes, particularly the SATACs, can be used to generate transgenic (non-human) animals, including mammals and birds, that stably express genes conferring desired traits, such as genes conferring resistance to pathogenic viruses. The artificial chromosomes can also be

-182-

used to produce transgenic (non-human) animals, such as pigs, that can produce immunologically humanized organs for xenotransplantation.

For example, transgenic mice containing a transgene encoding an anti-HIV ribozyme provide a useful model for the development of stable transgenic (non-human) animals using these methods. The artificial chromosomes can be used to produce transgenic (non-human) animals, particularly, cows, goats, mice, oxen, camels, pigs and sheep, that produce the proteins of interest in their milk; and to produce transgenic chickens and other egg-producing fowl, that produce therapeutic proteins or other proteins of interest in their eggs. For example, use of mammary gland-specific promoters for expression of heterologous DNA in milk is known [see, e.g. U.S. Patent No. 4,873,316]. In particular, a milk-specific promoter or a promoter, preferably linked to a milk-specific signal peptide, specifically activated in mammary tissue is operatively linked to the DNA of interest, thereby providing expression of that DNA sequence in milk.

1. Development of Control Transgenic Mice Expressing Anti-HIV Ribozyme

Control transgenic mice are generated in order to compare stability and amounts of transgene expression in mice developed using transgene DNA carried on a vector (control mice) with expression in mice developed using transgenes carried in an artificial megachromosome.

a. Development of Control Transgenic Mice Expressing β -galactosidase

One set of control transgenic mice was generated by microinjection of mouse embryos with the β -galactosidase gene alone. The microinjection procedure used to introduce the plasmid DNA into the mouse embryos is as described in Example 13, but modified for use with embryos [see, e.g., Hogan et al. (1994) *Manipulating the Mouse Embryo, A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring

-183-

Harbor, NY, see, especially pages 255-264 and Appendix 3]. Fertilized mouse embryos [Strain CB6 obtained from Charles River Co.] were injected with 1 ng of plasmid pCH110 (Pharmacia) which had been linearized by digestion with BamHI. This plasmid contains the β -galactosidase gene linked to the SV40 late promoter. The β -galactosidase gene product provides a readily detectable marker for successful transgene expression. Furthermore, these control mice provide confirmation of the microinjection procedure used to introduce the plasmid into the embryos. Additionally, because the megachromosome that is transferred to the mouse embryos in the model system (see below) also contains the β -galactosidase gene, the control transgenic mice that have been generated by injection of pCH110 into embryos serve as an analogous system for comparison of heterologous gene expression from a plasmid versus from a gene carried on an artificial chromosome.

After injection, the embryos are cultured in modified HTF medium under 5% CO₂ at 37°C for one day until they divide to form two cells. The two-cell embryos are then implanted into surrogate mother female mice [for procedures see, Manipulating the Mouse Embryo, A Laboratory Manual (1994) Hogan et al., eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pp. 127 et seq.].

b. Development of Control Transgenic Mice Expressing Anti-HIV Ribozyme

One set of anti-HIV ribozyme gene-containing control transgenic mice was generated by microinjection of mouse embryos with plasmid pCEPUR-132 which contains three different genes: (1) DNA encoding an anti-HIV ribozyme, (2) the puromycin-resistance gene and (3) the hygromycin-resistance gene. Plasmid pCEPUR-132 was constructed by ligating portions of plasmid pCEP-132 containing the anti-HIV ribozyme

-184-

gene (referred to as ribozyme D by Chang *et al.* [(1990) *Clin. Biotech.* 2:23-31]; see also U.S. Patent No. 5,144,019 to Rossi *et al.*, particularly Figure 4 of the patent) and the hygromycin-resistance gene with a portion of plasmid pCEPUR containing the puromycin-resistance gene.

5 Plasmid pCEP-132 was constructed as follows. Vector pCEP4 (Invitrogen, San Diego, CA; see also Yates *et al.* (1985) *Nature* 313:812-815) was digested with *Xho*I which cleaves in the multiple cloning site region of the vector. This ~10.4-kb vector contains the hygromycin-resistance gene linked to the thymidine kinase gene promoter and
10 polyadenylation signal, as well as the ampicillin-resistance gene and ColE1 origin of replication and EBNA-1 (Epstein-Barr virus nuclear antigen) genes and OriP. The multiple cloning site is flanked by the cytomegalovirus promoter and SV40 polyadenylation signal.

*Xho*I-digested pCEP4 was ligated with a fragment obtained by
15 digestion of plasmid 132 (see Example 4 for a description of this plasmid) with *Xho*I and *Sal*I. This *Xho*I/*Sal*I fragment contains the anti-HIV ribozyme gene linked at the 3' end to the SV40 polyadenylation signal. The plasmid resulting from this ligation was designated pCEP-132. Thus, in effect, pCEP-132 comprises pCEP4 with the anti-HIV ribozyme gene
20 and SV40 polyadenylation signal inserted in the multiple cloning site for CMV promoter-driven expression of the anti-HIV ribozyme gene.

 To generate pCEPUR-132, pCEP-132 was ligated with a fragment of pCEPUR. pCEPUR was prepared by ligating a 7.7-kb fragment generated upon *Nhe*I/*Nru*I digestion of pCEP4 with a 1.1-kb *Nhe*I/*Sna*BI
25 fragment of pBabe [see Morgenstern and Land (1990) *Nucleic Acids Res.* 18:3587-3596 for a description of pBabe] that contains the puromycin-resistance gene linked at the 5' end to the SV40 promoter. Thus, pCEPUR is made up of the ampicillin-resistance and EBNA1 genes, as well as the ColE1 and OriP elements from pCEP4 and the puromycin-

-185-

resistance gene from pBabe. The puromycin-resistance gene in pCEPUR is flanked by the SV40 promoter (from pBabe) at the 5' end and the SV40 polyadenylation signal (from pCEP4) at the 3' end.

Plasmid pCEPUR was digested with XhoI and Sall and the
5 fragment containing the puromycin-resistance gene linked at the 5' end to the SV40 promoter was ligated with XhoI-digested pCEP-132 to yield the ~12.1-kb plasmid designated pCEPUR-132. Thus, pCEPUR-132, in effect, comprises pCEP-132 with puromycin-resistance gene and SV40 promoter inserted at the XhoI site. The main elements of pCEPUR-132
10 are the hygromycin-resistance gene linked to the thymidine kinase promoter and polyadenylation signal, the anti-HIV ribozyme gene linked to the CMV promoter and SV40 polyadenylation signal, and the puromycin-resistance gene linked to the SV40 promoter and polyadenylation signal. The plasmid also contains the ampicillin-
15 resistance and EBNA1 genes and the ColE1 origin of replication and OriP.

Zygotes were prepared from (C57BL/6JxCBA/J) F1 female mice [see, e.g., Manipulating the Mouse Embryo, A Laboratory Manual (1994) Hogan et al., eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, p. 429], which had been previously mated with a
20 (C57BL/6JxCBA/J) F1 male. The male pronuclei of these F2 zygotes were injected [see, Manipulating the Mouse Embryo, A Laboratory Manual (1994) Hogan et al., eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY] with pCEPUR-132 (~3 µg/ml), which had been linearized by digestion with NruI. The injected eggs were then implanted
25 in surrogate mother female mice for development into transgenic offspring.

These primary carrier offspring were analyzed (as described below) for the presence of the transgene in DNA isolated from tail cells. Seven carrier mice that contained transgenes in their tail cells (but that may not

-186-

carry the transgene in all their cells, i.e., they may be chimeric) were allowed to mate to produce non-chimeric or germ-line heterozygotes. The heterozygotes were, in turn, crossed to generate homozygote transgenic offspring.

5 2. **Development of Model Transgenic Mice Using Mammalian Artificial Chromosomes**

Fertilized mouse embryos are microinjected (as described above) with megachromosomes (1-10 pL containing 0-1 chromosomes/pL) isolated from fusion cell line G3D5 or H1D3 (described above). The
10 megachromosomes are isolated as described herein. Megachromosomes isolated from either cell line carry the anti-HIV ribozyme (ribozyme D) gene as well as the hygromycin-resistance and β -galactosidase genes. The injected embryos are then developed into transgenic mice as described above.

15 Alternatively, the megachromosome-containing cell line G3D5* or H1D3* is fused with mouse embryonic stem cells [see, e.g., U.S. Patent No. 5,453,357, commercially available; see Manipulating the Mouse Embryo, A Laboratory Manual (1994) Hogan et al., eds., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, pages 253-289]
20 following standard procedures see also, e.g., Guide to Techniques in Mouse Development in Methods in Enzymology Vol. 25, Wassarman and De Pamphilis, eds. (1993), pages 803-932]. (It is also possible to deliver isolated megachromosomes into embryonic stem cells using the Microcell procedure [such as that described above].) The stem cells are cultured in
25 the presence of a fibroblast [e.g., STO fibroblasts that are resistant to hygromycin and puromycin]. Cells of the resultant fusion cell line, which contains megachromosomes carrying the transgenes [i.e., anti-HIV ribozyme, hygromycin-resistance and β -galactosidase genes], are then transplanted into mouse blastocysts, which are in turn implanted into a

-187-

surrogate mother female mouse where development into a transgenic mouse will occur.

Mice generated by this method are chimeric; the transgenes will be expressed in only certain areas of the mouse, e.g., the head, and thus
5 may not be expressed in all cells.

3. Analysis of Transgenic Mice for Transgene Expression

Beginning when the transgenic mice, generated as described above, are three-to-four weeks old, they can be analyzed for stable expression of the transgenes that were transferred into the embryos [or
10 fertilized eggs] from which they develop. The transgenic mice may be analyzed in several ways as follows.

a. Analysis of Cells Obtained from the Transgenic Mice

Cell samples [e.g., spleen, liver and kidney cells, lymphocytes, tail
15 cells] are obtained from the transgenic mice. Any cells may be tested for transgene expression. If, however, the mice are chimeras generated by microinjection of fertilized eggs or by fusion of embryonic stem cells with megachromosome-containing cells, only cells from areas of the mouse that carry the transgene are expected to express the transgene. If the
20 cells survive growth on hygromycin [or hygromycin and puromycin or neomycin, if the cells are obtained from mice generated by transfer of both antibiotic-resistance genes], this is one indication that they are stably expressing the transgenes. RNA isolated from the cells according to standard methods may also be analyzed by northern blot procedures
25 to determine if the cells express transcripts that hybridize to nucleic acid probes based on the antibiotic-resistance genes. Additionally, cells obtained from the transgenic mice may also be analyzed for β -galactosidase expression using standard assays for this marker enzyme [for example, by direct staining of the product of a reaction involving β -

-188-

galactosidase and the X-gal substrate, see, e.g., Jones (1986) EMBO 5:3133-3142, or by measurement of β -galactosidase activity, see, e.g., Miller (1972) in Experiments in Molecular Genetics pp. 352-355, Cold Spring Harbor Press]. Analysis of β -galactosidase expression is

- 5 particularly used to evaluate transgene expression in cells obtained from control transgenic mice in which the only transgene transferred into the embryo was the β -galactosidase gene.

Stable expression of the anti-HIV ribozyme gene in cells obtained from the transgenic mice may be evaluated in several ways. First, DNA

10 isolated from the cells according to standard procedures may be subjected to nucleic acid amplification using primers corresponding to the ribozyme gene sequence. If the gene is contained within the cells, an amplified product of pre-determined size is detected upon hybridization of the reaction mixture to a nucleic acid probe based on the ribozyme gene

15 sequence. Furthermore, DNA isolated from the cells may be analyzed using Southern blot methods for hybridization to such a nucleic acid probe. Second, RNA isolated from the cells may be subjected to northern blot hybridization to determine if the cells express RNA that hybridizes to nucleic acid probes based on the ribozyme gene. Third, the

20 cells may be analyzed for the presence of anti-HIV ribozyme activity as described, for example, in Chang et al. (1990) Clin. Biotech. 2:23-31. In this analysis, RNA isolated from the cells is mixed with radioactively labeled HIV gag target RNA which can be obtained by in vitro transcription of gag gene template under reaction conditions favorable to

25 in vitro cleavage of the gag target, such as those described in Chang et al. (1990) Clin. Biotech. 2:23-31. After the reaction has been stopped, the mixture is analyzed by gel electrophoresis to determine if cleavage products smaller in size than the whole template are detected; presence

-189-

of such cleavage fragments is indicative of the presence of stably expressed ribozyme.

b. Analysis of Whole Transgenic Mice

Whole transgenic mice that have been generated by transfer of the
5 anti-HIV ribozyme gene [as well as selection and marker genes] into
embryos or fertilized eggs can additionally be analyzed for transgene
expression by challenging the mice with infection with HIV. It is possible
for mice to be infected with HIV upon intraperitoneal injection with
high-producing HIV-infected U937 cells [see, e.g., Locardi et al. (1992)
10 J. Virol. 66:1649-1654]. Successful infection may be confirmed by
analysis of DNA isolated from cells, such as peripheral blood
mononuclear cells, obtained from transgenic mice that have been injected
with HIV-infected human cells. The DNA of infected transgenic mice
cells will contain HIV-specific gag and env sequences, as demonstrated
15 by, for example, nucleic acid amplification using HIV-specific primers. If
the cells also stably express the anti-HIV ribozyme, then analysis of RNA
extracts of the cells should reveal the smaller gag fragments arising by
cleavage of the gag transcript by the ribozyme.

Additionally, the transgenic mice carrying the anti-HIV ribozyme
20 gene can be crossed with transgenic mice expressing human CD4 (i.e.,
the cellular receptor for HIV) [see Gillespie et al. (1993) Mol. Cell. Biol.
13:2952-2958; Hanna et al. (1994) Mol. Cell. Biol. 14:1084-1094; and
Yeung et al. (1994) J. Exp. Med. 180:1911-1920, for a description of
transgenic mice expressing human CD4]. The offspring of these crossed
25 transgenic mice expressing both the CD4 and anti-HIV ribozyme
transgenes should be more resistant to infection [as a result of a
reduction in the levels of active HIV in the cells] than mice expressing
CD4 alone [without expressing anti-HIV ribozyme].

-190-

4. Development of transgenic chickens using artificial chromosomes

The development of transgenic chickens has many applications in the improvement of domestic poultry, an agricultural species of commercial significance, such as disease resistance genes and genes encoding therapeutic proteins. It appears that efforts in the area of chicken transgenesis have been hampered due to difficulty in achieving stable expression of transgenes in chicken cells using conventional methods of gene transfer via random introduction into recipient cells. Artificial chromosomes are, therefore, particularly useful in the development of transgenic chickens because they provide for stable maintenance of transgenes in host cells.

a. Preparation of artificial chromosomes for introduction of transgenes into recipient chicken cells

(i) Mammalian artificial chromosomes

Mammalian artificial chromosomes, such as the SATACs and minichromosomes described herein, can be modified to incorporate detectable reporter genes and/or transgenes of interest for use in developing transgenic chickens. Alternatively, chicken-specific artificial chromosomes can be constructed using the methods herein. In particular, chicken artificial chromosomes [CACs] can be prepared using the methods herein for preparing MACs; or, as described above, the chicken libraries can be introduced into MACs provided herein and the resulting MACs introduced into chicken cells and those that are functional in chicken cells selected.

As described in Examples 4 and 7, and elsewhere herein, artificial chromosome-containing mouse LMTK⁻-derived cell lines, or minichromosome-containing cell lines, as well as hybrids thereof, can be transfected with selected DNA to generate MACs [or CACs] that have

-191-

integrated the foreign DNA for functional expression of heterologous genes contained within the DNA.

To generate MACs or CACs containing transgenes to be expressed in chicken cells, the MAC-containing cell lines may be transfected with
5 DNA that includes λ DNA and transgenes of interest operably linked to a promoter that is capable of driving expression of genes in chicken cells. Alternatively, the minichromosomes or MACs [or CACs], produced as described above, can be isolated and introduced into cells, followed by targeted integration of selected DNA. Vectors for targeted integration
10 are provided herein or can be constructed as described herein.

Promoters of interest include constitutive, inducible and tissue (or cell)-specific promoters known to those of skill in the art to promote expression of genes in chicken cells. For example, expression of the lacZ gene in chicken blastodermal cells and primary chicken fibroblasts has
15 been demonstrated using a mouse heat-shock protein 68 (*hsp 68*) promoter [phspPTlacZpA; see Brazolot et al. (1991) Mol. Reprod. Devel. 30:304-312], a Zn²⁺-inducible chicken metallothionein (cMt) promoter [pCBcMtlacZ; see Brazolot et al. (1991) Mol. Reprod. Devel. 30:304-312], the constitutive Rous sarcoma virus and chicken β -actin promoters
20 in tandem [pmiwZ; see Brazolot et al. (1991) Mol. Reprod. Devel. 30:304-312] and the constitutive cytomegalovirus (CMV) promoter. Of particular interest herein are egg-specific promoters that are derived from genes, such as ovalbumin and lysozyme, that are expressed in eggs.

The choice of promoter will depend on a variety of factors,
25 including, for example, whether the transgene product is to be expressed throughout the transgenic chicken or restricted to certain locations, such as the egg. Cell-specific promoters functional in chickens include the steroid-responsive promoter of the egg ovalbumin protein-encoding gene [see Gaub et al. (1987) EMBO J. 6:2313-2320; Tora et al. (1988) EMBO

-192-

J. 7:3771-3778; Park et al. (1995) Biochem. Mol. Biol. Int. (Australia) 36:811-816].

(ii) **Chicken artificial chromosomes**

Additionally, chicken artificial chromosomes may be generated
5 using methods described herein. For example, chicken cells, such as
primary chicken fibroblasts [see Brazolot et al. (1991) Mol. Reprod.
Devel. 30:304-312], may be transfected with DNA that encodes a
selectable marker [such as a protein that confers resistance to
antibiotics] and that includes DNA (such as chicken satellite DNA) that
10 targets the introduced DNA to the pericentric region of the endogenous
chicken chromosomes. Transfectants that survive growth on selection
medium are then analyzed, using methods described herein, for the
presence of artificial chromosomes, including minichromosomes, and
particularly SATACs. An artificial chromosome-containing transfectant
15 cell line may then be transfected with DNA encoding the transgene of
interest [fused to an appropriate promoter] along with DNA that targets
the foreign DNA to the chicken artificial chromosome.

b. **Introduction of artificial chromosomes carrying
transgenes of interest into recipient chicken cells**

20 Cell lines containing artificial chromosomes that harbor
transgene(s) of interest (i.e., donor cells) may be fused with recipient
chicken cells in order to transfer the chromosomes into the recipient
cells. Alternatively, the artificial chromosomes may be isolated from the
donor cells, for example, using methods described herein [see, e.g.,
25 Example 10], and directly introduced into recipient cells.

Exemplary chicken recipient cell lines include, but are not limited
to, stage X blastoderm cells [see, e.g., Brazolot et al. (1991) Mol.
Reprod. Dev. 30:304-312; Etches et al. (1993) Poultry Sci. 72:882-889;

-193-

Petitte *et al.* (1990) Development 108:185-189] and chick zygotes [see, *e.g.*, Love *et al.* (1994) Biotechnology 12:60-63].

For example, microcell fusion is one method for introduction of artificial chromosomes into avian cells [see, *e.g.*, Dieken *et al.* [(1996) 5 Nature Genet. 12:174-182 for methods of fusing microcells with DT40 chicken pre-B cells]. In this method, microcells are prepared [for example, using procedures described in Example 1.A.5] from the artificial chromosome-containing cell lines and fused with chicken recipient cells.

Isolated artificial chromosomes may be directly introduced into 10 chicken recipient cell lines through, for example, lipid-mediated carrier systems, such as lipofection procedures [see, *e.g.*, Brazolot *et al.* (1991) Mol. Reprod. Dev. 30:304-312] or direct microinjection. Microinjection is generally preferred for introduction of the artificial chromosomes into chicken zygotes [see, *e.g.*, Love *et al.* (1994) Biotechnology 12:60-63].

15 c. Development of transgenic chickens

Transgenic chickens may be developed by injecting recipient Stage X blastoderm cells (which have received the artificial chromosomes) into embryos at a similar stage of development [see, *e.g.*, Etches *et al.* (1993) Poultry Sci. 72:882-889; Petitte *et al.* (1990) Development 20 108:185-189; and Carsience *et al.* (1993) Development 117: 669-675]. The recipient chicken embryos within the shell are candled and allowed to hatch to yield a germline chimeric chicken that will express the transgene(s) in some of its cells.

Alternatively, the artificial chromosomes may be introduced into 25 chick zygotes, for example through direct microinjection [see, *e.g.*, Love *et al.* (1994) Biotechnology 12:60-63], which thereby are incorporated into at least a portion of the cells in the chicken. Inclusion of a tissue-specific promoter, such as an egg-specific promoter, will ensure appropriate expression of operatively-linked heterologous DNA.

-194-

The DNA of interest may also be introduced into a minichromosome, by methods provided herein. The minichromosome may either be one provided herein, or one generated in chicken cells using the methods herein. The heterologous DNA will be introduced
5 using a targeting vector, such as those provided herein, or constructed as provided herein.

Since modifications will be apparent to those of skill in this art, it is intended that this invention be limited only by the scope of the
10 appended claims.

-195-

SEQUENCE LISTING

(1) GENERAL INFORMATION

- (i) APPLICANT:
 (A) NAME: The Biological Research Center of the Hungarian Academy of Sciences
 (B) STREET: Post Office Box 521
 (C) CITY: H-6701 Szeged
 (D) STATE:
 (E) COUNTRY: Hungary
 (F) POSTAL CODE (ZIP):
- (i) APPLICANT:
 (A) NAME: Loma Linda University
 (B) STREET:
 (C) CITY: Loma Linda
 (D) STATE: California
 (E) COUNTRY: USA
 (F) POSTAL CODE (ZIP): 92350
- (i) APPLICANT:
 (A) NAME: AMERICAN GENE THERAPY, INC.
 (B) STREET: 5022 154 Street
 (C) CITY: Edmonton, T6H5PE
 (D) STATE:
 (E) COUNTRY: CANADA
 (F) POSTAL CODE (ZIP):
- (i) INVENTOR/APPLICANT:
 (A) NAME: Gyula Hadlaczky
 (B) STREET: H.6723, Szeged
 (C) CITY: Szamos U.I.A. IX 36
 (D) STATE:
 (D) COUNTRY: Hungary
 (E) POSTAL CODE (ZIP):
- (i) INVENTOR/APPLICANT:
 (A) NAME: Aladar Szalay
 - (B) STREET: 7327 Fairwood
 (C) CITY: Highland
 (D) STATE: California
 (D) COUNTRY: USA
 (E) POSTAL CODE (ZIP): 92346
- (ii) TITLE OF THE INVENTION: ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS FOR PREPARING ARTIFICIAL CHROMOSOMES
- (iii) NUMBER OF SEQUENCES: 34
- (iv) CORRESPONDENCE ADDRESS:
 (A) ADDRESSEE: Brown, Martin, Haller & McClain
 (B) STREET: 1660 Union Street
 (C) CITY: San Diego
 (D) STATE: CA
 (E) COUNTRY: USA
 (F) ZIP: 92101-2926

-196-

- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette
 - (B) COMPUTER: IBM Compatible
 - (C) OPERATING SYSTEM: DOS
 - (D) SOFTWARE: FastSEQ Version 1.5
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER:
 - (B) FILING DATE: 10-04-1997
 - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/695,191
 - (B) FILING DATE: 07-AUG-1996
- (vi) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/682,080
 - (B) FILING DATE: 15-JUL-1996
 - (C) CLASSIFICATION:
- (vi) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/629,822
 - (B) FILING DATE: 10-APR-1996
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Seidman, Stephanie L
 - (B) REGISTRATION NUMBER: 33,779
 - (C) REFERENCE/DOCKET NUMBER: 6869-402PC
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 619-238-0999
 - (B) TELEFAX: 619-238-0062
 - (C) TELEX:
- (2) INFORMATION FOR SEQ ID NO:1:
 - (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 1293 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
 - (ii) MOLECULE TYPE: Genomic DNA
 - (iii) HYPOTHETICAL: NO
 - (iv) ANTISENSE: NO
 - (v) FRAGMENT TYPE:
 - (vi) ORIGINAL SOURCE:
 - (ix) FEATURE:
- (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

GAATTCATCA	TTTTTCANGT	CCTCAAGTGG	ATGTTTCTCA	TTTNCCATGA	TTTTAAGTTT	60
TCTCGCCATA	TTCTGGTCC	TACAGTGTGC	ATTTCTCCAT	TTNCACGTT	TTNCAGTGAT	120
TTCGTCAATTT	TCAAGTCCTC	AAGTGGATGT	TTCTCATTTN	CCATGAATTT	CAGTTTTCTN	180
GCCATATTCC	ACGTCCTACA	GNGGACATTT	CTAAATTTNC	CACCTTTTTC	AGTTTTCTC	240
GCCATATTTT	ACGTCCTAAA	ATGTGTATTT	CTCGTTTNCC	GTGATTTTCA	GTTTTCTCGC	300
CAGATTCAG	GTCTATAAT	GTGCATTTCT	CATTTNNCAC	GTTTTTCAGT	GATTTTCGTCA	360
TTTTTTCAAG	TCGGCAAGTG	GATGTTTCTC	ATTTNCCATG	ATTTNCAGTT	TTCTTGNAAT	420
ATTCCATGTC	CTACAATGAT	CATTTTAAAT	TTTCCACCTT	TTCATTTTTC	CACGCCATAT	480

-197-

TTCATGTCCT	AAAGTGTATA	TTTCTCCTTT	TCCGCGATTT	TCAGTTTTCT	CGCCATATTC	540
CAGGTCCTAC	AGTGTGCATT	CCTCATTTTT	CACCTTTTTT	ACTGATTTCG	TCATTTTTCA	600
AGTCGTCAAC	TGGATCTTTC	TAATTTTCCA	TGATTTTCAG	TTATCTTGTC	ATATTCCATG	660
TCCTACAGTG	GACATTTCTA	AATTTTCCAA	CTTTTTCAAT	TTTTCTCGAC	ATATTGACG	720
TGCTAAAGTG	TGTATTTCTT	ATTTTCCGTG	ATTTTCAGTT	TTCTCGCCAT	ATTCCAGGTC	780
CTAATAGTGT	GCATTTCTCA	TTTTTCACGT	TTTTTCAGTGA	TTTCGTCAAT	TTTTCCAGTT	840
GTCAAGGGGA	TGTTTCTCAT	TTTCCATGAG	TGTCAGTTTT	CTTGCTATAT	TCCATGTCCT	900
ACAGTGACAT	TTCTAAATAT	TATACCTTTT	TCAGTTTTTC	TCACCATATT	TCACGTCCTA	960
AAGTATATAT	TTCTCATTTT	CCCTGATTTT	CAGTTTCCTT	GCCATATTCC	AGGTCCTACA	1020
GTGTGCATTT	CTCATTTTTC	ACGTTTTTCA	GTAATTTCTT	CATTTTTTAA	GCCCTCAAAT	1080
GGATGTTTCT	CATTTTCCAT	GATTTTCAGT	TTTCTTGCCA	TATACCATGT	CCTACAGTGG	1140
ACATTTCTAA	ATTATCCACC	TTTTTCAGTT	TTTCATCGGC	ACATTTACAG	TCCTAAAGTG	1200
TGTATTTCTA	ATTTTCAGTG	ATTTTCAGTT	TTCTCGCCAT	ATTCCAGGAC	CTACAGTGTG	1260
CATTTCTCAT	TTTTTCACGT	TTTCAGTGAA	TTC			1293

(2) INFORMATION FOR SEQ ID NO:2:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1044 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

AGGCCATATGG	TGAAAAAGGA	AATATCTTCC	CCTGAAAAC	AGACAGAAGG	ATTCTCAGAA	60
TCTTATTTGT	GATGTGCGCC	CCTCAACTAA	CAGTGTTGAA	GCTTCTTTT	GATAGAGCAG	120
TTTTGAAACA	CTCTTTTGT	AAAATCTGCA	AGAGGATATT	TGGATAGCTT	TGAGGATTTT	180
CGTTGGAAAC	GGGATTGTCT	TCATATAAAC	CCTAGACAGA	AGCATTCTCA	GAAGCTTCAT	240
TGGGATGTTT	CAGTTGAAGT	CACAGTGTGG	AACAGTCCCC	TTTCATAGAG	CAGGTTTGAA	300
ACACTGTTTT	TTGTAGTATC	TGGAAGTGGA	CATTTGGAGC	GATCTCAGGA	CTGCGGTGAA	360
AAAGGAAATA	TCTTCCAATA	AAAGCTAGAT	AGAGGCAATG	TCAGAAACCT	TTTTCATGAT	420
GTATCTACTC	AGCTAACAGA	GTTGAACCTT	CCTTTGAGAG	AGCAGTTTTG	AAACACTCTT	480
TTTGTGGAAAT	CTGCAAGTGG	ATATTTGTCT	AGCTTTGAGG	ATTTCGTTGG	GAAACGGGAT	540
TACATATAAA	AAGCAGACAG	CAGCATTCCC	AGAAACTTCT	TTGTGATGTT	TGCATTCAAG	600
TCACAGAGTT	GAACATTCCC	TTTCATAGAG	CAGGTTTGAA	ACACACTTTT	TGATGTATCT	660
GGATGTGGAC	ATTTGCAGCG	CTTTCAGGCC	TAAGGTGAAA	AGGAAATATC	TTCCCCTGAA	720
AACTAGACAG	AAGCATTCTC	AGAAACTTAT	TTGTGATGTG	CGCCCTCAAC	TAACAGTGTG	780
GAAGCTTTCT	TTTGATAGAG	GCAGTTTTGA	AACACTCTTT	TGTGGAATCT	GCAAGTGGAT	840
ATTTGTCTAG	CTTTGAGGAT	TTCTTTGGAA	ACGGGATTAC	ATATAAAAAG	CAGACAGCAG	900
CATTCCCAGA	ATCTTGTTTG	TGATGTTTGC	ATTCAAGTCA	CAGAGTTGAA	CATTCCCTTT	960
CAGAGAGCAG	GTTTGAACAC	TCTTTTATA	GTATCTGGAT	GTGGACATTT	GGAGCGCTTT	1020
CAGGGGGGAT	CCTCTAGAAT	TCCT				1044

(2) INFORMATION FOR SEQ ID NO:3:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 2492 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

-198-

(ii) MOLECULE TYPE: Genomic DNA
 (iii) HYPOTHETICAL: NO
 (iv) ANTISENSE: NO
 (v) FRAGMENT TYPE:
 (vi) ORIGINAL SOURCE:
 (ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

CTGCAGCTGG	GGGTCTCCAA	TCAGGCAGGG	GCCCCTTACT	ACTCAGATGG	GGTGGCCGAG	60
TAGGGGAAGG	GGGTGCAGGC	TGCATGAGTG	GACACAGCTG	TAGGACTACC	TGGGGGCTGT	120
GGATCTATGG	GGGTGGGGAG	AAGCCCAGTG	ACAGTGCCTA	GAAGAGACAA	GGTGGCCTGA	180
GAGGGTCTGA	GGAACATAGA	GCTGGCCATG	TTGGGGCCAG	GTCTCAAGCA	GGAAGTGAGG	240
AATGGGACAG	GCTTGAGGAT	ACTCTACTCA	GTAGCCAGGA	TAGCAAGGAG	GGCTTGGGGT	300
TGCTATCCTG	GGGTTCACC	CCCCAGGTTG	AAGGCCCTGG	GGGAGATGGT	CCCAGGACAT	360
ATTACAATGG	ACACAGGAGG	TTGGGACACC	TGGAGTCACC	AAACAAAACC	ATGCCAAGAG	420
AGACCATGAG	TAGGGGTGTC	CAGTCCAGCC	CTCTGACTGA	GCTGCATTGT	TCAAATCCAA	480
AGGGCCCCCTG	CTGCCACCTA	GTGGCTGATG	GCATCCACAT	GACCCTGGGC	CACACGCGTT	540
TAGGGTCTCT	GTGAAGACCA	AGATCCTTGT	TACATTGAAC	GACTCCTAAA	TGAGCAGAGA	600
TTTCCACCTA	TTCGAAACAA	TCACATAAAA	TCCATCCTGG	AAAAAGCCTG	GGGGATGGCA	660
CTAAGGCTAG	GGATAGGGTG	GGATGAAGAT	TATAGTTACA	GTAAGGGGTT	TAGGGTTAGG	720
GATCAACGTT	GGTTAGGAGT	TAGGGATACA	GTAGGGTACC	GGTAGGGTTA	GGGGTTAGGG	780
TTAGGGGTGA	GGGTAGGGT	TAGGGTTAGG	GTTAGGGTTA	GGGGTTAGGG	GTTAGGGTTA	840
GGGTAGGTT	TTGGGGTGGC	GTATTTTGGT	CTTATACGCT	GTGTTCCACT	GGCAATGAAA	900
AGAGTTCTTG	TTTTTCCTTC	AGCAATTTGT	CATTTTTTAA	AGAGTTTAGC	AATTCTAACA	960
GATATAGACC	AGCTGTGCTA	TCTCATTGTG	GTTTTCAATT	GTAACCACAT	TGTGGTTTCA	1020
ATGTGTTTAC	TTGCCATCTG	TAGATCTTCT	TTGCGTGAGG	TGTCTGTTCA	GATGTGTGTG	1080
CATTCTTGN	NTTTNGGCTG	TTAACTTAT	TGTTTAGTTT	TAATAATTTT	TTATATATTT	1140
GAAGACAAAT	CTTTCTCAGA	TGTGTATTTG	CAAATATTTT	TTCAATATGA	GGCTTGCTTT	1200
TGTCTCTAAC	AAGGTCTCTT	CAGAGATAAC	TTAAATATAA	GAAATCCACA	CTGTCACTTC	1260
TTTTGTGTAT	ATCTACCTTT	TGTGTCATTT	GTTAAAATTC	ATTACCAAAC	CCAAAGGCAG	1320
ATAGCTTTTC	TTCTATTGTT	TCTTCTAGAA	ATTTGTATAG	TTTTGCATTT	TTAGTGTAAG	1380
GATGATTTTG	AGTGATTATT	TGTGTAAGTT	GTAAAGTTTT	CGTCTATATC	CATATCATTT	1440
CTTATGGTTT	CCAATTAATC	GTTCCCTCAC	TATTTTTGGG	AAAGACACAG	GATAGTGGGC	1500
TTTGTTAGAG	TAGATAGGTA	GCTAGACATG	AACAGGAGGG	GGCCTCCTGG	AAAAGGGAAA	1560
GTCTGGGAAG	GCTCACCTGG	AGGACCACCA	AAAATTCACA	TATTAGTAGC	ATCTCTAGTG	1620
CTGGAGTGGA	TGGGCACTTG	TCAATTGTGG	GTAGGAGGGA	AAAGAGGTCC	TATGCAGAAA	1680
GAAACTCCCT	AGAACTCCTC	TGAAGATGCC	CCAATCATTC	ACTCTGCAAT	AAAAATGTCA	1740
GAATATTGCT	AGCTACATGC	TGATAAGNN	AAAGGGGACA	TTCTTAAGTG	AAACCTGGCA	1800
CCATAAGTAC	AGATTAGGGC	AGAGAAGGAC	ATTCAAAAGA	GGCAGGCGCA	GTAGGTACAA	1860
ACGTGATCGC	TGTCAGTGTG	CCTGGGATGG	CGGGAAGGAG	GCTGGTGCCA	GAGTGGATTC	1920
GTATTGATCA	CCACACATAT	ACCTCAACCA	ACAGTGAGGA	GGTCCCACAA	GCCTAAGTGG	1980
GGCAAGTTGG	GGAGCTAAGG	CAGTAGCAGG	AAAACCAGAC	AAAGAAAACA	GGTGGAGACT	2040
TGAGACAGAG	GCAGGAATGT	GAAGAAATCC	AAAATAAAAT	TCCCTGCACA	GGACTCTTAG	2100
GCTGTTTAAT	GCATCGCTCA	GTCCCACTCC	TCCCTATTTT	TCTACAATAA	ACTCTTTACA	2160
CTGTGTTTCT	TTTCAATGAA	GTTATCTGCC	ATCTTTGTAT	TGCCTCTTGG	TGAAAATGTT	2220
TCTTCCAAGT	TAAACAAGAA	CTGGGACATC	AGCTCTCCCC	AGTAATAGCT	CCGTTTCAGT	2280
TTGAATTTAC	AGAAGTATG	GGCTTAATAA	CTGGCGCTCT	GACTTTAGTG	GTGCAGGAGG	2340
CCGTACACAC	GGGACCAAGA	GTGCCCTGCC	TAGTCCCCAT	CTGCCCGCAG	GTGGCGGCTG	2400
CCTCGACACT	GACAGCAATA	GGGTCCGGCA	TGTCCCCAG	CTGCCAGCAG	GGGGCGTACG	2460
ACGACTACAC	TGTGAGCAAG	AGGGCCCTGC	AG			2492

(2) INFORMATION FOR SEQ ID NO:4:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 28 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

-199-

(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

GGGGAATTCA TTGGGATGTT TCAGTTGA

28

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CGAAAGTCCC CCCTAGGAGA TCTTAAGGA

29

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 47 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO
(v) FRAGMENT TYPE:
(vi) ORIGINAL SOURCE:
(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CCGCTTAATA CTCTGATGAG TCCGTGAGGA CGAAACGCTC TCGCACC

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 25 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA
(iii) HYPOTHETICAL: NO
(iv) ANTISENSE: NO

-200-

- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:
- (ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CGATTTAAAT TAATTAAGCC CGGGC

25

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Genomic DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:
- (ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

TAAATTTAAT TAATTCGGGC CCGTCGA

27

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 69 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Genomic DNA
 - (D) OTHER INFORMATION IL-2 signal sequence

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

ATG TAC AGG ATG CAA CTC CTG TCT TGC ATT GCA CTA AGT CTT GCA CTT
Met Tyr Arg Met Gln Leu Leu Ser Cys Ile Ala Leu Ser Leu Ala Leu

48

GTC ACA AAC AGT GCA CCT ACT
Val Thr Asn Ser Ala Pro Thr

69

(2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 945 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (vi) ORIGINAL SOURCE:
- (ix) FEATURE:

-201-

(A) NAME/KEY: Coding Sequence

(B) LOCATION: 1...942

(D) OTHER INFORMATION: Renilla Reinformis Luciferase

(x) PUBLICATION INFORMATION:

PATENT NO.: 5,418,155

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

AGC	TTA	AAG	ATG	ACT	TCG	AAA	GTT	TAT	GAT	CCA	GAA	CAA	AGG	AAA	CGG	48
Ser	Leu	Lys	Met	Thr	Ser	Lys	Val	Tyr	Asp	Pro	Glu	Gln	Arg	Lys	Arg	
1				5				10						15		
ATG	ATA	ACT	GGT	CCG	CAG	TGG	TGG	GCC	AGA	TGT	AAA	CAA	ATG	AAT	GTT	96
Met	Ile	Thr	Gly	Pro	Gln	Trp	Trp	Ala	Arg	Cys	Lys	Gln	Met	Asn	Val	
			20					25						30		
CTT	GAT	TCA	TTT	ATT	AAT	TAT	TAT	GAT	TCA	GAA	AAA	CAT	GCA	GAA	AAT	144
Leu	Asp	Ser	Phe	Ile	Asn	Tyr	Tyr	Asp	Ser	Glu	Lys	His	Ala	Glu	Asn	
			35				40						45			
GCT	GTT	ATT	TTT	TTA	CAT	GGT	AAC	GCG	GCC	TCT	TCT	TAT	TTA	TGG	CGA	192
Ala	Val	Ile	Phe	Leu	His	Gly	Asn	Ala	Ala	Ser	Ser	Tyr	Leu	Trp	Arg	
			50				55					60				
CAT	GTT	GTG	CCA	CAT	ATT	GAG	CCA	GTA	GCG	CGG	TGT	ATT	ATA	CCA	GAT	240
His	Val	Val	Pro	His	Ile	Glu	Pro	Val	Ala	Arg	Cys	Ile	Ile	Pro	Asp	
65					70					75					80	
CTT	ATT	GGT	ATG	GGC	AAA	TCA	GGC	AAA	TCT	GGT	AAT	GGT	TCT	TAT	AGG	288
Leu	Ile	Gly	Met	Gly	Lys	Ser	Gly	Lys	Ser	Gly	Asn	Gly	Ser	Tyr	Arg	
				85					90					95		
TTA	CTT	GAT	CAT	TAC	AAA	TAT	CTT	ACT	GCA	TGG	TTG	AAC	TTC	TTA	ATT	336
Leu	Leu	Asp	His	Tyr	Lys	Tyr	Leu	Thr	Ala	Trp	Leu	Asn	Phe	Leu	Ile	
			100					105						110		
TAC	CAA	AGA	AGA	TCA	TTT	TTT	GTC	GGC	CAT	GAT	TGG	GGT	GCT	TGT	TTG	384
Tyr	Gln	Arg	Arg	Ser	Phe	Phe	Val	Gly	His	Asp	Trp	Gly	Ala	Cys	Leu	
			115				120						125			
GCA	TTT	CAT	TAT	AGC	TAT	GAG	CAT	CAA	GAT	AAG	ATC	AAA	GCA	ATA	GTT	432
Ala	Phe	His	Tyr	Ser	Tyr	Glu	His	Gln	Asp	Lys	Ile	Lys	Ala	Ile	Val	
			130				135					140				
CAC	GCT	GAA	AGT	GTA	GTA	GAT	GTG	ATT	GAA	TCA	TGG	GAT	GAA	TGG	CCT	480
His	Ala	Glu	Ser	Val	Val	Asp	Val	Ile	Glu	Ser	Trp	Asp	Glu	Trp	Pro	
145						150					155				160	
GAT	ATT	GAA	GAA	GAT	ATT	GCG	TTG	ATC	AAA	TCT	GAA	GAA	GGA	GAA	AAA	528
Asp	Ile	Glu	Glu	Asp	Ile	Ala	Leu	Ile	Lys	Ser	Glu	Glu	Gly	Glu	Lys	
				165					170					175		
ATG	GTT	TTG	GAG	AAT	AAC	TTC	TTC	GTG	GAA	ACC	ATG	TTG	CCA	TCA	AAA	576
Met	Val	Leu	Glu	Asn	Asn	Phe	Phe	Val	Glu	Thr	Met	Leu	Pro	Ser	Lys	
			180					185					190			
ATC	ATG	AGA	AAG	TTA	GAA	CCA	GAA	GAA	TTT	GCA	GCA	TAT	CTT	GAA	CCA	624
Ile	Met	Arg	Lys	Leu	Glu	Pro	Glu	Glu	Phe	Ala	Ala	Tyr	Leu	Glu	Pro	
			195				200						205			

TTC AAA GAG AAA GGT GAA GTT CGT CGT CCA ACA TTA TCA TGG CCT CGT	672
Phe Lys Glu Lys Gly Glu Val Arg Arg Pro Thr Leu Ser Trp Pro Arg	
210 215 220	
GAA ATC CCG TTA GTA AAA GGT GGT AAA CCT GAC GTT GTA CAA ATT GTT	720
Glu Ile Pro Leu Val Lys Gly Gly Lys Pro Asp Val Val Gln Ile Val	
225 230 235 240	
AGG AAT TAT AAT GCT TAT CTA CGT GCA AGT GAT GAT TTA CCA AAA ATG	768
Arg Asn Tyr Asn Ala Tyr Leu Arg Ala Ser Asp Asp Leu Pro Lys Met	
245 250 255	
TTT ATT GAA TCG GAT CCA GGA TTC TTT TCC AAT GCT ATT GTT GAA GGC	816
Phe Ile Glu Ser Asp Pro Gly Phe Phe Ser Asn Ala Ile Val Glu Gly	
260 265 270	
GCC AAG AAG TTT CCT AAT ACT GAA TTT GTC AAA GTA AAA GGT CTT CAT	864
Ala Lys Lys Phe Pro Asn Thr Glu Phe Val Lys Val Lys Gly Leu His	
275 280 285	
TTT TCG CAA GAA GAT GCA CCT GAT GAA ATG GGA AAA TAT ATC AAA TCG	912
Phe Ser Gln Glu Asp Ala Pro Asp Glu Met Gly Lys Tyr Ile Lys Ser	
290 295 300	
TTC GTT GAG CGA GTT CTC AAA AAT GAA CAA TAA	945
Phe Val Glu Arg Val Leu Lys Asn Glu Gln	
305 310	

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

TTTGAATTC A TGTACAGGAT GCAACTCCTG

30

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 30 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(ix) FEATURE:

-203-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

TTTGAATTCA GTAGGTGCAC TGTTTGTAC

30

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1434 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

CCTCCACGCA	CGTTGTGATA	TGTAGATGAT	AATCATTATC	AGAGCAGCGT	TGGGGGATAA	60
TGTCGACATT	TCCACTCCCA	ATGACGGTGA	TGTATAATGC	TCAAGTATTC	TCCTGCTTTT	120
TTACCACTAA	CTAGGAACTG	GGTTTGGCCT	TAATTCAGAC	AGCCTTGGCT	CTGTCTGGAC	180
AGGTCCAGAC	GACTGACACC	ATTAACACTT	TGTCAGCCTC	AGTGA CTACA	GTCATAGATG	240
AACAGGCCTC	AGCTAATGTC	AAGATACAGA	GAGGTCTCAT	GCTGGTTAAT	CAACTCATAG	300
ATCTTGTCCA	GATACA ACTA	GATGTATTAT	GACAAATAAC	TCAGCAGGGA	TGTGAACAAA	360
AGTTTCCGGG	ATTGTGTGTT	ATTTCCATTC	AGTATGTTAA	ATTTACTAGG	ACAGCTAATT	420
TGTCAAAAG	TCTTTTTCAG	TATATGTTAC	AGAATTGGAT	GGCTGAATTT	GAACAGATCC	480
TTCGGGAATT	GAGACTTCAG	GTCAACTCCA	CGCGCTTGGA	CCTGTCGCTG	ACCAAAGGAT	540
TACCCAATTG	GATCTCCTCA	GCATTTTCTT	TCTTTAAAAA	ATGGGTGGGA	TTAATATTAT	600
TTGGAGATAC	ACTTTGCTGT	GGATTAGTGT	TGCTTCTTTG	ATTGGTCTGT	AAGCTTAAGG	660
CCCAAAC TAG	GAGAGACAAG	GTGGTTATTG	CCCAGGCGCT	TGCAGGACTA	GAACATGGAG	720
CTTCCCCTGA	TATATGGTTA	TCTATGCTTA	GGCAATAGGT	CGCTGGCCAC	TCAGCTCTTA	780
TATCCCACGA	GGCTAGTCTC	ATTGTACGGG	ATAGAGTGAG	TGTGCTTCAG	CAGCCCCGAGA	840
GAGTTGCAAG	GCTAAGCACT	GCAATGGAAA	GGCTCTGCGG	CATATATGTG	CCTATTCTAG	900
GGGGACATGT	CATCTTTCAT	GAAGGTTTCAG	TGTCCTAGTT	CCCTTCCCCC	AGGCAAAAACG	960
ACACGGGAGC	AGGTCAGGGT	TGCTCTGGGT	AAAAGCCTGT	GAGCCTGGGA	GCTAATCCTG	1020
TACATGGCTC	CTTTACCTAC	ACACTGGGGA	TTTGACCTCT	ATCTCCACTC	TCATTAATAT	1080
GGGTGGCCTA	TTTGCTCTTA	TTAAAAGGAA	AGGGGGAGAT	GTTGGGAGCC	GCGCCACAT	1140
TCGCCGTTAC	AAGATGGCGC	TGACAGCTGT	GTTCTAAGTG	GTAAACAAAT	AATCTGCGCA	1200
TGTGCCGAGG	GTGGTTCTTC	ACTCCATGTG	CTCTGCCTTC	CCCGTGACGT	CAACTCGGCC	1260
GATGGGCTGC	AGCCAATCAG	GGAGTGACAC	GTCTTAGGCG	AAGGAGAATT	CTCCTTAATA	1320
GGGACGGGGT	TTCGTTCTCT	CTCTCTCTCT	TGCTTCTCTC	TCTTGCTTTT	TCGCTCTCTT	1380
GCTTCCCGTA	AAGTGATAAT	GATTATCATC	TACATATCAC	AACGTGCGTG	GAGG	1434

(2) INFORMATION FOR SEQ ID NO:14:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1400 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

-204-

CCTCCACGCA	CGTTGTGATA	TGTAGATGAT	AATCATTATC	AGAGCAGCGT	TGGGGGATAA	60
TGTCGACATT	TCCACTCCCA	ATGACGGTGA	TGTATAATGC	TCAAGTATTC	TCCTGCTTTT	120
TTACCACTAA	CTAGGAACTG	GGTTTGGCCT	TAATTCAGAC	AGCCTTGGCT	CTGTCTGGAC	180
AGGTCCAGAT	ACAAC TAGAT	GTATTATGAC	AAATAACTCA	GCAGGGATGT	GAACAAAAGT	240
TTCCGGGATT	GCGTGTATT	TCCATCCAGT	ATGTTAAATT	TACTAGGGCA	GCTAATTTGT	300
CAAAAAGTCT	TTTCCAGTAT	ATGTTACAGA	ATTGGATGGC	TGAATTTGAA	CAGATCCTTC	360
GGGAATTGAG	ACTTCAGGTC	AACTCCACGC	GCTTGGACCT	GTCCCTGACC	AAAGGATTAC	420
CCAATTGGAT	CTCCTCAGCA	TTTTCTTTCT	TTAAAAAATG	GGTGGGATTA	ATATTATTTG	480
GAGATACACT	TTGCTGTGGA	TTAGTGTTCG	TTCTTTGATT	GGTCTGTAAG	CTTAAGGCC	540
AAACTAGGAG	AGACAAGGTG	GTTATTGCC	AGGCGCTTGC	AGGACTAGAA	CATGGAGCTT	600
CCCCTGATAT	ATCTATGCTT	AGGCAATAGG	TCGCTGGCCA	CTCAGCTCTT	ATATCCCATG	660
AGGCTAGTCT	CATTGCACGG	GATAGAGTGA	GTGTGCTTCA	GCAGCCCGAG	AGAGTTGCAC	720
GGCTAAGCAC	TGCAATGGAA	AGGCTCTGCG	GCATATATGA	GCCTATTCTA	GGGAGACATG	780
TCATCTTTCA	AGAAGGTTGA	GTGTCCAAGT	GTCTTCTCTC	CAGGCAAAAC	GACACGGGAG	840
CAGGTCAGGG	TTGCTCTGGG	TAAAAGCCTG	TGAGCCTAAG	AGCTAATCCT	GTACATGGCT	900
CCTTTACCTA	CACACTGGGG	ATTTGACCTC	TATCTCCACT	CTCATTAATA	TGGGTGGCCT	960
ATTTGCTCTT	ATTAAAAGGA	AAGGGGGAGA	TGTTGGGAGC	CGCGCCCA	TTCGCCGTTA	1020
CAAGATGGCG	CTGACAGCTG	TGTTCTAAGT	GGTAAACAAA	TAATCTGCGC	ATGCGCCGAG	1080
GGTGGTTCTT	CACTCCATGT	GCTCTGCCTT	CCCCGTGACG	TCAACTCGGC	CGATGGGCTG	1140
CAGTCAATCA	GGGAGTGACA	CGTCCTAGGC	GAAGGAAAAT	TCTCCTTAAT	AGGGACGGGG	1200
TTTCGTTTTT	TCTCTCTCTT	GCTTCGCTCT	CTCTTGCTTC	TTGCTCTCTT	TTCTGAAGA	1260
TGTAAGAATA	AAGCTTTGCC	GCAGAAGATT	CTGGTCTGTG	GTGTTCTTCC	TGGCCGGTGC	1320
TGAGAACGCG	TCTAATAACA	ATTGGTGCCG	AAACCCGGGT	GATAATGATT	ATCATCTACA	1380
TATCAACG	TGCGTGGAGG					1400

(2) INFORMATION FOR SEQ ID NO:15:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 1369 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CCTCCACGCA	CGTTGTGATA	TGTAGATGAT	AATCATTATC	ACTTTACGGG	TCCTTTCACT	60
ACAAC TGCCA	CGAGGCCCGG	TGCTCTGGTA	ATAGATCTTT	GCTGAAAAGG	CACACACATG	120
ACACATTACT	CAAGGTGGGC	TCATCTGAGC	TGCAGATTCA	GCTTAATATG	AATCTTGCCA	180
ATTGTGTGAA	ATCATAAATC	TTCAAAGTGA	CACTCATTGC	CAGACACAGG	TGCCCACCTT	240
TGGCATAATA	AACAAACACA	AATTATCTAT	TATATAAAGG	GTGTTAGAAG	ATGCTTTAGA	300
ATACAAATAA	ATCATGGTAG	ATAACAGTAA	GTTGAGAGCT	TAAATTTAAT	AAAGTGATAT	360
ACCTAATAAA	AATTAAATTA	AGAAGGTGTG	AATATACTAC	AGTAGGTAAG	TTATTTTCATT	420
AATTTATTTT	CTTTCTTAAT	CCTTTATAAT	GTTTTCTGCT	ATTGTCAATT	GCACATCCAT	480
ATGTTCAATT	CTTCACTGTA	ATGAAGAAAT	GTAGTAAATA	TACTTTCCGA	ACAAGTTGTA	540
TCAAATATGT	TACACTTGAT	TCCGTGTGTT	ACTTATCATT	TTATTATTAT	ATTGATTGCA	600
TTCTTTCGTT	ACTTGATATT	ATTACAAGGT	ACATATTTAT	TCTCTCAGAT	CTTCATTATA	660
CTCTAACCAT	TTTATAACAT	ACTTTATTTA	TTCAATTTCTT	ATGTGTGCTG	TGAGGCACAA	720
ATGCCAGAGA	GAAC TTGAGC	AGATAAGAGG	ACAAATTGCA	AGAGTCAGTT	ACCTCCTGCT	780
GTTCCTTGGA	AACTCAGGAT	CAAATTCAGG	TTGTCAGGCT	TGGCAGCATG	CACTTTTTTAC	840
CAGTGCCCTC	ATCTTGCTAG	CCCTGAACAT	CAAGCTTTGC	AGACAGACAG	GCTACACTAA	900
GTGAAC TGGT	CATTACAGC	ATGCATGGTG	ATTTATTGTT	ACTTTCTATT	CCATGCCTTT	960
ACTATTTCTA	CTAGGTGCTA	GCTAGTACTG	TATTTTCGAGA	TAGAAGTTAC	TGAAAGAAAA	1020
TTACATTGTT	TTCTATAGAT	CCTTGATACT	CTTTTCAGCAG	ATATAGAGTT	TTAATCAGGT	1080
CCTAGACCCT	TTCTTCACTC	TTATTAAATA	CTAAGTACAA	ATTAAGTTTA	TCCAAAACAG	1140
TACGGATGTT	GATTTTGTGC	AGTTC TACTA	TGATAATAGT	CTAGCTTCAT	AAATCTGACA	1200

-205-

CACTTATTGG	GAATGTTTTT	GTTAATAAAA	GATTCAGGTG	TACTCTAGG	TCAAGAGAAT	1260
ATTAAACATC	AGTCCCAAAT	TACAACTTC	AATAAAAGAT	TTGACTCTCC	AGTGGTGGCA	1320
ATATAAGTG	ATAATGATTA	TCATCTACAT	ATCACAACGT	GCGTGGAGG		1369

(2) INFORMATION FOR SEQ ID NO:16:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22118 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

GAATCCCCCT	ATCCCTAATC	CAGATTGGTG	GAATAACTTG	GTATAGATGT	TTGTGCATTA	60
AAAACCCTGT	AGGATCTTCA	CTCTAGGTCA	CTGTTTCAGCA	CTGGAACCTG	AATTGTGGCC	120
CTGAGTGATA	GGTCTGGGA	CATATGCAGT	TCTGCACAGA	CAGACAGACA	GACAGACAGA	180
CAGACAGACA	GACAGACGTT	ACAAACAAAC	ACGTTGAGCC	GTGTGCCAAC	ACACACACAA	240
ACACCACTCT	GGCCATAATT	ATTGAGGACG	TTGATTTATT	ATTCTGTGTT	TGTGAGTCTG	300
TCTGTCTGTC	TGTCTGTCTG	TCTGTCTGTC	TATCAAACCA	AAAGAAACCA	AACAATTATG	360
CCTGCCTGCC	TGCCTGCCTG	CCTACACAGA	GAAATGATTT	CTTCAATCAA	TCTAAAACGA	420
CCTCCTAAGT	TTGCCTTTTT	TCTCTTTCTT	TATCTTTTTC	TTTTTTCTTT	TCTTCTTCCT	480
TCCTTCCTTC	CTTCCTTCCT	TCCTTCCTTT	CTTTCTTTCT	TTCTTTCTTT	CTTACTTTCT	540
TTCTTTCTCT	CTTACATTTA	TTCTTTTCAT	ACATAGTTTC	TTAGTGTAAG	CATCCCTGAC	600
TGTCTTGAAG	ACACTTTGTA	GGCCTCAATC	CTGTAAGAGC	CTTCCTCTGC	TTTTCAAATG	660
CTGGCATGAA	TGTTGTACCT	CACTATGACC	AGCTTAGTCT	TCAAGTCTGA	GTTACTGGAA	720
AGGAGTTCCA	AGAAGACTGG	TTATATTTTT	CATTTATTAT	TGCATTTTAA	TTAAATTTTA	780
ATTTACCAAA	AAGAATTTAG	ACTGACCAAT	TCAGAGTCTG	CCGTTTAAAA	GCATAAGGAA	840
AAAGTAGGAG	AAAAACGTGA	GGCTGTCTGT	GGATGGTCGA	GGCTGCTTTA	GGGAGCCTCG	900
TCACCAATCT	GCACTTGCAA	ACCGGGCCAC	TAGAACCCGG	TGAAGGGAGA	AACCAAGCG	960
ACCTGGAAC	AATAGGTCAC	ATGAAGGCCA	GCCACCTCCA	TCTTGTTGTG	CGGGAGTTCA	1020
GTTAGCAGAC	AAGATGGCTG	CCATGCACAT	GTTGTCTTTC	AGCTTGGTGA	GGTCAAAGTA	1080
CAACCGAGTC	ACAGAACAAG	GAAGTATACA	CAGTGAGTTC	CAGGTCAGCC	AGAGTTTACA	1140
CAGAGAAACC	ACATCTTGAA	AAAAACAAAA	AAATAAATTA	AATAAATATA	ATTTAAAAAT	1200
TTAAAAATAG	CCGGGAGTGA	TGGCGCATGT	CTTTAATCCC	AGCTCTCTTC	AGGCAGAGAT	1260
GGGAGGATTT	CTAGTTTGA	GGCCAGCCTG	GTCTGCAAG	TGAGTTCCAG	GACAGTCAGG	1320
GCTATACAGA	GTAACCTGT	CTTGAAACT	AAACTAAAT	AACTAAACT	AAACTAAAAA	1380
AATATAAAAT	AAAAATTTTA	AAGAATTTTA	AAAAACTACA	GAAATCAAAC	ATAAGCCCAC	1440
GAGATGGCAA	GTAAGTCAA	TCATAGCAGA	AATATTATAC	ACACACACAC	ACACAGACTC	1500
TGTCATAAAA	TCCAATGTGC	CTTCATGATG	ATCAAATTTT	GATAGTCAGT	AATACTAGAA	1560
GAATCATATG	TCTGAAAATA	AAAGCCAGAA	CCTTTTCTGC	TTTTGTTTTT	TTTTGCCCCA	1620
AGATAGGGTT	TCTCTCAGT	TATCCCTGGC	ATCCCTGCCT	GGAAGTTTCT	TTGTAGGTTT	1680
GGTAGCCTCA	GAAGTCAAGA	GGTCTCTCT	GCCTGCCTGC	CTGCCTGCCT	GCCTGCCTGC	1740
CTGCCTGCCT	GCCTGCCTCA	CTTCTTCTGC	CACCCACACA	ACCGAGTCGA	ACCTAGGATC	1800
TTTATTTCTT	TCTCTTCTC	TCTTCTTCT	TTCTTCTTT	CTTTCTTCT	TTCTTCTTT	1860
CTTTCTTCT	TTCTTATTCA	ATTAGTTTTT	AATGTAAGTG	TGTGTTTGTG	CTCTATCTGC	1920
TGCCTATAGG	CCTGCTTGCC	AGGAGAGGGG	AACAGAACCT	AGGAGAAACC	ACCATGCAGC	1980
TCCTGAGAA	AAGTGAAAAA	ACAACAAAAA	AAGGAAATTC	TAATCACATA	GAATGTAGAT	2040
ATATGCCGAG	GCTGTCAGAG	TGCTTTTTAA	GGCTTAGTGT	AAGTAATGAA	AATTGTTGTG	2100
TGTCTTTTAT	CCAAACACAG	AAGAGAGGTG	GCTCGGCCTG	CATGTCTGTT	GTCTGCATGT	2160
AGACCAGGCT	GGCCTTGAAC	ACATTAATCT	GTCTGCCTCT	GCTTCCCTAA	TGCTGCGATT	2220
AAAGGCATGT	GCCACCACTG	CCCGGACTGA	TTTCTTCTTT	TTTTTTTTTT	TGGAAAATAC	2280
CTTTCTTCT	TTTCTCTCT	CTCTTTCTTC	CTTCCTTCCT	TTCTTTCTAT	TCTTTTCTTC	2340
TTTCTTTTTT	CTTTTTTTTT	TTTTTTTTTA	AATTTGCCTA	AGGTTAAAGG	TGTGCTCCAC	2400
AATTGCCTCA	GCTCTGCTCT	AATTCTCTTT	AAAAAAAAC	AAACAAAAAA	AAAACCAAAA	2460

-206-

CAGTATGTAT	GTATGTATAT	TTAGAAGAAA	TACTAATCCA	TTAATAACTC	TTTTTTCCTA	2520
AAATTCATGT	CATTCTTGTT	CCACAAAGTG	AGTTCCAGGA	CTTACCAGAG	AAACCCTGTG	2580
TTCAAATTTT	TGTGTTCAAG	GTCAACCCTGG	CTTACAAAGT	GAGTTCCAAG	TCCGATAGGG	2640
CTACACAGAA	AAACCATATC	TCAGAAAAAA	AAAAAGTTCC	AAACACACAC	ACACACACAC	2700
ACACACACAC	ACACACACAC	ACACACACAC	ACACACACAG	CGCGCCGCGG	CGATGAGGGG	2760
AAGTCGTGCC	TAAAATAAAT	ATTTTTCTGG	CCAAAGTGAA	AGCAAATCAC	TATGAAGAGG	2820
TACTCCTAGA	AAAAATAAAT	ACAAACGGGC	TTTTTAATCA	TTCCAGCACT	GTTTTAATTT	2880
AACCTCTGAAT	TTAGTCTTGG	AAAAGGGGGC	GGGTGTGGGT	GAGTGAGGGC	GAGCGAGCAG	2940
ACGGGCGGGC	GGGCGGGTGA	GTGGCCGGCG	GCGGTGGCAG	CGAGCACCAG	AAAACAACAA	3000
ACCCCAAGCG	GTAGAGTGTT	TTAAAAATGA	GACCTAAATG	TGGTGGAACG	GAGGTCGCCG	3060
CCACCCTCCT	CTTCCACTGC	TTAGATGCTC	CCTTCCCCTT	ACTGTGCTCC	CTTCCCCTAA	3120
CTGTGCCTAA	CTGTGCCTGT	TCCCTCACCC	CGCTGATTCC	CCAGCGACGT	ACTTTGACTT	3180
CAAGAACGAT	TTTGCTGTG	TTCCACCGCTC	CCTGTCTATC	TTTCGTTTTT	GGGTGCCCGA	3240
GTCTAGCCCC	TTTCGTATGT	TCGGGCGGGA	CGATGGGGAC	CGTTTGTGCC	ACTCGGGAGA	3300
AGTGGTGGGT	GGGTACGCTG	CTCCGTCGTG	CGTGCGTGAG	TGCCGGAACC	TGAGCTCGGG	3360
AGACCCTCCG	GAGAGACAGA	ATGAGTGAGT	GAATGTGGCG	GCGCGTGACG	GATCTGTATT	3420
GGTTTGTATG	GTTGATCGAG	ACCATTGTCT	GGCGACACCT	AGTGGTGACA	AGTTTCGGGA	3480
ACGCTCCAGG	CCTCTCAGGT	TGGTGACACA	GGAGAGGGAA	GTGCCTGTGG	TGAGGCGACC	3540
AGGGTGACAG	GAGGCCGGGC	AAGCAGGGCG	GAGCGTCTCG	GAGATGGTGT	CGTGTTTAAG	3600
GACGCTCTCT	AACAAGGAGG	TCGTACAGGG	AGATGGCCAA	AGCAGACCGA	GTTGCTGTAC	3660
GCCCTTTTGG	GAATAATGCT	AGGGTTGGTG	GCAACGTTAC	TAGGTCGACC	AGAAGGCTTA	3720
AGTCCTACCC	CCCCCCCCCT	TTTTTTTTTT	TTTCCTCCAG	AAGCCCTCTC	TTGTCCCCGT	3780
CACCGGGGGC	ACCGTACATC	TGAGGCCGAG	AGGACGCGAT	GGGCCCCGCT	TCCAAGCCGG	3840
TGTGGCTCGG	CCAGCTGGCG	CTTCGGGTCT	TTTTTTTTTT	TTTTTTTTTT	TTTTCTCCA	3900
GAAGCCTTGT	CTGTGCTGTG	CACCGGGGGC	GCTGTACTTC	TGAGGCCGAG	AGGACGCGAT	3960
GGGCCCGGGC	TTCCAAGCCG	GTGTGGCTCG	GCCAGCTGGA	GCTTCGGGTC	TTTTTTTTTT	4020
TTTTTTTTTT	TTTTTTTCTC	CAGAAGCCTT	GTCTGTGCTG	GTCACCGGGG	GCGCTGTACT	4080
TCTGAGGCCG	AGAGGACGCG	ATGGGTGCGC	TTCCAAGCCG	ATGTGGCGGG	GCCAGCTGGA	4140
GCTTCGGGTT	TTTTTTTTTT	CTCCAGAAGC	CCTCTCTTGT	CCCCGTCACC	GGGGGCGCTG	4200
TACTTCTGAG	GCCGAGAGGA	CGTGATGGGC	CCGGGTTCCA	GGCGGATGTC	GCCCGGTGAG	4260
CTGGAGCTTT	GGATCTTTTT	TTTTTTTTTT	CCTCCAGAAG	CCCTCTCTTG	TCCCCGTCAC	4320
CGGGGGCACC	TTACATCTGA	GGGCGAGAGG	ACGTGATGGG	TCCGGCTTCC	AAGCCGATGT	4380
GGCGGGGCCA	GCTGGAGCTT	CGGGTTTTTT	TTTTTTCCTC	CAGAAGCCCT	CTCTTGTCCT	4440
CGTCAACGGG	GGCGCTGTAC	TTCTGAGGCC	GAGAGGACGT	GATGGGCCCC	GGTTCAGGC	4500
GGATGTCGCC	CGGTGAGCTG	GAGCTTTGGA	TCATTTTTTT	TTTCCCTCC	AGAAGCCCTC	4560
TCTTGTCCCC	GTCACCGGGG	GCACCGTACA	TCTGAGGCCG	AGAGGACACG	ATGGGCCTGT	4620
CTTCCAAGCC	GATGTGGCCC	GGCCAGCTGG	AGCTTCGGGT	CTTTTTTTTT	TTTTTTCCTC	4680
CAGAAGCCTT	GTCGTGCGCT	GTCACCGGGG	GCGCTGTACT	TCTGAGGCCG	AGAGGACGCG	4740
ATGGGCCCGG	CTTCCAAGCC	GGTGTGGCTC	GGCCAGCTGG	AGCTTCGGGT	CTTTTTTTTT	4800
TTTTTTTTTT	TTCTCCAGA	AACCTTGTCT	GTCGCTGTCA	CCCGGGGCGC	TTGTACTTCT	4860
GATGCCGAGA	GGACGCGATG	GGCCCGTCTT	CCAGGCCGAT	GTGGCCCGGT	CAGCTGGAGC	4920
TTTGGATCTT	TTTTTTTTTT	TTTCCCTCCA	GAAGCCCTCT	CTTGTCCTCG	TCACCGGGGG	4980
CACCTTACAT	CTGAGGCCTA	GAGGACACGA	TGGGCCCGGG	TTCCAGGCCG	ATGTGGCCCC	5040
GTCAGCTGGA	GCTTTGGATC	TTTTTTTTTT	TTTTCTTCCA	GAAGCCCTCT	TGTCCCCGTC	5100
ACCGGTGGCA	CTGTACATCT	GAGGCGGAGA	GGACATTATG	GGCCCGGCTT	CCAATCCGAT	5160
GTGGCCCGGT	CAGCTGGAGC	TTTGGATCTT	ATTTTTTTTT	TAATTTTTTT	TTCCAGAAGC	5220
CCTCTTGTC	CTGTCAACGG	TGGCACGGTA	CATCTGAGGC	CGAGAGGACA	TTATGGGCCC	5280
GGCTTCCAGG	CCGATGTGGC	CCGGTCAGCT	GGAGCTTTGG	ATCTTTTTTT	TTTTTTTTTCT	5340
TTTTTCTCC	AGAAGCCCTC	TCTGTCCCTG	TCACCGGGGG	CCCTGTACGT	CTGAGGCCGA	5400
GGGAAAGCTA	TGGGCGCGGT	TTTCTTTTCT	TGACCTGTCT	GTCTTATCAG	TTCTCCGGGT	5460
TGTCAGGGTC	GACCAGTTGT	TCCTTTGAGG	TCCGGTTCTT	TTCTGTTATG	GGTCATTTTT	5520
GGGCCACCTC	CCCAGGTATG	ACTTCCAGGC	GTCGTTGCTC	GCCTGTCACT	TTCTTCCCTG	5580
TCTCTTTTAT	GCTTGTGATC	TTTTCTATCT	GTTCTTATTT	GACCTGGAGA	TAGGTACTGA	5640
CACGCTGTCC	TTTCCCTATT	AACACTAAAG	GACACTATAA	AGAGACCCCT	TCGATTTAAG	5700
GCTGTTTTGC	TTGTCCAGCC	TATTTCTTTT	ACTGGCTTGG	GTCTGTGCGG	GTGCCTGAAG	5760
CTGTCCCCGA	GCCACGCTTC	CTGCTTTCCC	GGGCTTGTCT	CTTGCGTGTG	CTTGCTGTGG	5820
GCAGCTTGTC	ACAACCTGGG	GCTGTGACTT	TGCTGCGTGT	CAGACGTTTT	TCCCAGATTC	5880
CCCGAGGTGT	CGTTGTACAC	CCTGTCCCGG	TTGGAATGGT	GGAGCCAGCT	GTGGTTGAGG	5940
GCCACCTTAT	TTCCGGCTCAC	TTTTTTTTTT	TTTTTTTTCT	TTGGAGTCCC	GAACCTCCGC	6000
TCTTTTCTCT	TCCCGGTCTT	TCTTCCACAT	GCCTCCCGAG	TGCATTTCTT	TTTGTTTTTT	6060
TTCTTTTTTT	TTTTTTTTTT	TTGGGGAGGT	GGAGAGTCCC	GAGTACTTCA	CTCCTGTCTG	6120

-207-

TGGTGTCCAA	GTGTTTCATGC	CACGTGCCTC	CCGAGTGCAC	TTTTTTTTTGT	GGCAGTCGCT	6180
CGTTGTGTTT	TCTTGTCTCT	TGTCTGCCCG	TATCAGTAAC	TGTCTTGCCC	CGCGTGTAAG	6240
ACATTCCCTAT	CTCGCTTGTT	TCTCCCGATT	GCGCGTCGTT	GCTCACTCTT	AGATCGATGT	6300
GGTGCTCCGG	AGTTCTCTTC	GGGCCAGGGC	CAAGCCGCGC	CAGGCGAGGG	ACGGACATTC	6360
ATGGCGAATG	GCGGCCGCTC	TTCTCGTTCT	GCCAGCGGGC	CCTCGTCTCT	CCACCCCATC	6420
CGTCTGCCGG	TGGTGTGTGG	AAGGCAGGGG	TGCGGCTCTC	CGGCCCGACG	CTGCCCCGCG	6480
CGCACTTTTC	TCAGTGGTTC	GCGTGGTCCT	TGTGGATGTG	TGAGGCGCCC	GGTTGTGCCC	6540
TCACGTGTTT	CACTTTGGTC	GTGTCTCGCT	TGACCATGTT	CCCAGAGTCG	GTGGATGTGG	6600
CCGGTGGCGT	TGCATACCTT	TCCCGTCTGG	TGTGTGCACG	CGCTGTTTCT	TGTAAGCGTC	6660
GAGGTGCTCC	TGGAGCGTTC	CAGGTTTGTC	TCCTAGGTGC	CTGCTTCTGA	GCTGGTGGTG	6720
GCGCTCCCCA	TTCCCTGGTG	TGCCCTCCGT	GCTCCGTCGT	GCTGTGTGCC	TTCCCGTTTG	6780
TGTCTGAGAA	GCCCGTGAGA	GGGGGGTCTG	GGAGAGAAGG	AGGGGCAAGA	CCCCCTTCT	6840
TCGTCCGGTG	AGGCGCCAC	CCCGCGACTA	GTACGCCTGT	GCGTAGGGCT	GGTGCTGAGC	6900
GGTCGCGGCT	GGGGTTGGAA	AGTTTCTCGA	GAGAQTCAAT	GCTTTCCCGT	GGGGAGCTTT	6960
GAGAGGCCTG	GCTTTCCGGG	GGGACCGGTT	GCAGGGTCTC	CCCTGTCCGC	GGATGCTCAG	7020
AATGCCCTTG	GAAGAGAACC	TTCTGTGTC	CGCAGACCCG	CCCGCGCGGT	CGCCCGCGTG	7080
TTGGTCTTCT	GGTTTCCCTG	TGTGCTCGTC	GCATGCATCC	TCTCTCGGTG	GCCGCGGCTC	7140
GTCCGGGTTT	TGGGTCCGTC	CCGCCCTCAG	TGAGAAAGTT	TCCTTCTCTA	GCTATCTTCC	7200
GGAAAGGGTG	CGGGCTTCTT	ACGGTCTCGA	GGGGTCTCTC	CCGAATGGTC	CCCTGGAGGG	7260
CTCGCCCCCT	GACCGCCTCC	CGCGCGCGCA	GCCTTTGCTC	TCTCGTCTAC	CGCGGCCCGC	7320
GGCCTCCCCG	CTCCGAGTTC	GGGGAGGGAT	CACGCGGGGC	AGAGCCTGTC	TGTCGTCTCT	7380
CCGTTGCTGC	GGAGCATGTG	GCTCGGCTTG	TGTGGTTGGT	GGCTGGGGAG	AGGGCTCCGT	7440
GCACACCCCG	GCGTGCGCGT	ACTTTCCTCC	CCTCTGAGG	GCCGCCGTGC	GGACGGGGTG	7500
TGGGTAGGCG	ACGGTGGGCT	CCCGGGTCCC	CACCCGTCTT	CCCGTGCCCT	ACCCGTGCCT	7560
TCCGTCCGCT	GCGTCCCTCT	CGCTCGCGTC	CACGACTTTG	GCCGCTCCCG	CGACGGCGGC	7620
CTGCGCCGCG	CGTGGTGCGT	GCTGTGTGCT	TCTCGGGCTG	TGTGGTTGTG	TCGCCTCGCC	7680
CCCCCTTCTC	CGCGGCAGCG	TTCCACCGGC	TGGCGAAATC	GCGGGAGTCC	TCCTTCCCTC	7740
CCTCGGGGTC	GAGAGGGTCC	GTGTCTGGCG	TTGATTGATC	TGCTCTCGG	GGACGGGACC	7800
GTTCTGTGGG	AGAACGGCTG	TTGGCCGCGT	CCGGCGCGAC	GTCGGACGTG	GGGACCCACT	7860
GCCGCTCGGG	GCTCTTCGTC	GGTAGGCATC	GGTGTGTGCG	CATCGGTCTC	TCTCTCGTGT	7920
CGGTGTGCGC	TCCTCGGGCT	CCCGGGGGGC	CGTCGTGTTT	CGGGTCGGCT	CGGCGCTGCA	7980
GGTGTGGTGG	GAATGCTCAG	GGGAGTGGTG	CAGTGTGATT	CCCGCCGGTT	TTGCCTCGCG	8040
TGCCCTGACC	GGTCCGACGC	CCGAGCGGTC	TCTCGGTCCC	TTGTGAGGAC	CCCCTCCCG	8100
GAGGGGCCCC	TTTCGGCCGC	CCTTGCCGTC	TTCCGCGGCT	CTCGTTCTGC	TGTGTCTGTC	8160
CCCCCTCCCC	GTCGTCGCGA	GCCGGTCTTT	GTTCTCTCT	CCCCCTCTCT	CCTCTGACTG	8220
ACCCGTGGCC	GTGCTGTGCG	ACCCCCCGCA	TGGGGGCGGC	CGGGCACGTA	CGCGTCCGGG	8280
CGGTCAACCG	GGTCTTGGGG	GGGGGCCGAG	GGGTAAGAAA	GTGCGCTCGG	CGGGCGGGAG	8340
GAGCTGTGGT	TTGGAGGGCG	TCCCGGCCCC	GCGGCCGTGG	CGGTGTCTTG	CGCGGTCTTG	8400
GAGAGGGCTG	CGTGCGAGGG	GAAAAGGTTG	CCCGCGGAGG	GCAAAGGGAA	AGAGGCTAGC	8460
AGTGGTCAAT	GTCCCGACGG	TGTGGTGGTG	TGTTGGCCGA	GGTGCGTCTG	GGGGGCTCGT	8520
CCGCGCTTGT	CGTCCGTCGG	GAAGCGCGCT	GTTGGGCGCT	GCCGGAGTGC	CGAGGTGGGT	8580
ACCCTGGCGG	TGGGATTAAC	CCCGCGCGCG	TGTCCCGGTG	TGGCGGTGGG	GGCTCCGGTC	8640
GATGTCTACC	TCCCTCTCCC	CGAGGTCTCA	GGCCTTCTCC	GCGCGGGCTC	TCGGCCCTCC	8700
CCTCGTTTCT	CCCTCTCGCG	GGGTCAAGT	CGCTCGTCGA	CCTCCCCCTC	TCCGTCTTTC	8760
CATCTCTCGC	GCAATGGCGC	CGCCCGAGTT	CACGGTGGGT	TCGTCTCTCC	CCTCCGCTTC	8820
TCGCGGGGGG	CTGGCCGCTG	TCCGGTCTCT	CCTGCCCGAC	CCCCGTTGGC	GTGGTCTTCT	8880
CTCGCCGGCT	TCGCGGACTC	CTGGCTTCCG	CCGGAGGGTC	AGGGGGCTTC	CCGGTTCCCC	8940
GACGTTGCGC	CTCGCTGCTG	TGTGCTTGGG	GGGGGCCCGC	TGCGGCCTCC	GCCCGCCCGT	9000
GAGCCCCCTG	CGCACCCGCC	GGTGTGCGGT	TTGCGCGCGC	GGTCAGTTGG	GCCCTGGCGT	9060
TGTGTGCGGT	CGGGAGCGTG	TCCGCCCTCG	GGCGGCTAGA	CGCGGGTGTC	GCCGGGCTCC	9120
GACGGGTGGC	CTATCCAGGG	CTCGCCCCCG	CCGACCCCCG	CCTGCCCGTC	CCGGTGGTGG	9180
TCGTTGGTGT	GGGGAGTGAA	TGGTGCTACC	GGTCATTCCC	TCCCGCGTGG	TTTGACTGTC	9240
TCGCGCGTGT	CGCGCTTCTC	TTTCCGCAAA	CCCCACGCC	AACCCACCAC	CCTGCTCTCC	9300
CGGCCCGGTG	CGGTTCGACG	TCCGGCTCTC	CCGATGCCGA	GGGGTTCGGG	ATTTGTGCCG	9360
GGGACGGAGG	GGAGAGCGGG	TAAGAGAGGT	GTCGGAGAGC	TGTCCCGGGG	CGACGCTCGG	9420
GTTGGCTTTG	CCGCGTGCGT	GTGCTCGCGG	ACGGGTTTTG	TCGGACCCCG	ACGGGGTTCG	9480
TCCGGCCGCA	TGCACTCTCC	CGTTCCGCGC	GAGCGCCCCG	CCGGCTCACC	CCCGGTTTGT	9540
CCTCCGCGCA	GGCTCTCCGC	CGCCGCCGCC	TCCTCTCTCT	CTCTCGCGCT	CTCTGTCCCG	9600
CCTGGCTCTG	TCCCACCCCG	GACGCTCCCG	TCGCGCTTCC	TTACCTGGTT	GATCTGTCCA	9660
GGTAGCATAT	GCTTGTCTCA	AAGATTAAGC	CATGCATGTC	TAAGTACGCA	CGGCCGGTAC	9720
AGTGAAACTG	CGAATGGCTC	ATTAAATCAG	TTATGGTTCC	TTTGGTCTCT	CGCTCCTCTC	9780

CTACTTGGAT	AACTGTGGTA	ATTCTAGAGC	TAATACATGC	CGACGGGCGC	TGACCCCCCT	9840
TCCCGGGGGG	GGATGCGTGC	ATTTATCAGA	TCAAACCAA	CCCGGTGAGC	TCCCTCCCGG	9900
CTCCGGCCGG	GGGTGCGGCG	CCGGCGGCTT	GGTGACTCTA	GATAACCTCG	GGCCGATCGC	9960
ACGCCCCCGG	TGGCGGCGAC	GACCCATTCC	AACGTCTGCC	CTATCAACTT	TCGATGGTAG	10020
TCGCCGTGCC	TACCATGGTG	ACCACGGGTG	ACGGGGAATC	AGGGTTTCGAT	TCCGGAGAGG	10080
GAGCCTGAGA	AACGGCTACC	ACATCCAAGG	AAGGCAGCAG	GCGCGCAAAT	TACCCACTCC	10140
CGACCCGGGG	AGGTAGTGAC	GAAAAATAAC	AATACAGGAC	TCTTTTCGAGG	CCCTGTAATT	10200
GGAATGAGTC	CACTTTAAAT	CCTTTAACGA	GGATCCATTG	GAGGGCAAGT	CTGGTGCCAG	10260
CAGCCGCGGT	AATTCAGCT	CCAATAGCGT	ATATTAAAGT	TGCTGCAGTT	AAAAAGCTCG	10320
TAGTTGGATC	TTGGGAGCGG	GCGGGCGGTC	CGCCGCGAGG	CGAGTCACCG	CCCGTCCCCG	10380
CCCCTTGCCT	CTCGGCGCCC	CCTCGATGCT	CTTAGCTGAG	TGTCCCGCGG	GGCCCGAAGC	10440
GTTTACTTTG	AAAAAATTAG	AGTGTTCAAA	GCAGGCCCGA	GCCGCCTGGA	TACCGCAGCT	10500
AGGAATAATG	GAATAGGACC	GCGGTTCTAT	TTTGTGGTTT	TTCGGAACTG	AGGCCATGAT	10560
TAAGAGGGAG	GGCCGGGGGC	ATTTCGTATT	CGCCGCTAGA	GGTGAAATTC	TTGGACCGGC	10620
GCAAGACGGA	CCAGAGCGAA	AGCATTTGCC	AAGAATGTTT	TCATTAATCA	AGAACGAAAG	10680
TCGGAGGTTT	GAAGACGATC	AGATACCGTC	GTAGTTCCGA	CCATAAACGA	TGCCGACTGG	10740
CGATGCGGCG	GCGTTATTCC	CATGACCCGC	CGGGCAGCTT	CCGGGAAACC	AAAGTCTTTG	10800
GGTTCCGGGG	GGAGTATGGT	TGCAAAGCTG	AACTTAAAG	GAATTGACGG	AAGGGCACCA	10860
CCAGGAGTGG	GCCTGCGGCT	TAATTTGACT	CAACACGGGA	AACCTCACCC	GGCCCGGACA	10920
CGGACAGGAT	TGACAGATTG	ATAGCTCTTT	CTCGATTCCG	TGGGTGGTGG	TGCTATGGCCG	10980
TTCTTAGTTG	TGGGAGCGAT	TTGTCTGGTT	AATTCGATA	ACGAACGAGA	CTCTGGCATG	11040
CTAACTAGTT	ACGCGACCCC	CGAGCGGTCG	GCGTCCCCCA	ACTTCTTAGA	GGGACAAGTG	11100
GCGTTCAGCC	ACCCGAGATT	GAGCAATAAC	AGGTCTGTGA	TGCCCTTAGA	TGTCCGGGGC	11160
TGCACGCGCG	CTACACTGAC	TGGCTCAGCG	TGTGCCTACC	CTGCGCCGGC	AGGCGCGGGT	11220
AAACCGTTGA	ACCCCATTCG	TGATGGGGAT	CGGGGATTGC	AATTATTCCC	CATGAACGAG	11280
GAATTCCCG	TAAGTGCGGG	TCATAAGCTT	CGTTTGATTA	AGTCCCTGCC	CTTTGTACAC	11340
ACCGCCCGTC	GCTACTACCG	ATTGGATGGT	TTAGTGAGGC	CCTCGGATCG	GCCCCCGCGG	11400
GGTCGGCCCA	CGGCCCTGGC	GGAGCGCTGA	GAAGACGGTC	GAACCTGACT	ATCTAGAGGA	11460
AGTAAAAGTC	GTAACAAGGT	TTCCGTAGGT	GAACCTGCGG	AAGGATCATT	AAACGGGAGA	11520
CTGTGGAGGA	GCGGCGGCGT	GGCCCGCTCT	CCCCGTCTTG	TGTGTGTCTT	CGCCGGGAGG	11580
CGCGTGCGTC	CCGGGTCCCC	TGCCCCGCGT	GTGGAGCGAG	GTGTCTGGAG	TGAGGTGAGA	11640
GAAGGGGTGG	TCTGGGTCCG	TCTGGGTCCG	CCTGAGGACTA	CCTCCGATTT	CCCCCGCCCC	11700
TCCCTCTCTC	CTCGTCCGGC	TCTGACCTCG	CCACCTTACC	GCGGCGGCGG	CTGCTCGCGG	11760
GCGTCTTGCC	TCTTTCCCGT	CCGGCTCTTC	CGTGTCTACG	AGGGGCGGTA	CGTCTTACG	11820
GGTTTTGTAC	CCGTCCCGGG	GGCGTTCGGT	CGTCGGGGCG	CGCGCTTTGC	TCTCCCGGCA	11880
CCCATCCCCG	CCGCGGCTCT	GGCTTTTCTA	CGTTGGCTGG	GGCGGTTGTC	GCGTGTGGGG	11940
GGATGTGAGT	GTCCGCTGTG	GGCTCGCCCG	TCCCGATGCC	ACGCTTTTCT	GGCCTCGCGT	12000
GTCTCTCCCG	CTCCTGTCCC	GGGTACCTAG	CTGTCCGCTT	CCGGCGCGGA	GGTTTAAGGA	12060
CCCCGGGGGG	GTCGCCCTGC	CGCCCCCAGG	GTCGGGGGGC	GGTGGEGCCC	GTAGGGAAGT	12120
CGGTCTGTCG	GGCGGCTCTC	CCTCAGACTC	CATGACCCTC	CTCCCCCGGC	TGCCGCGGTT	12180
CCCGAGGCGG	CGGTCTGTGT	GGGGGGTGGA	TGTCTGGAGC	CCCCTCGGGC	GCCGTGGGGG	12240
CCCGACCCGC	GCCGCCGGCT	TGCCCGATTT	CCGCGGGTCG	GTCCTGTCTG	TGCCGGTCTG	12300
GGGTTCCCGT	GTCGTTCCCC	TGTTTTTCCG	CTCCCGACCC	TTTTTTTTTT	CTCCCCCCCC	12360
CACGTGTCTC	GTTTCGTTCC	TGCTGGCCCG	CCTGAGGCTA	CCCCTCGGTC	CATCTGTTCT	12420
CCTCTCTCTC	CGGGGAGAGG	AGGGCGGTGG	TCGTTGGGGG	ACTGTGCCGT	CGTCAGCACC	12480
CGTGAGTTCG	CTCACACCCG	AAATACCGAT	ACGACTCTTA	GCGGTGGATC	ACTCGGCTCG	12540
TGCGTCGATG	AAGAACGCAG	CTAGCTGCGA	GAATTAATGT	GAATTGCAGG	ACACATTGAT	12600
CATCGACACT	TCGAACGCAC	TTGCGGCCCC	GGGTTCTCTC	CGGGGCTACG	CCTGTCTGAG	12660
CGTCGGTTGA	CGATCAATCG	CGTCACCCGC	TGCGGTGGTG	GCTGCGCGGC	TGGGAGTTTG	12720
CTCGCAGGGC	CAACCCCCCA	ACCCGGGTCG	GGCCCTCCGT	CTCCCGAAGT	TCAGACGTGT	12780
GGGCGGTTGT	CGGTGTGGCG	CGCGCGCCCG	CGTCGCGGAG	CCTGGTCTCC	CCCGCGCATC	12840
CGCGCTCGCG	GCTTCTTCCC	GCTCCGCCGT	TCCCGCCCTC	GCCCGTGCAC	CCCGGTCTTG	12900
GCCTCGCGTC	GGCGCCTCCC	GGACCGCTGC	CTCACCAGTC	TTTCTCGGTC	CCGTGCCCCG	12960
TGGGAACCCA	CCGCGCCCCC	GTGGCGCCCC	GGGGTGGGCG	CGTCCGCATC	TGCTCTGGTC	13020
GAGGTGGCGG	GTTGAGGGTG	TGCGTGCGCC	GAGGTGGTGG	TGCGTCCCTT	GCGGCCGCGG	13080
GGTTGTGCGG	TGGCGGGTCG	ACGAGGGCCG	GTCGGTCCGC	TGCGGTGGTT	GTCTGTGTGT	13140
GTTTGGGTCT	TGCGCTGGGG	GAGGCGGGGT	CGACCGCTCG	CGGGGTGGGC	GCGGTGCCCC	13200
GGCGCCGCGC	ACCCTCCGGC	TTGTGTGGAG	GGAGAGCGAG	GGCGAGAACG	GAGAGAGGTG	13260
GTATCCCCGG	TGGCGTTGCG	AGGGAGGGTT	TGGCGTCCCG	CGTCCGTCCG	TCCCTCCCTC	13320
CCTCGGTGGG	CGCCTTCGCG	CCGCACGCGG	CCGCTAGGGG	CGGTGCGGGC	CCGTGGCCCC	13380
CGTGGCTCTT	CTTCGTCTCC	GCTTCTCTCT	CACCCGGGCG	GTACCCGCTC	CGGCGCCGGC	13440

-209-

CCGCGGGACG	CCGCGGCGTC	CGTGCGCCGA	TGCGAGTCAC	CCCCGGGTGT	TGCGAGTTCC	13500
GGGAGGGAGA	GGGCCTCGCT	GACCCGTTGC	GTCCCGGCTT	CCCTGGGGGG	GACCCGGCGT	13560
CTGTGGGCTG	TGCGTCCCGG	GGGTTGCGTG	TGAGTAAGAT	CCTCCACCCC	CGCCGCCCTC	13620
CCCTCCCGCC	GGCCTCTCGG	GGACCCCTTG	AGACGGTTCC	CCGGCTCGTC	CTCCCGTGCC	13680
GCCGGGTGCC	GTCTCTTTCC	CGCCCGCCTC	CTCGCTCTCT	TCTTCCCGCG	GCTGGGCGCG	13740
TGTCCCCCTT	TTCTGACCGC	GACCTCAGAT	CAGACGTGGC	GACCCGCTGA	ATTTAAGCAT	13800
ATTAGTCAGC	GGAGGAAAAG	AAACTAACCA	GGATTCCCTC	AGTAACGGCG	AGTGAACAGG	13860
GAAGAGCCCA	GCGCCGAATC	CCCGCCGCGC	GTCCGCGCGT	GGGAAATGTG	GCGTACGGAA	13920
GACCCACTCC	CCGGCGCCGC	TCGTGGGGGG	CCCAAGTCCT	TCTGATCGAG	GCCCAGCCCC	13980
TGGACGGTGT	GAGGCCGGTA	GCGGCCCCGG	CGCGCCGGGC	TCGGGTCTTC	CCGGAGTCGG	14040
GTTGCTTGGG	AATGCAGCCC	AAAGCGGGTG	GTAAACTCCA	TCTAAGGCTA	AATACCGGCA	14100
CGAGACCGAT	AGTCAACAAG	TACCGTAAGG	GAAAGTTGAA	AAGAACTTTG	AAGAGAGAGT	14160
TCAAGAGGGC	GTGAAACCGT	TAAGAGSTAA	ACGGGTGGGG	TCCGCGCAGT	CCGCCCGGAG	14220
GATTCAACCC	GTGGCGCGCG	GTCCGGCCGT	GCCCGTGGT	CCCGGCGGAT	CTTTCCCGCT	14280
CCCCGTTCTT	CCCGACCCCT	CCACCCGCGC	GTCGTTCCCC	TCTTCTCTCC	CGCGTCCGGC	14340
GCCTCCGGCG	GCGGGCGCGG	GGGTGGTGT	GGTGGTGGCG	CGCGGGCGGG	GCCGGGGGTG	14400
GGGTCCGGCG	GGGACCGCCC	CCGGCCGGCG	ACCGGCCGCC	GCCGGGCGCA	CTTCCACCGT	14460
GGCGGTGCGC	CGCGACCGGC	TCCGGGACGG	CCGGGAAGGC	CCGGTGGGGA	AGGTGGCTCG	14520
GGGGGGCGCG	CGCGTCTCAG	GGCGCGCCGA	ACCACCTCAC	CCCGAGTGTT	ACAGCCCTCC	14580
GGCCGCGCTT	TCGCCGAATC	CCGGGGCCGA	GGAAGCCAGA	TACCCGTGCG	CGCGCTCTCC	14640
CTCTCCCCCC	GTCCGCTCTC	CGGGCGGGCG	TGGGGGTGGG	GGCCGGGCGG	CCCCCTCCAC	14700
GGCGCGACCG	CTCTCCCACC	CCCCTCCGTC	GCCTCTCTCG	GGGCCCGGTG	GGGGGCGGGG	14760
CGGACTGTCC	CCAGTGCGCC	CCGGGCGTCC	TCGCGCCGTC	GGGTCCCGGG	GGGACCGTCC	14820
GTCACGCGTC	TCCCAGCGAA	GCCGAGCGCA	CGGGGTCCGG	GGCGATGTCC	GCTACCCACC	14880
CGACCCGTCT	TGAAACACGG	ACCAAGGAGT	CTAACGCGTG	CGCGAGTCAG	GGGCTCGTCC	14940
GAAAGCCGCC	GTGGCGCAAT	GAAGGTGAAG	GGCCCCCGCC	GGGGGCCCCA	GGTGGGATCC	15000
CGAGGCCTCT	CCAGTCCGCC	GAGGGCGCAC	CACCGGCCCG	TCTCGCCCCG	CGCGCCGGGG	15060
AGGTGGAGCA	CGAGCGTACG	CGTTAGGACC	CGAAAGATGG	TGAACTATGC	TTGGGCAGGG	15120
CGAAGCCAGA	GGAAACTCTG	GTGGAGGTCC	GTAGCGGTCC	TGACGTGCAA	ATCGGTCTGC	15180
CGACCTGGGT	ATAGGGGCGA	AAGACTAATC	GAACCATCTA	GTAGCTGGTT	CCCTCCGAAG	15240
TTTCTCTCAG	GATAGCTGGC	GCTCTCGCTC	CCGACGTACG	CAGTTTTATC	CGGTAAAGCG	15300
AATGATTAGA	GGTCTTGGGG	CCGAAACGAT	CTCAAACCTT	TCTCAAACCT	TAAATGGGTA	15360
AGAAGCCCGG	CTCGCTGGCG	TGGAGCCGGG	CGTGGAATGC	GAGTGCCTAG	TGGGCCACTT	15420
TTGGTAAGCA	GAAGTGGCGC	TGCGGGATGA	ACCGAACGCC	GGGTAAAGGC	GCCCCGATCC	15480
GACGCTCATC	AGACCCCAGA	AAAGGTGTTG	GTTGATATAG	ACAGCAGGAC	GGTGGCCATG	15540
GAAGTCGGAA	TCCGCTAAGG	AGTGTGTAAC	AACTCACCTG	CCGAATCAAC	TAGCCCTGAA	15600
AATGGATGGC	GCTGGAGCGT	CGGGCCCCAT	CCCGGCCGTG	GCCGCAGTCG	GAACGGAACG	15660
GGACGGGAGC	GGTCCGGGGT	GCGCGTCTCT	CGGGCTCGGG	GGTGCCTGGC	GGGGGCCCGT	15720
CCCCCGCCTC	CCCTCCGCGC	GCCGGGTTCG	CCCCCGCGGC	GTCGGGCCCC	GCGGAGCCTA	15780
CGCCGCGACG	AGTAGGAGGG	CCGCTGCGGT	GAGCCTTGAA	GCCTAGGGCG	CGGGCCCCGG	15840
TGGAGCCGCC	GCAGGTGCAG	ATCTTGGTGG	TAGTAGCAAA	TATTCAAACG	AGAACTTTGA	15900
AGGCCGAAGT	GGAGAAGGGT	TCCATGTGAA	CAGCAGTTGA	ACATGGGTCA	GTCGGTCTCT	15960
AGAGATGGGC	GAGTGCCGTT	CCGAAGGGAC	GGGCGATGGC	CTCCGTTGCC	CTCGGCCGAT	16020
CGAAAGGGAG	TCGGGTTCAG	ATCCCCGAAT	CCGAGTGGC	GGAGATGGGC	GCCGCGAGGC	16080
CAGTGCGGTA	ACGCGACCGA	TCCCGGAGAA	GCCGCGGGGA	GGCCTCGGGG	AGAGTTCTCT	16140
TTTCTTTGTG	AAGGGCAGGG	CGCCCTGGAA	TGGGTTCGCC	CCGAGAGAGG	GGCCCGTGCC	16200
TTGGAAAGCG	TCGCGGTTCC	GGCGGCGTCC	GGTGAGCTCT	CGCTGGCCCT	TGAAAATCCG	16260
GGGGAGAGGG	TGTAAATCTC	GCGCCGGGCC	GTACCCATAT	CCGCAGCAGG	TCTCCAAGGT	16320
GAACAGCCTC	TGGCATGTTG	GAACAATGTA	GGTAAGGGAA	GTCGGCAAGC	CGGATCCGTA	16380
ACTTCGGGAT	AAGGATTGGC	TCTAAGGGCT	GGGTCCGTCC	GGCTGGGGCG	CGAAGCGGGG	16440
CTGGGCGCGC	GCCGCGGCTG	GACGAGGCGC	CGCCGCCCTC	TCCCACGTCC	GGGGAGACCC	16500
CCCGTCTTTT	CCGCCCCGGG	CCGCCCTCCC	CTCTTCCCCG	CGGGGCCCCG	TCGTCCCCCG	16560
CGTCGTGCGC	ACCTCTCTTC	CCCCCTCCTT	CTTCCCCTCG	GGGGGCGGGT	CGGGGGTCGG	16620
CGCGCGGCGC	GGGCTCCGGG	GCGGCGGGTC	CAACCCCGCG	GGGGTTCCGG	AGCGGGAGGA	16680
ACCAGCGGTC	CCCGGTGGGG	CGGGGGGCGC	GGACACTCGG	GGGGCCGGCG	GCGGCGGCGA	16740
CTCTGGAGGC	GAGGCGGGCC	CTTCCCCTGG	ATCCGCTCAG	CTGCGGCGGG	CGTCGCGGCC	16800
GCTCCCGGGG	AGCCCGGCGG	GTGCCGGCGC	GGGTCCCCCT	CCCGCGGGGC	CTCGCTCCAC	16860
CCCCCATCG	CCTCTCCCGA	GGTGCCTGGC	GGGGGCGGGC	GGGCGTGTCC	CGCGCGTGTG	16920
GGGGGAACCT	CCGCGTCCGT	GTTCCCCCGC	CGGGTCCGCC	CCCCGGGCCG	CGGTTTTCCG	16980
CGCGGCGCCC	CCGCCTCGGC	CGGCGCCTAG	CAGCCGACTT	AGAACTGGTG	CGGACCAGGG	17040
GAATCCGACT	GTTTAATTAA	AACAAAGCAT	CGCGAAGGCC	CGCGGCGGGT	GTTGACGCGA	17100

-210-

TGTGATTTCT	GCCAGTGCT	CTGAATGTCA	AAGTGAAGAA	ATTCAATGAA	GCGCGGGTAA	17160
ACGGCGGGAG	TAAGGCGAGC	TCTCTTAAGG	TAGCCAAATG	CCTCGTCATC	TAATTAGTGA	17220
CGCGCATGAA	TGGATGAACG	AGATTCCCAC	TGTCCCTACC	TACTATCCAG	CGAAACCACA	17280
GCCAAGGGAA	CGGGCTTGGC	GGAAATCAGC	GGGAAAGAAG	ACCCTGTTGA	GCTTGACTCT	17340
AGTCTGGCAC	GGTGAAGAGA	CATGAGAGGT	GTAGAATAAG	TGGGAGGCC	CCGGCGCCCG	17400
GCCCCGTCCT	CGCGTCGGGG	TCGGGGCACC	CCGGCCTCGC	GGGCGGCCCG	TGAAATACCA	17460
CTACTCTCAT	CGTTTTTTCA	CTGACCCGGT	GAGGCGGGGG	GGCGAGCCCC	GAGGGGCTCT	17520
CGCTTCTGGC	GCCAAGCGTC	CGTCCCGCGC	GTGCGGGCGG	GCGCGACCCG	CTCCGGGGAC	17580
AGTGCCAGGT	GGGGAGTTTG	ACTGGGGCGG	TACACCTGTC	AAACGGTAAC	GCAGGTGTCC	17640
TAAGGCGAGC	TCAGGGAGGA	CAGAAACCTC	CCGTGGAGCA	GAAGGGCAAA	AGCTCGCTTG	17700
ATCTTGATTT	TCAGTACGAA	TACAGACCGT	GAAAGCGGGG	CCTCACGATC	CTTCTGACCT	17760
TTTGGGTTTT	AAGCAGGAGG	TGTCAGAAAA	GTTACCACAG	GGATAACTGG	CTTGTGGCGG	17820
CCAAGCGTTC	ATAGCGACGT	CGTTTTTTGA	TCCTTCGATG	TCGGCTCTTC	CTATCATTGT	17880
GAAGCAGAA	TCACCAAGCG	TTGGATTGTT	CACCCACTAA	TAGGGAACGT	GAGCTGGGTT	17940
TAGACCGTCG	TGAGACAGGT	TAGTTTTACC	CTACTGATGA	TGTGTTGTTG	CCATGGTAAT	18000
CCTGCTCAGT	ACGAGAGGAA	CCGCAGGTTT	AGACATTTGG	TGTATGTGCT	TGGCTGAGGA	18060
GCCAATGGGG	CGAAGCTACC	ATCTGTGGGA	TTATGACTGA	ACGCCTCTAA	GTCAGAATCC	18120
GCCCAAGCGG	AACGATACGG	CAGCGCCGAA	GGAGCCTCGG	TTGGCCCCCG	ATAGCCGGGT	18180
CCCCGTCGGT	CCCGCTCGGC	GGGGTCCCCG	CGTCGCCCCG	CGGCGGCGCG	GGGTCTCCCC	18240
CCGCCGGGCG	TCGGGACCGG	GGTCCGGTGC	GGAGAGCCGT	TCGTCTTGGG	AAACGGGGTG	18300
CGGCGGAA	GGGGCCCGCC	CTCTCGCCCC	TCACGTTGAA	CGCACGTTCC	TGTGGAACCT	18360
GGCGCTAAAC	CATTGCTAGA	CGACCTGCTT	CTGGGTCCGG	GTTTCGTACG	TAGCAGACGA	18420
GCTCCCTCGC	TGCGATCTAT	TGAAAGTCAG	CCCTCGACAC	AAGGGTTTGT	CTCTGCGGGC	18480
TTTCCCGTCG	CACGCCCGCT	CGCTCGCACC	CGACCGTGTG	GCCGCCCGGG	CGTCACGGGG	18540
GCGGTCGCCT	CGGCCCCCGC	GCGGTTGCCC	GAACGACCGT	GTGGTGGTTG	GGGGGGGGAT	18600
CGTCTTCTCC	TCCGTCTCCC	GAGGACGGTT	CGTTTCTCTT	TCCCCTTCCG	TCGCTCTCCT	18660
TGGGTGTGGG	AGCCTCGTGC	CGTCGCGACC	GCGGCCCTGCC	GTCGCCTGCC	GCCGCAGCCC	18720
CTTGCCCTCC	GGCCTTGGCC	AAGCCGGAGG	GCGGAGGAGG	GGGATCGGCG	GCGGCGGCGA	18780
CCGCGGCGCG	GTGACGCACG	GTGGGATCCC	CATCCTCGGC	GCGTCCGTCC	GGGACGGCCG	18840
GTGGAGGGGG	CGGGAGGGGT	TTTTCCCGTG	AACGCCGCGT	TCGGCGCCAG	GCCTCTGGCG	18900
GCCGGGGGGG	CGCTCTCTCC	GCCCCGAGAT	CCCCACTCCC	GCCCCCTCCT	TTCGCGCGCC	18960
GCGGCGGCGG	CGTCCGTACG	AGGGGAGGAT	GTCCGCGTGT	GGAGGCGGAG	AGGGTCCGGC	19020
GCGGCGCCTC	TTCCATTTTT	TCCCCCCCCA	CTTCGGAGGT	CGACCAAGTAC	TCCGGGCGAC	19080
ACTTTGTTTT	TTTTTTTTTCC	CCCGATGCTG	GAGGTCGACC	AGATGTCCGA	AAGTGTCCCC	19140
CCCCCCCCCC	CCCCCGGGCG	CGGAGCGGCG	GGGCCACTCT	GGACTCTTTT	TTTTTTTTTTT	19200
TTTTTTTTTT	TTAAATTCCT	GGAACCTTTA	GGTCGACCAG	TTGTCCGTCT	TTTACTCCTT	19260
CATATAGGTC	GACCAAGTCT	CCGGGTGGTA	CTTTGTCTTT	TTCTGAAAAT	CCCAGAGGTC	19320
GACCAGATAT	CCGAAAGTCC	TCTCTTTCCC	TTTACTCTTC	CCCACAGCGA	TTCTCTTTTT	19380
TTTTTTTTTT	TTTGGTGTGC	CTCTTTTTGA	CTTATATACA	TGTAATAGT	GTGTACGTTT	19440
ATATACTTAT	AGGAGGAGGT	CGACCAGTAC	TCCGGGCGAC	ACTTTGTTTT	TTTTTTTTTTT	19500
TCCACCGATG	ATGGAGGTCC	ACCAGATGTC	CGAAAGTGTC	CCGTCCCCCC	CCTCCCCCCC	19560
CCGCGACCGG	GCGGGCTCAC	TCTGGACTCT	TTTTTTTTTT	TTTTTTTTTT	TTTAAATTTT	19620
TGGAACCTTA	AGGTCGACCA	GTTGTCCGTC	TTTCACTCAT	TCATATAGGT	CGACCGGTGG	19680
TACTTTGTCT	TTTTCTGAAA	ATCGCAGAGG	TCGACCAGAT	GTCAGAAAGT	CTGGTGGTCG	19740
ATAAATTATC	TGATCTAGAT	TTGTTTTTCT	GTTTTTCAGT	TTTGTGTTGT	TTTGTGTTGT	19800
TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	19860
TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	19920
TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	19980
TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	TTTGTGTTGT	20040
TACACAAACA	TGACTTTTTT	TTAAATAAAA	TTTAAAAAAT	AAATGCGAAA	ATCGACCAAT	20100
TATCCCTTTC	CTTCTCTCTC	TTTTTTAAAA	ATTTTCTTTG	TGTGTGTGTG	TGTGTGTGTG	20160
TGTGTGTGTG	TGCGTGTGTG	TGTGTGTGTG	CGTGCAGCGT	GCGCGCGCTC	GTTTTATAAA	20220
TACTTATAAT	AATAGGTCGC	CGGGTGGTGG	TAGCTTCCCC	GACTCCAGAG	GCAGAGGCAG	20280
CGAGACTTCT	GAGTTCGAGG	CCAGCCTGGT	CTACAGAGGA	ACCCTGTCTC	GAAAAATGAA	20340
AATAAATAGT	TACATACATA	CATACATACA	TACATACATA	CATACATACA	TACATATGAG	20400
GTTGACCAGT	TGTCATTCCT	TTAGAAATTT	GTTTTTAATT	AATGTGATAG	AGAGATAGAT	20460
AATAGATAGA	TGGATAGAGT	GATACAAATA	TAGGTTTTTT	TTTCAGTAAA	TATGAGGTTG	20520
ATTAACCACT	TTTCCCTTTT	TAGGTTTTTT	TTTTTTTCCC	CTGTCCATGT	GGTTGCTGGG	20580
ATTTGAACTC	AGGACCCTGG	CAGGTCAACT	GGAAAACGTG	TTTTCTATAT	ATATAAATAG	20640
TGGTCTGTCT	GCTGTTTGTT	TGTTTGCTTG	CTTGCTTGCT	TGCTTGCTTG	CTTGCTTGCT	20700
TGCTTTTTTT	TTTCTTCTGA	GACAGTATTT	CTCTGTGTAA	CCTGGTGCCC	TGAAACTCAC	20760

-211-

TCTGTAGACC	AGCCTGGCCT	CAATCGAACT	CAGAAATCCT	CCTGCCTCTT	GTCTACCTCC	20820
CAATTTTGGG	GTAAAGGTGT	GCTACACCAC	TGCCTGGCAT	TATTATCATT	ATCATTATTA	20880
ATTTTATTAT	TAGACAGAAC	GAAATCAACT	AGTTGGTCCT	GTTTCGTTAA	TTCATTTGAA	20940
ATTAGTTGGA	CCAATTAGTT	GGCTGGTTTG	GGAGGTTTCT	TTTGTTTCCG	ATTTGGGTGT	21000
TTGTGGGGCT	GGGGATCAGG	TATCTCAACG	GAATGCATGA	AGGTTAAGGT	GAGATGGCTC	21060
GATTTTGTGA	AAGATTACTT	TTCTTAGTCT	GAGGAAAAAA	TAAAATAATA	TTGGGCTACG	21120
TTTCATTGCT	TCATTTCTAT	TTCTCTTTCT	TTCTTTCTTT	CTTTCAGATA	AGGAGGTCGG	21180
CCAGTTCCTC	CTGCCCTTCTG	GAAGATGTAG	GCATTGCATT	GGGAAAAGCA	TTGTTTGAGA	21240
GATGTGCTAG	TGAACCAGAG	AGTTTGGATG	TCAAGCCGTA	TAATGTTTAT	TACAATATAG	21300
AAAAGTTCTA	ACAAAGTGAT	CTTTAACTTT	TTTTTTTTTT	TTTCTCCTTC	TACTTCTACT	21360
TGTTCTCACT	CTGCCACCAA	CGCGCTTTGT	ACATTGAATG	TGAGCTTTGT	TTTGCTTAAC	21420
AGACATATAT	TTTTTCTTTT	GGTTTTGCTT	GACATGGTTT	CCCTTTCTAT	CCGTGCAGGG	21480
TTCCCAGACG	GCCTTTTGAG	AATAAAAATGG	GAGGCCAGAA	CCAAAGTCTT	TTGAATAAAG	21540
CACCACAAC	CTAACCTGTT	TGGCTGTTTT	CCTTCCCAAG	GCACAGATCT	TTCCCAGCAT	21600
GGAAAAGCAT	GTAGCAGTTG	TAGGACACAC	TAGACGAGAG	CACCAGATCT	CATTGTGGGT	21660
GGTTGTGAAC	CACCCACCAT	GTGGTTGCCT	GGGATTGAA	CTCAGGATCT	TCAGAAGACG	21720
AGTCAGGGCT	CTAAACCGAT	GAGCCATCTC	TCCAGCCCTC	CTACATTCTC	TCTTAAGGCA	21780
TGAATGATCC	CAGCATGGGA	AGACAGTCTG	CCCTCTTTGT	GGTATATCAC	CATATACTCA	21840
ATAAAATAAT	GAAATGAATG	AAGTCTCCAC	GTATTTATTT	CTTCGAGCTA	TCTAAATTCT	21900
CTCACAGCAC	CTCCCCCTCC	CCCACACTGC	CTTTCTCCCT	ATGTTTGGGT	GGGGCTGGGG	21960
GAGGGGTGGG	GTGGGGGCAG	GGATCTGCAT	GTCTTCTTGC	AGGTCTGTGA	ACTATTTGCG	22020
ATGGCCTGGT	TCTCTGAACT	GTTGAGCCTT	GTCTATCCAG	AGGCTGACTG	GCTAGTTTTC	22080
TACCTGAAGT	CCCTGAGTGA	TGATTTCCCT	GTGAATTC			22118

(2) INFORMATION FOR SEQ ID NO:17:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 42999 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

GCTGACACGC	TGTCTCTGCG	CGACCTGTCTG	TCCGAGAGGT	TGGGCCTCCG	GATGCGCGCG	60
GGGCTCTGGC	CTCACGGTGA	CCGGCTAGCC	GGCCGCGCTC	CTGCCTTGAG	CCGCCTGCCG	120
CGGCCCGCGG	GCCTGCTGTT	CTCTCGCGCG	TCCGAGCGTC	CCGACTCCCG	GTGCCGGCCC	180
GGGTCCGGGT	CTCTGACCCA	CCCGGGGGCG	GCGGGGAAGG	CGGCGAGGGC	CACCGTGCCC	240
CGTGCGCTCT	CCGCTGCGGG	CGCCCGGGGC	GCCGCACAAC	CCCACCCGCT	GGCTCCGTGC	300
CGTGCGTGTC	AGGCGTTCTC	GTCTCCGCGG	GGTTGTCCGC	CGCCCCCTTC	CCGGAGTGGG	360
GGGTGGCCGG	AGCCGATCGG	CTCGCTGGCC	GGCCGGCCTC	CGCTCCCGGG	GGGCTCTTCG	420
ATCGATGTGG	TGACGTCTGT	CTCTCCGGGG	CCGGGTCCGA	GCCGCGACGG	GCGAGGGGCG	480
GACGTTCTGT	GCGAACGGGA	CCGTCTTTCT	CGCTCCGCCC	GCGCGGTCCC	CTCGTCTGCT	540
CCTCTCCCCG	CCCGCCGGCC	GGCGTGTGGG	AAGGCGTGGG	GTGCGGACCC	CGGCCCGACC	600
TCGCCGTCCC	GCCCGCCGCC	TTCGCTTCGC	GGGTGCGGGC	CGGCGGGGTC	CTCTGACCGG	660
GCAGACAGCC	CTGCCTGTCTG	CCTCCAGTGG	TTGTGCACTT	GCGGGCGGGC	CCCCTCCGCG	720
GCGGTGGGGG	TGCCGTCCCC	CCGGCCCGTC	GTGCTGCCCT	CTCGGGGGGG	GTTTTCGCGA	780
CGCTCGGCTC	CGCCTGGGCC	CTTGCGGTGC	TCTTGAGCGT	CTCCGGGTTG	TCCCTCAGGT	840
GCCCGAGGCC	GAACGGTGGT	GTGTGCTTCC	CGCCCCCGGC	GCCCCCTCCT	CCGGTCCGCC	900
CCGCGGTGTC	CGCGCGTGGG	TCCTGAGGGA	GCTCGTCCGT	GTGGGGTTCTG	AGGCGGTTTG	960
AGTGAGACGA	GACGAGACGC	GCCCCCTCCA	CGCGGGGAAG	GGCGCCCGCC	TGCTCTCGGT	1020
GAGCGCACGT	CCCGTGCTCC	CCTCTGGCGG	GTGCGCGCGG	GCCGTGTGAG	CGATCGCGGT	1080
GGGTCTGGGG	CGGTGTGACG	CGTGCGCGCG	CCGGCCCGCG	AGGGGCTGCC	GTTCTGCCTC	1140
CGACCGGTCTG	TGTGTGGGTT	GACTTCCGAG	CGCTCTGCC	TCGGAAGGAA	GAGAGTGGGT	1200
GGACGGGGGG	GCCTGGTGGG	GTTGCGCGCA	CGCGCGCACC	GGCCGGGGCC	CCGCCCTGAA	1260

-212-

CGCGAACGCT	CGAGGTGGCC	GCGCGCAGGT	GTTTCCTCGT	ACCGCAGGGC	CCCCTCCCTT	1320
CCCCAGGCGT	CCCTCGGCGC	CTCTGCGGGC	CCGAGGAGGA	GCGGCTGGCG	GGTGGGGGGA	1380
GTGTGACCCA	CCCTCGGTGA	GAAAAGCCTT	CTCTAGCGAT	CTGAGAGGCG	TGCCTTGGGG	1440
GTACCGGATC	CCCCGGGGCCG	CCGCCTCTGT	CTCTGCCTCC	GTTATGGTAG	CGCTGCCGTA	1500
GCGACCCGCT	CGCAGAGGAC	CCTCCTCCGC	TTCCCCCTCG	ACGGGGTTGG	GGGGGAGAAG	1560
CGAGGGTTCC	GCCGGCCACC	GCGGTGGTGG	CCGAGTGC GG	CTCGTCGCCT	ACTGTGGCCC	1620
GCGCCTCCCC	CTTCCGAGTC	GGGGGAGGAT	CCCGCCGGGC	CGGGCCCGGC	GCTCCACCCC	1680
AGCGGGTTGG	GACGCGGCGG	CCGGCGGGCG	GTGGGTGTGC	GCGCCCGGCG	CTCTGTCCGG	1740
CGCGTGACCC	CCTCCGTCCG	CGAGTCGGCT	CTCCGCCCGC	TCCCGTGCCG	AGTCGTGACC	1800
GGTGCCGACG	ACCGCGTTTG	CGTGGCACGG	GGTCGGGCCC	GCCTGGCCCT	GGGAAAGCGT	1860
CCCACGGTGG	GGGCGCGCCG	GTCTCCCGGA	GCGGGACCGG	GTCCGAGGAT	GGACGAGAAT	1920
CACGAGCGAC	GGTGTGGTGG	GCGTGTCCGG	TTCTGTGGCTG	CGGTTCGCTCC	GGGGCCCCCG	1980
GTGGCGGGGC	CCCGGGGCTC	GCGAGGCGGT	TCTCGGTGGG	GGCCGAGGGC	CGTCCGGCGT	2040
CCCAGGCGGG	GCGCCGCGGG	ACCGCCCTCG	TGTCTGTGGC	GGTGGGATCC	CGCGGCCCGTG	2100
TTTTCTTGGT	GGCCCCGGCCG	TGCCGTAGGT	TTCTCCCCGA	GCCGCCGCCT	CTGCGGGCTC	2160
CCGGGTGCCC	TTGCCCTCGC	GGTCCCCGGC	CCTCGCCCGT	CTGTGCCCTC	TTCCCCGCCC	2220
GCCGCGCCCG	GATCCTCTTC	TTCCCCCGGA	CGGGCTCACC	GGCTTCACGT	CCGTGTGGTG	2280
CCCCGCCTGG	GACCGAACCC	GGCACCGCCT	CGTGGGGCGC	CGCCGCCGGC	CACGTATCGG	2340
CCCGGCGTCC	GCGTCCCCCG	GCGCGCGCCT	TGGGGACCGG	GTGCGTGGCG	CGCCGCGTGG	2400
GGCCCGGTGG	GCTTCCCGGA	GGGTTCGGGG	GGTCGGCCTG	CGGCGCGTGC	GGGGGAGGAG	2460
ACGGTTCGGG	GGGACCGGCC	GCGGCTGCGG	CGGCGGCGGT	GGTGGGGGGA	GCCGCGGGGA	2520
TCGCGGAGGG	CCGGTCGGCC	GCCCCGGGTG	CCCCGCGGTG	CCGCCGGCGG	CGGTGAGGCC	2580
CCGCGCGTGT	TCTCCCGGCTG	CGGTCCGGCC	CGCTCGAGGG	GTCCCCGTGG	CGTCCCTTTC	2640
CCCGCCGGCC	GCCTTTCTCG	CGCCTTCCCC	GTCCGCCCGG	CCTCGCCCGT	GGTCTCTCGT	2700
CTTCTCCCGG	CCCGCTCTTC	CGAACCGGGT	CGGCGCGTCC	CCCGGGTGCG	CCTCGCTTCC	2760
CGGGCCTGCC	GCGGCCCTTC	CCCGAGGCGT	CCGTCCCGGG	CGTCGGCGTC	GGGGAGAGCC	2820
CGTCTCTCCC	GCGTGGCGTC	GCCCCGTTTC	GCGCGCGCGT	GCGCCCGAGC	GCGGCCCGGT	2880
GGTCCCTCCC	GGACAGGCGT	TCGTGCGACG	TGTGGCGTGG	GTGACCTTCC	GCCTTGCCGG	2940
TCGCTCGCCC	TCTCCCCGGG	TCGGGGGGTG	GGGCCCGGGC	CGGGGCCTCG	GCCCCGGTCG	3000
CTGCCTCCCG	TCCCGGGCGG	GGGCGGGCGC	GCCGGCCGGC	CTCGGTCCGC	CTCCCTTGGC	3060
CGTCGTGTGG	CGTGTGCCAC	CCCTGCGCCG	GCGCCCGCCG	GCGGGGCTCG	GAGCCGGGCT	3120
TCGGCCGGGC	CCCGGGCCCT	CGACCGGACC	GGCTGCGCGG	GCGCTGCGGC	CGCACGGCGC	3180
GACTGTCCCC	GGGCCGGGCA	CCGCGGTCCG	CCTCTCGCTC	GCCGCCCGGA	CGTCGGGGCC	3240
GCCCCCGGGG	GCGGGCGGAG	CGCCGTCCCC	GCCTCGCCCG	CGCCCGCGGG	CGCCGGCCGC	3300
GCGCGCGCGC	CGGTGGCCGC	CGGTCCCTCC	GCGCCGCGCG	GCGCGGGTCG	GGCCGTCCGC	3360
CTCCTCGCGG	GCGGGCGCGA	CGAAGAAGCG	TCGCGGGTCT	GTGGCGCGGG	GCCCCCGGTG	3420
GTCGTGTGCG	GTGGGGGGCG	GGTGGTTGGG	GCGTCCGGTT	CGCCGCGCCC	CGCCCCGGCC	3480
CCACCGGTCC	CGGCCGCGGC	CCCCGCGCCC	GCTCGCTCCC	TCCCGTCCGC	CCGTCCGCGG	3540
CCCGTCCGTC	CGTCCGTCCG	TCGTCTCTCT	CGCTTGCGGG	GCGCCGGGCC	CGTCTCTCGG	3600
AGGCCCCCGG	CGGCCCGGTC	CGGCCCGGTC	GGGGCTTCGC	CGCGCTCTAC	CTTACCTACC	3660
TGGTTGATCC	TGCCAGTAGC	ATATGCTTGT	CTCAAAGATT	AAGCCATGCA	TGTCTAAGTA	3720
CGCACGGCCG	GTACAGTGAA	ACTGCGAATG	GCTCATTAAT	TCAGTTATGG	TTCTTTTGGT	3780
CGCTCGCTCC	TCTCTACTTT	GGATAACTGT	GGTAATTCTA	GAGCTAATAC	ATGCCGACGG	3840
GCGCTGACCC	CCTTCGCGGG	GGGGATGCGT	GCATTTATCA	GATCAAAACC	AACCCGGTCA	3900
GCCCTCTCTC	GGCCCCGGCC	GGGGGGCGGG	CGCCGGCGGC	TTTGGTGACT	CTAGATAACC	3960
TCGGGCGGAT	CGCACGCCCC	CCGTGGCGGC	GACGACCCAT	TCGAACGTCT	GCCCTATCAA	4020
CTTTCGATGG	TAGTCGCCGT	GCCTACCATG	GTGACCACGG	GTGACGGGGA	ATCAGGGTTC	4080
GATTCCGGAG	AGGGAGCCTG	AGAAACGGCT	ACCACATCCA	AGGAAGGCAG	CAGGCGCGCA	4140
AATTACCCAC	TCCCGACCCG	GGGAGGTAGT	GACGAAAAAT	AACAATACAG	GACTCTTTTCG	4200
AGGCCCTGTA	ATTGGAATGA	GTCCACTTTA	AATCCTTTAA	CGAGGATCCA	TTGGAGGGCA	4260
AGTCTGGTGC	CAGCAGCCGC	GGTAATTCCA	GCTCCAATAG	CGTATATTAA	AGTTGCTGCA	4320
GTTAAAAAGC	TCGTAGTTGG	ATCTTGGGAG	CGGGCGGGCG	GTCCGCCGCG	AGGCGAGCCA	4380
CCGCCCGTCC	CCGCCCTTTC	CCTCTCGGCG	CCCCCTCGAT	GCTCTTAGCT	GAGTGTCCCG	4440
CGGGGCCCGA	AGCGTTTACT	TTGAAAAAAT	TAGAGTGTTT	AAAGCAGGCC	CGAGCCGCCT	4500
GGATACCGCA	GCTAGGAATA	ATGGAATAGG	ACCGCGGTTT	TATTTTGTTG	GTTTTTCGGAA	4560
CTGAGGCCAT	GATTAAGAGG	GACGGCCGGG	GGCATTCGTA	TTGCGCCGCT	AGAGGTGAAA	4620
TTCTTGGGAC	GGCGCAAGAC	GGACCAAGAG	GAAAGCATTT	GCCAAGAATG	TTTTTCATTAA	4680
TCAAGAACGA	AAGTCGGAGG	TTCAAGAGAC	ATCAGATACC	GTCTAGTTTC	CGACCATAAA	4740
CGATGCCGAC	CGGCGATGCG	GCGGCGTTAT	TCCCATGACC	CGCCGGGCAG	CTTCCGGGAA	4800
ACCAAAGTCT	TTGGGTTCCG	GGGGGAGTAT	GGTTGCAAAG	CTGAAACTTA	AAGGAATTGA	4860
CGGAAGGGCA	CCACCAGGAG	TGGAGCCTGC	GGCTTAATTT	GACTCAACAC	GGGAAACCTC	4920

-213-

ACCCGGCCCG	GACACGGACA	GGATTGACAG	ATTGATAGCT	CTTCTCGAT	TCCGTGGGTG	4980
GTGGTGCATG	GCCGTTCTTA	GTTGGTGGAG	CGATTTGTCT	GGTTAATTCC	GATAACGAAC	5040
GAGACTCTGG	CATGCTAACT	AGTTACGCGA	CCCCCGAGCG	GTCGGCGTCC	CCCAACTTCT	5100
TAGAGGGACA	AGTGGCGTTC	AGCCACCCGA	GATTGAGCAA	TAACAGGTCT	GTGATGCCCT	5160
TAGATGTCCG	GGGCTGCACG	CGCGCTACAC	TGACTGGCTC	AGCGTGTGCC	TACCCTACGC	5220
CGGCAGGCGC	GGGTAAACCCG	TTGAACCCCA	TTCGTGATGG	GGATCGGGGA	TTGCAATTAT	5280
TCCCCATGAA	CGAGGGAATT	CCCAGTAAG	TGCGGGTCAT	AAGCTTGCGT	TGATTAAGTC	5340
CCTGCCCTTT	GTACACACCG	CCCGTCGCTA	CTACCGATTG	GATGGTTTAG	TGAGGCCCTC	5400
GGATCGGCC	CGCCGGGGTC	GGCCACGGC	CCTGGCGGAG	CGCTGAGAAG	ACGGTCGAAC	5460
TTGACTATCT	AGAGGAAGTA	AAAGTCGTAA	CAAGGTTTCC	GTAGGTGAAC	CTGCGGAAGG	5520
ATCATTAAAG	GAGCCCGGAG	GGCGAGGCC	GCGGCGGCGC	CGCCGCCGCC	GCGCGCTTCC	5580
CTCCGCACAC	CCACCCCCC	ACCGCGACGC	GGCGCGTGCG	CGGGCGGGGC	CCGCGTGCCC	5640
GTTCTGTTTCG	TGCTCGTTC	GTTGCGCCGC	CGGCCCGGCC	GCCGCGAGAG	CCGAGAAGTC	5700
GGGAGGGAGA	CGGGGGGGAG	AGAGAGAGAG	AGAGAGAGAG	AGAGAGAGAG	AGAGAGAGAA	5760
AGAAGGGCGT	GTCGTTGGTG	TGCGCGTGTC	GTGGGGCCGG	CGGGCGGCGG	GGAGCGGTCC	5820
CCGGCCGCGG	CCCCGACGAC	GTGGGTGTCT	GCGGGCGCGG	GGGCGGTTCT	CGGCGGCGTC	5880
GCGGCGGGTC	TGGGGGGGTC	TGCTGCCCC	CCTCCCCGCC	GGGGCCCGTC	GTCCGGCCCC	5940
GCCGCGCCGG	CTCCCCGTCT	TCGGGGCCGG	CCGGATTCCC	GTCGCCTCCG	CCGCGCCGCT	6000
CCGCGCCGCG	GGGACGCGCC	CCGCTCGCTC	TCCCCGGCCT	TCCCGCTAGG	GCGTCTCGAG	6060
GGTCGGGGGC	CGGACGCGCG	TCCCCTCCCC	CGCCTCCTCG	TCCGCCCCCC	CGCCGTCCAG	6120
GTACCTAGCG	CGTTCCGGCG	CGGAGGTTTA	AAGACCCCTT	GGGGGGATCG	CCCGTCCGCC	6180
CGTGGGTCCG	GGGCGGTGGT	GGGCCCCGCG	GGGAGTCCCG	TCGGGAGGGG	CCCGGCCCTT	6240
CCGCGCCCTC	CACCGCGGAC	TCCGCTCCCC	GGCCGGGGCC	GCGCCGCCGC	CGCCGCCGCG	6300
GCGGCCGTCT	GGTGGGGGCT	TTACCCGGCG	GCCGTCCGCG	GCCTGCCGCG	CGTGTGGCGT	6360
GCGCCCCGCG	CCGTGGGGGC	GGGAACCCCC	GGGCGCCTGT	GGGGTGGTGT	CCGCGCTCGC	6420
CCCCGCGTGG	GCGGCGCGCG	CCTCCCCGTG	GTGTGAAACC	TTCCGACCCC	TCTCCGGAGT	6480
CCGGTCCCGT	TTGCTGTCTC	GTCTGGCCGG	CCTGAGGCAA	CCCCCTCTCC	TCTTGGGCGG	6540
GGGGGGCGGG	GGGACGTGCC	GCGCCAGGAA	GGGCCTCCTC	CCGGTGCCTC	GTGCGGAGCG	6600
CCCTCGCCAA	ATCGACCTCG	TACGACTCTT	AGCGGTGGAT	CACTCGGCTC	GTGCGTGCAT	6660
GAAGAACGCA	GCTAGCTGCG	AGAATTAATG	TGAATTGACG	GACACATTGA	TCATCGACAC	6720
TTCGAACGCA	CTTGCGGCC	CGGGTTCCCT	CCGGGGCTAC	GCCTGTCTGA	GCATCTGTTG	6780
CCGATCAATC	GCCCCGGGGG	TGCTTCCGGG	CTCCTCGGGG	TGCGCGGCTG	GGGGTTCCCT	6840
CGCAGGGCCC	GCCGGGGGCC	CTCCGTCCCC	CTAAGCGCAG	ACCCGGCGGC	GTCCGCCCTC	6900
CTCTTGCCGC	CGCGCCCGCC	CCTTCCCCCT	CCCCCGCGCG	GCCCTGCGTG	GTACGCGCTC	6960
GGGTGGCGGG	GGGGAGAGGG	GGGCGCGCCC	GGCTGAGAGA	GACGGGGAGG	GCGGCGCCGC	7020
CGCCGGAAGA	CGGAGAGGGA	AAGAGAGAGC	CGGCTCGGGC	CGAGTTCCCG	TGGCCGCCGC	7080
CTGCGGTCCG	GGTTCCCTCC	TCGGGGGGCT	CCCTCGCGCC	GCGCGCGGCT	CGGCGGTTCCG	7140
GGTTCGTCCG	CCCCGGCCGG	GTGGAAGGTC	CCGTGCCCCG	CGTCGTCTGC	GTGCGCGCTC	7200
GTCGGCGGTG	GGGGCGTGTT	GCGTGCGGTG	TGGTGGTGGG	GGAGGAGGAA	GGCGGGTCCG	7260
GAAGGGGAAG	GGTGCCGGCG	GGGAGAGAGG	GTGCGGGGAG	CGCGTCCCGG	TCGCGCGGGT	7320
TCCGCCGCC	GCCCCCGGTG	GCGGCCCGGC	GTCCGGCCGA	CCGGCCGCTC	CCCGCGCCCC	7380
TCTCTCTCCC	CGCCCGCCCT	CCTCCGAGGC	CCCGCCCGTC	CTCCTCGCCC	TCCCGCGCGC	7440
TACGCGCGCG	GCCCGGCCCG	CCCGGCTCGC	CTCGCGCGCG	GTGCGCCGGG	GCCGGGAGCC	7500
CGCCCCGCGG	CCCGCCCGTG	GCCGCGGCGC	CGGGGTTTCG	GTGTCCCCCG	CGGCGACCCG	7560
CGGGACGCGG	CGGTGTCTGC	CGCCGTCTGC	CGCCCGCCTC	CGGCTCGCGG	CCGCGCCGCG	7620
CCGCGCCGGG	GCCCCGTCCC	GAGCTTCCGC	GTCGGGGCGG	CGCGGCTCCG	CCGCGCGCTC	7680
CTCGGACCCG	TCCCCCCGAC	CTCCGCGGGG	GAGACGCGCC	GGGGCGTGCG	GCGCCCGTCC	7740
CGCCCCCGGC	CCGTGCCCC	CCCTCCGGTC	GTCCCGCTCC	GGCGGGGCGG	CGCGGGGGCG	7800
CCGTCCGGCG	GCGCTCTCT	CTCCGCTCGC	TCTCCCCCTC	CGCCGGGGCC	GTCTCCCGAC	7860
GGAGCGTCCG	GCGGGCGGTC	GGGCCGCGCG	GATTCCGTCC	GTCCGTCCGC	CGAGCGGCCC	7920
GTCCCCCTCC	GAGACGCGAC	CTCAGATCAG	ACGTGGCGAC	CCGCTGAATT	TAAGCATATT	7980
AGTCAGCGGA	GGAAAAGAAA	CTAACCAGGA	TTCCCTCAGT	AACGGCGAGT	GAACAGGGAA	8040
GAGCCACAGC	CCGAATCCCC	GCCCCGCGGG	GCGCGGGACA	TGTGGCGTAC	GGAAGACCCG	8100
CTCCCCGGCG	CCGCTCGTGG	GGGGCCCCAAG	TCCTTCTGAT	CGAGGCCAG	CCCGTGGACG	8160
GTGTGAGGCC	GGTAGCGGCC	GCGCGCGGCC	CGGGTCTTCC	CGAGTCCGGG	TTGCTTGGGA	8220
ATGCAGCCCA	AAGCGGGTGG	TAAACTCCAT	CTAAGGCTAA	ATACCGGCAC	GAGACCGATA	8280
GTCAACAAGT	ACCGTAAGGG	AAAGTTGAAA	AGAACTTTGA	AGAGAGAGTT	CAAGAGGGCG	8340
TGAAACCGTT	AAGAGGTAAA	CGGGTGGGGT	CCGCGCAGTC	CGCCCGGAGG	ATTCAACCCG	8400
GCGGCGGGTC	CGGCCGTGTC	GGCGGCCCGG	CGGATCTTTC	CCGCCCCCCG	TTCCTCCCGA	8460
CCCCTCCACC	CGCCCTCCCT	TCCCCCGCCG	CCCCTCCTCC	TCTCCCCCGG	AGGGGGCGGG	8520
CTCCGGCGGG	TGCGGGGGTG	GGCGGGCGGG	GCCGGGGGTG	GGGTCGGCGG	GGGACCGTCC	8580

-214-

CCCGACCGGC	GACCGGCCGC	CGCCGGGCGC	ATTTCCACCG	CGGCGGTGCG	CCGCGACCGG	8640
CTCCGGGACG	GCTGGGAAGG	CCCGGCGGGG	AAGGTGGCTC	GGGGGGCCCC	GTCCGTCCGT	8700
CCGTCTCTCT	CCTCCCCCGT	CTCCGCCCCC	CGGCCCCGCG	TCCTCCCTCG	GGAGGGCGCG	8760
CGGGTCGGGG	CGGCGGCGGC	GGCGGCGGTG	GCGGCGGCGG	CGGGGGCGGC	GGGACCGAAA	8820
CCCCCCCCGA	GTGTTACAGC	CCCCCGGCA	GCAGCACTCG	CCGAATCCCG	GGGCCGAGGG	8880
AGCGAGACCC	GTCGCGCGCG	TCTCCCCCT	CCCGGCGCCC	ACCCCGCGCG	GGAATCCCCC	8940
GCGAGGGGGG	TCTCCCCCGC	GGGGGCGCGC	CGGCGTCTCC	TCGTGGGGGG	GCCGGGCCAC	9000
CCCTCCCACG	GCGCGACCGC	TCTCCCACCC	CTCCTCCCCG	CGCCCCCGCC	CCGGCGACCG	9060
GGGGGGTGCC	GCGCGCGGGT	CGGGGGGCGG	GGCGGACTGT	CCCCAGTGCG	CCCCGGGCGG	9120
GTCGCGCCGT	CGGGCCCCGG	GGAGGTTCTC	TCGGGGCCAC	GCGCGCGTCC	CCCGAAGAGG	9180
GGGACGGCGG	AGCGAGCGCA	CGGGGTCGGC	GGCGACGTCT	GCTACCCACC	CGACCCGTCT	9240
TGAAACACGG	ACCAAGGAGT	CTAACACGTG	CGCGAGTCGG	GGGCTCGCAC	GAAAGCCGCC	9300
GTGGCGCAAT	GAAGGTGAAG	GCCGGCGCGC	TCGCCGGCCG	AGGTGGGATC	CCGAGGCCTC	9360
TCCAGTCCGC	CGAGGGCGCA	CCACCGGCCC	GTCTCGCCCG	CCGCGCCGGG	GAGGTGGAGC	9420
ACGAGCGCAC	GTGTTAGGAC	CCGAAAGATG	GTGAACTATG	CCTGGGCAGG	GCGAAGCCAG	9480
AGGAAACTCT	GGTGGAGGTC	CGTAGCGGTG	CTGACGTGCA	AATCGGTCTG	CCGACCTGGG	9540
TATAGGGGCG	AAAGACTAAT	CGAACCCTCT	AGTAGCTGGT	TCCCTCCGAA	GTTCCTCTCA	9600
GGATAGCTGG	CGCTCTCGCA	GACCCGACGC	ACCCCGGCCA	CGCAGTTTTA	TCCGGTAAAG	9660
CGAATGATTA	GAGGTCTTGG	GGCCGAAACG	ATCTCAACCT	ATTCTCAAAC	TTTAAATGGG	9720
TAAGAAGCCC	GGCTCGCTGG	CGTGGAGCCG	GGCGTGGAAT	GCGAGTGCCT	AGTGGGCCAC	9780
TTTTGGTAAG	CAGAACTGGC	GCTGCGGGAT	GAACCGAACG	CCGGGTAAAG	GCGCCCGATG	9840
CCGACGCTCA	TCAGACCCCA	GAAAAGGTGT	TGGTTGATAT	AGACAGCAGG	ACGGTGGCCA	9900
TGGAAGTCGG	AATCCGCTAA	GGAGTGTGTA	ACAACCTACC	TGCCGAATCA	ACTAGCCCTG	9960
AAAATGGATG	GCGCTGGAGC	GTCGGGCCCA	TACCCGGCCG	TCGCCGGCAG	TCGAGAGTGG	10020
ACGGGAGCGG	CGGGGGCGGC	GCGCGCGCGC	GCGCGTGTGG	TGTGCGTCGG	AGGGCGGCGG	10080
CGGCGGCGGC	GGCGGGGGTG	TGGGGTCCTT	CCCCCGCCCC	CCCCCCCCAC	CCTCCTCCCC	10140
TCCTCCCGCC	CACGCCCCGC	TCCCCGCCCC	CGGAGCCCCG	CGGACGCTAC	GCCGCGACGA	10200
GTAGGAGGGC	AGCTGCGGTG	AGCCTTGAAG	CTAGGGCGCG	GGGCCCCGGT	GGAGCCGCGG	10260
CAGGTGCAGA	TCTTGGTGGT	AGTAGCAAAT	ATTCAAACGA	GAACCTTGAA	GGCCGAAGTG	10320
GAGAAGGGTT	CCATGTGAAC	AGCAGTTGAA	CATGGGTCAG	TCGGTCCTGA	GAGATGGGCG	10380
AGCGCCGTTT	CGAAGGGACG	GGCGATGGCC	TCCGTTGCCC	TCGGCCGATC	GAAAGGGAGT	10440
CGGGTTCAGA	TCCCCGAATC	CGGAGTGGCG	GAGATGGGCG	CCGCGAGGCG	TCCAGTGCAG	10500
TAACGCGACC	GATCCCGGAG	AAGCCGGCGG	GAGCCCCGGG	GAGAGTTCTC	TTTTCTTTGT	10560
GAAGGGCAGG	ATGGCTTGGG	CCCGAGAGAG	CCCGAGAGAG	GGGCCCGTGC	CTTGGAAAGC	10620
GTCGCGGTTT	CGGCGGCGTC	CGGTGAGCTC	TCGTTGGCCC	TTGAAAATCC	GGGGGAGAGG	10680
GTGTAAATCT	CGCGCCGGGC	CGTACCCATA	TCCGCAGCAG	GTCTCCAAGG	TGAACAGCCT	10740
CTGGCATGTT	GGAACAATGT	AGGTAAGGGA	AGTCGGCAAG	CCGGATCCGT	AACTTCGGGA	10800
TAAGGATTGG	CTCTAAGGGC	TGGGTCGGTC	GGGCTGGGGC	GCGAAGCGGG	GCTGGGCGCG	10860
CGCCGCGGCT	GGACGAGGCG	CGCGCCCCCT	CCACGCCCGG	GGCACCCCCC	TCGCGGCCCT	10920
CCCCCGCCCC	ACCCGCGCGC	GCCGCTCGCT	CCCTCCCCAC	CCCGCGCCCT	CTCTCTCTCT	10980
CTCTCCCCCG	CTCCCCGTCC	TCCCCCTCTC	CCGGGGGAGC	GCCGCGTGGG	GGCGGGGCGG	11040
GGGGAGAAGG	GTCGGGGCGG	CAGGGGCCGC	GCGGCGGCCG	CCGGGGCGGC	CGGCGGGGGC	11100
AGGTCCCCCG	GAGGGGGGGC	CCGGGGACCC	GGGGGGCCGG	CGGCGGCGCG	GACTCTGGAC	11160
GCGAGCCGGG	CCCTTCCCGT	GGATCGCCCC	AGCTGCGGCG	GGCGTCGCGG	CCGCCCCCGG	11220
GGAGCCCGCG	GGCGGCGCGG	CGCGCCCCCC	ACCCCCACCC	CACGTCTCGG	TCGCGCGCGC	11280
GTCCGCTGGG	GGCGGGAGCG	GTCGGGCGGC	GGCGGTCGGC	GGGCGGCGGG	GCGGGGCGGT	11340
TCGTCCCCCC	GCCCTACCCC	CCCGGCCCCG	TCCGCCCCCC	GTTCCCCCCT	CCTCCTCGGC	11400
GCGCGGCGGC	GGCGGCGGCA	GGCGGCGGAG	GGGCCGCGGG	CCGGTCCCCC	CCGCGGGGTC	11460
CGCCCCCGGG	GCCGCGGTTT	CGCGCGCGCC	TCGCCTCGGC	CGGCGCCTAG	CAGCCGACTT	11520
AGAAGTGGTG	CGGACCAGGG	GAATCCGACT	GTTTAATTAA	AACAAAGCAT	CGCGAAGGCC	11580
CGCGGCGGGT	GTTGACGCGA	TGTGATTTCT	GCCCAGTGCT	CTGAATGTCA	AAGTGAAGAA	11640
ATTCAATGAA	ACGGCGGGAG	ACGGCGGGAG	TAACATGAC	TCTCTTAAGG	TAGTCAAATG	11700
CCTCGTCATC	TAATTAGTGA	CGCGCATGAA	TGGATGAACG	AGATTCCAC	TGTCCCTACC	11760
TACTATCCAG	CGAAACCACA	GCCAAGGGAA	CGGGCTTGCC	GGAATCAGCG	GGGAAAGAAG	11820
ACCCTGTTGA	GCTTGACTCT	AGTCTGGCAC	GGTGAAGAGA	CATGAGAGGT	GTAAGAATAAG	11880
TGGGAGGCCC	CCGGCGCCCC	CCCGGTGTCC	CCGCGAGGGG	CCCGGGGCGG	GGTCCGCGGC	11940
CCTGCGGGCC	GCCGGTGAAA	TACCACTACT	CTGATCGTTT	TTTCACTGAC	CCGGTGAGGC	12000
GGGGGGGCGA	CCCCGAGGGG	CTCTCGCTTC	TGGCGCCAAG	CGCCCGCCCC	CGGGGGCGCG	12060
ACCCGCTCCG	GGGACAGTGC	CAGGTGGGGA	GTTTGACTGG	GGCGGTACAC	CTGTCAAACG	12120
GTAACGCAGG	TGTCCTAAGG	CGAGCTCAGG	GAGGACAGAA	ACCTCCCGTG	GAGCAGAAGG	12180
GCAAAAGCTC	GCTTGATCTT	GATTTTCAGT	ACGAATACAG	ACCGTGAAAG	CGGGGCCCTCA	12240

-215-

CGATCCTTCT	GACCTTTTGG	GTTTTAAGCA	GGAGGTGTCA	GAAAAGTTAC	CACAGGGATA	12300
ACTGGCTTGT	GGCGGCCAAG	CGTTCATAGC	GACGTCGCTT	TTTGATCCTT	CGATGTCGGC	12360
TCTTCCTATC	ATTGTGAAGC	AGAATTTCGC	AAGCGTTGGA	TTGTTACACC	ACTAATAGGG	12420
AACGTGAGCT	GGGTTTAGAC	CGTCGTGAGA	CAGGTTAGTT	TTACCCTACT	GATGATGTGT	12480
TGTTGCCATG	GTAATCCTGC	TCAGTACGAG	AGGAACCGCA	GGTTCAGACA	TTTGGTGTAT	12540
GTGCTTGGCT	GAGGAGCCAA	TGGGGCGAAG	CTACCATCTG	TGGGATTATG	ACTGAACGCC	12600
TCTAAGTCAG	AATCCCGCCC	AGGCGAACGA	TACGGCAGCG	CCGCGGAGCC	TCGGTGGGCC	12660
TCGGATAGCC	GGTCCCCCGC	CTGTCCCCGC	CGGCGGGCCG	CCCCCCCCCT	CACGCGCCCC	12720
GCCGCGGGAG	GGCGCGTGCC	CCGCGCGCGC	CCGGGACCGG	GGTCCGGTGC	GGAGTGCCCT	12780
TCGTCTGGG	AAACGGGGCG	CGGCCGGAAA	GGCGGCCGCC	CCCTCGCCCC	TCACGCACCG	12840
CACGTTCTGT	GGGAACCTGG	CGCTAAACCA	TCGTAGACG	ACCTGCTTCT	GGGTCGGGGT	12900
TTCGTACGTA	GCAGAGCAGC	TCCCTCGCTG	CGATCTATTG	AAAGTCAGCC	CTCGACACAA	12960
GGGTTTGTCC	CGCGCGCGCT	GCGTCCGGGG	GGCCCGGCGG	GCGTGCCTGT	TCGGCGCCGT	13020
CCGTCTTCTC	GTTCTCTTTC	CTCCCTCCCC	GCCTCTCCCC	CCGACCGCGG	CGTGGTGGTG	13080
GGGTGGGGGG	GAGGGCGCGC	GACCCCGGTC	GGCCGCCCCG	CTTCTTCGGT	TCCCGCCTCC	13140
TCCCCGTTCA	CGCCGGGGCG	GCTCGTCCGC	TCCGGGCCGG	GACGGGGTCC	GGGGAGCGTG	13200
GTTTGGGAGC	CGCGGAGGCG	CCGCGCCGAG	CCGGGCCCCG	TGGCCCCCGG	GTCCCCGTCC	13260
CGGGGGTTGG	CCGCGCGGCG	CGGTGGGGGG	CCACCCGGGG	TCCCGGCCCT	CGCGCGTCTT	13320
TCCTCCTCGC	TCCTCCGCAC	GGGTCCGACG	ACGAACCGCG	GCTGGCGGGC	GGCGGGCGGC	13380
GAGCCCCACG	GGCGTCCCCG	CACCCGGCCG	ACCTCCGCTC	GCGACCTCTC	CTCGGTCCGG	13440
CCTCCGGGGT	CGACCGCCTG	CGCCCGCGGG	CGTGAGACTC	AGCGGCGTCT	CGCCGTGTCC	13500
CGGGTCGACC	GCGGCCTTCT	CCACCGAGCG	GCGGTGTAGG	AGTGCCCGTC	GGGACGAACC	13560
GCAACCGGAG	CGTCCCCGTC	TCGGTTCGGCA	CCTCCGGGGT	CGACCAGCTG	CCGCCCCGCA	13620
GCTCCGGACT	TAGCCGGCGT	CTGCACGTGT	CCCGGGTCGA	CCAGCAGGCG	GCCGCCGGAC	13680
GCAGCGGCGC	ACGCACGCGA	GGGCGTCGAT	TCCCTTCGCG	GCGCCCCGCG	CTCCACCGGC	13740
CTCGGGCCGC	GGTGGAGCTG	GGACCACGCG	GAACCTCCCT	TCCCACATTT	TTTTTCAGCCC	13800
CACCGCGAGT	TTGCGTCCGC	GGGACCTTTA	AGAGGGAGTC	ACTGCTGCCG	TCAGCCAGTA	13860
CTGCCTCCTC	CTTTTTCGCT	TTTAGGTTTT	GCTTGCCCTT	TTTTTTTTTT	TTTTTTTTTT	13920
TTTTTTCTTT	CTTCTTTTCT	TTCTTTCTTT	CTTCTTTTCT	TTCTTTCTTT	CGCTTGCTCT	13980
CTTCTTGTGT	TCTCTTCTTG	CTCTTCTCTT	GTCTGTCTCT	CTCTCTCTCT	CTCTCTCTGT	14040
CTCTCGCTCT	CGCCCTCTCT	CTCTCTCTCT	TCTCTCTCTC	TCTCTCTCTG	TCTCTCTCTC	14100
TCGCCCTCTC	TCTCTCTCTT	CTCTCTCTCT	CTCTCTCTCT	CTCTCTCTCT	CTCTCTCTCT	14160
GTGCTCTCTG	CCCTCTCTCT	CTCTCTCTGT	CTCTGTCTGT	GTCTCTCTCT	CTCCCTCCCT	14220
CCCTCCCTCC	CTCCCTCCCT	CCCTCCCTCT	CCTTGCGCGC	TTCTCGGCTC	TTGAGACTTA	14280
GCCGCTGTCT	CGCCGTACCC	CGGGTCGACC	GGCGGGCCTT	CTCCACCGAG	CGGCGTGCCA	14340
CAGTGCCTGT	CGGGACGAGC	CGGACCCGCC	GCGTCCCCGT	CTCGGTCCGG	ACCTCCGGGG	14400
TCGACCAGCT	CGCGCCCGCG	AGCTCCGGAG	TTAGCCGGCG	TCTGCACGTG	TCCCGGCTCG	14460
ACCAGCAGGC	GGCCGCCGGA	CGCAGCGGCG	CACCGACGGA	GGGCGCTGAT	TCCCGTTTAC	14520
GCGCCCGCGC	CTCCACCGGC	CTCGGCCCGC	CGTGGAGCTG	GGACCACGCG	GAACCTCCCT	14580
TCCTACATTT	TTTTTCAGCCC	CACCGCGAGT	TTGCGTCCGC	GGGACCTTTA	AGAGGGAGTC	14640
ACTGCTGCCG	TCAGCCAGTA	CTGCCTCCTC	CTTTTTCGCT	TTTAGGTTTT	GCTTGCCCTT	14700
TTTTTTTTTT	TTTTTTTTTT	TTTTTTCTTT	CTTTCTTTCT	TTCTTTCTTT	CTTTCTTTCT	14760
TTCTTTCTTT	CTTTCTTTCT	CGCTCTCTCG	CTCTCTCTCT	CGCTCGTTTC	TTTCTTTCTC	14820
TTTCTCTCTC	TCTCTCTCTC	TCTCTCTCTC	TCTGTCTCTC	GCTCTCGCCC	TCTCTCTCTC	14880
TTTCTCTCTC	TCTCTGTCTC	TCTCTCTCTC	TCTCTCTCTC	TCTCTCTCTC	CCTCCCTCCC	14940
TCCCCCTCCC	TCCCTCTCTC	CCCTTCCCTG	GCGCCTTCTC	GGCTCTTGAG	ACTTAGCCGC	15000
TGTCTCGCCG	TGTCCCGGGT	CGACCGGCGG	GCCTTCTCCA	CCGAGCGGCG	TGCCACAGTG	15060
CCCGTCGGGA	CGAGCCGGAC	CCGCCCGGTC	CCCGCTCTCG	TCGGCACCTC	CGGGGTCGAC	15120
CAGCTGCCGC	CCGCGAGCTC	CGGACTTAGC	CGGCTCTGCG	ACGTGTCCCG	GCTCGACCAG	15180
CAGGCGGCCG	CCGGACGCTG	CGGCGCACCG	ACGCGAGGGC	GTCGATTCCG	GTTACGCGCG	15240
CGGCGACCTC	CACCGGCCCT	GGCCCGCGGT	GGAGCTGGGA	CCACGCGGAA	CTCCCTCTCC	15300
CACATTTTTT	TCAGCCCCAC	CGCGAGTTTG	CGTCCGCGGG	ACTTTTAAGA	GGGAGTCACT	15360
GCTGCCGTCA	GCCAGTAATG	CTTCTCTCTT	TTTTGCTTTT	TGGTTTTGCC	TTGCGTTTTT	15420
TTTCTTTCTT	TCTTTCTTTT	TTTCTTTCTT	TCTTTCTTTT	TCTCTCTCTC	TCTCTCTCTC	15480
TCTCTGTCTC	TCTCTCTCTG	TCTCTCTCTC	CTCCCTCCCT	CCTTGGTGCC	TTCTCTCTCT	15540
GCTGTCTCTG	CTGCCCTCTG	CTCCACGGTT	CAAGCAAACA	GCAAGTTTTT	TATTTTCGAGT	15600
AAAGACGTAA	TTTCACCATT	TTGGCCGGGC	TGGTCTCGAA	CTCCCGACCT	AGTGATCCGC	15660
CCGCCTCGGC	CTCCCAAAGA	CTGCTGGGAG	TACAGATGTG	AGCCACCATG	CCCGGCCGAT	15720
TCCTTCTCTT	TTTCAATCTT	ATTTTCTGAA	CGCTGCCGTG	TATGAACATA	CATCTACACA	15780
CACACACACA	CACACACACA	CACACACACA	CACACACACA	CACACACCCC	GTAAGTGTAA	15840
AACTATGTAA	ATGATATTTT	CATAATTAAT	ACGTTTATAT	TATGTTACTT	TTAATGGATG	15900

-216-

AATATGTATC	GAAGCCCCAT	TTCATTTACA	TACACGTGTA	TGTATATCCT	TCCTCCCTTC	15960
CTTCATTCAT	TATTTATTAA	TAATTTTCGT	TTATTTATTT	TCCTTTCTTT	TGGGGCCGGC	16020
CCGCCTGGTC	TTCTGTCTCT	GCGCTCTGGT	GACCTCAGCC	TCCCAAATAG	CTGGGACTAC	16080
AGGGATCTCT	TAAGCCCGGG	AGGAGAGGTT	AACGTGGGCT	GTGATCGCAC	ACTTCCACTC	16140
CAGCTTACGT	GGGCTGCGGT	GCGGTGGGGT	GGGGTGGGGT	GGGGTGGGGT	GCAGAGAAAA	16200
CGATTGATTG	CGATCTCAAT	TGCCCTTTAG	CTTCATTCAT	ACCCTGTTAT	TTGCTCGTTT	16260
ATTCTCATGG	GTTCTTCTGT	GTCATTGTCA	CGTTCATCGT	TTGCTTGCCCT	GCTTGCCTGT	16320
TTATTTCTTT	CCTTCCTTCC	TTCCTTCTCT	CCTTCTCTCC	TTCCTTCTCT	CCCTCCCTTA	16380
CTGGCAGGGT	CTTCTCTGT	CTCTGCCGCC	CAGGATCACC	CCAACCTCAA	CGCTTTGGAC	16440
CGACCAAACG	GTCGTTCTGC	CTCTGATCCC	TCCCATCCCC	ATTACCTGAG	ACTACAGGCG	16500
CGCACCACCA	CACCGGCTGA	CTTTTATGTT	GTTTCTCATG	TTTTCCGTAG	GATAGTATGT	16560
GTGTGTGTGT	GTGTGTGTGT	GTGTGTGTGT	GTGTGTGTGT	GTGTGTGTGT	GTGTGTATCT	16620
ATGTATGTAC	GTATGTATGT	ATGTATGTGA	GTGAGATGGG	TTTCGGGGTT	CTATCATGTT	16680
GCCCCAGCTG	GTCTCGAACT	CCTGTCTCTA	AGCAATCCGC	CTGCCTGCCT	CGGCCGCCCA	16740
CACTGCTGCT	ATTACAGGCG	TGAGACGCTG	CGCCTGGCTC	CTTCTACATT	TGCCTGCCTG	16800
CCTTGCCTGC	TGCCTGCCTA	TCAATCGTCT	TCTTTTGTAG	ACGGATGTCT	TCTCGCTTTA	16860
TTGTCCATGC	TCTGGGCACA	CGTGGTCTCT	TTTCAAACTT	CTATGATTAT	TATTATTGTA	16920
GGCGTCATCT	CACGTGTCTG	GGTGATCTCG	AACTTTTAGG	CTCCAGAGAT	CCTCCCGCAT	16980
CGGCCTCCCG	GAGTGCTGTG	ATGACACGCG	TGGGCACGGT	ACGCTCTGGT	CGTGTGTTGTC	17040
GTGGGTCCGGT	TCTTTCCGTT	TTTAATACGG	GGACTGCGAA	CGAAGAAAAAT	TTTCAGACGC	17100
ATCTCACCCG	TCCGCCTTTT	CGTTCCTTCT	TTTTATTCTC	TTTAGACGGA	GTTTCACTCT	17160
TGTCGCCCGA	GGTGGAGTAC	GATGGCGGCT	CTCGGCTCAC	CGCACCTTCC	GCTCCCAAGT	17220
TTCAAGTGAT	TCTCCTGCCT	CAGCCTTCCC	GAGTAGCTGG	AATGACAGAG	ATGAGCCATC	17280
GTGCCCCGGT	AATTTTTCTA	TTTTTAGTAC	AGATGGGGTT	TCTCCATCTT	GGTCAGGCTG	17340
GTCTTCAACT	TCCGACCGTT	GGAGAATCTT	AACTTTCTTG	GTGGTGGTTG	TTTTCTTTTT	17400
TCTTTTTTTT	TCTTTTCTTT	TCTTTCTCTC	TCTTCCCCCC	CCCACCCCCC	TTGTCTGCTG	17460
CCTCCTCCTC	CTCCTCCTCC	TCTCCTCCTC	CCTCCTCCTC	CTCCTCCTCC	TCTTTCACTT	17520
CTTTTCAGCTG	GGCTCTCCTA	CTTGTGTTGC	TCTGTGCTCT	ACGCTGGTCT	CAAACTCCTG	17580
GCCTTGACTC	TTCTCCCGTC	ACATCCGCCG	TCTGGTTGTT	GAAATGAGCA	TCTCTCGTAA	17640
AATGGAAAAG	ATGAAAGAAA	TAAACACGAA	GACGGAAAGC	ACGGTGTGAA	CGTTTCTCTT	17700
GCCGTCTCCC	GGGGTGTACC	TTGGACCCGG	AAACACGGAG	GGAGCTTGGC	TGAGTGGGTT	17760
TTCCGTGCGG	AAACCTCCCG	AGGGCCTCCT	TCCCTCTCCC	CCTTGTCCCC	GCTTCTCCGC	17820
CAGCCGAGCG	TCCCACCGCC	GCCCCCTGCA	TTTTCCATAG	GAGAGGTATG	GGAGCCGACT	17880
GACACGCCCT	CCAGATCTAT	ATCCTGCCCG	ACGTCTCTGG	CTCGGCGTGC	CCCACCGGCT	17940
ACCTGCCACC	TTCCAGGGAG	CTCTGAGGCG	GATGCGACCC	CCACCCCCCC	GTCACGTCCC	18000
GCTACCCTCC	CCCGGCTGGC	CTTTGCCGGG	CGACCCCGAG	GGAACCGCGT	TGATGCTGCT	18060
TCGGATCCTC	CGGCGAAGAC	TTCCACCGGA	TGCCCCGGGT	GGGCCGGTTG	GGATCAGACT	18120
GGACCAACCC	GGACCGTGCT	GTTCTTGGGG	GTGGGTTGAC	GTACAGGGTG	GACTGGCAGC	18180
CCCAGCATTT	TAAAGGGTGC	GTGGGTATGG	AAATGTCACC	TAGGATGCCC	TCTTCTCCTT	18240
CGGTCTGCCT	TCAGCTGCCT	CAGGCGTGAA	GACAACTTCC	CATCGGAACC	TCTTCTCTTC	18300
CCTTTCTCCA	GCACACAGAT	GAGACGCACG	AGAGGGAGAA	ACAGCTCAAT	AGATAACGCT	18360
GACCTTCATT	TGTGGAATCC	TCAGTCATCG	ACACACAAGA	CAGGTGACTA	GGCAGGGACA	18420
CAGATCAAAC	ACTATTTCCG	GGTCCTCGTG	GTGGGATTGG	TCTCTCTCTC	TCTCTCTCTC	18480
TCTCTCTCTC	TCTCTCTCTC	TCTCGCACGC	GCACGCGCGC	ACACACACAC	ACAATTTCCA	18540
TATCTAGTTC	ACAGAGCACA	CTCACTTCCC	CTTTTCACAG	TACGCAGGCT	GAGTAAAACG	18600
CGCCCCACCC	TCCACCCGTT	GGCTGACGAA	ACCCCTTCTC	TACAATTGAT	GAAAAAGATG	18660
ATCTGGGCCG	GGCACGCTAG	CTCACGCCTG	TCACTCCGGC	ACTTTGGGAG	GCCGAGGCGG	18720
GTGGATCGCT	TGGGGCCGGG	AGTTCGAGAC	CAGGCTGGCC	GACGTGGCGA	AACCCCGTCT	18780
CTCTGAAAAA	TAGAACGATT	AGCCGGGCCT	GGTGGCGTGG	GCTTGGAAATC	ACGACCGCTC	18840
GGGAGACTTG	GGCGGGCGAC	TTGTTCCAAC	CGGGGAGGCC	GAGGCCGCGA	TGAAGTCAGA	18900
TCGTGACCTG	GCGATGCGGC	CTGGATGACG	GAGCGAGACC	CCGTCTCGAG	AGAATCATGA	18960
TGTTATTATA	AGATGAGTTG	TGCGCGGTGA	TGGCCGCCTG	TAGTCGCGGC	TACTCGGGAG	19020
GCTGAGACGA	GGAGAAGATC	ACTTGAGGCC	CCACAGGTCG	AGGCTTCGGT	CGGCCGTGAC	19080
CCACTGTATC	CTGGGCAGTC	ACCGGTCAAG	GAGATATGCC	CCTTCCCCGT	TTGCTTTTCT	19140
TTCTTCCCTT	TCTCTTTTCT	TCTTTTTTGT	TCTCTTTTCT	TTCTTTCTTT	CTTTCTTTCT	19200
TTCTTTCTTT	CTTTCTTTCT	TTTTCTTTTT	CTCTCTTCCC	CTCTTTCTTT	CTGCTCTTCC	19260
TGCCCTTCTT	CTTTCTTTCT	TTCCTCCCTT	CTCTCTTCCC	TTCTTTCTCT	CCGCCCTCAGC	19320
CTCCCAAAGT	GCTGGGATGA	CTGGCGGGAG	GCACCATGCC	TGCTTGGCCC	AAAGAGACCC	19380
TCTTGGAAAAG	TGAGACGCAG	AGAGCGCCTT	CCAGTGATCT	CATTGACTGA	TTTAGAGACG	19440
GCACTCTCGCT	CCGTCACCCC	GGCAGTGGTG	CCGTCTGTAAC	TCACTCCCTG	CAGCGTGGAC	19500
GCTCCTGGAC	TCGAGCGATC	CTTCCACCTC	AGCCTCCAGA	GTACAGAGCC	TGGGACCGCG	19560

-217-

GGCACGCGCC	ACTGTGCCCA	CACCGTTTTT	AATTGTTTTT	TTTCCCCCG	AGACAGAGTT	19620
TCACTCTCGT	GGCCTAGACT	GCAGTGCGGT	GGCGCGATCT	TGGCTCACC	CAACCTCTGC	19680
CTCCCGGTTT	CAAGCGATT	TCCTGCATCG	GCCTCCTGAG	TAGCCGGGAT	TGCGGGCATG	19740
CGCTGCCACG	TCTGGCTGAT	TTCTGATTTT	TAGTGGAGAC	GGGGCTTCTC	CATGTCTGATC	19800
GGGCTGGTTT	CGAACTCCCG	ACCTCAGGTG	ATCCGCCCTC	CCCGGCCTCC	GGAAGTGCTG	19860
GGATGACAGG	CGTGAGCCAC	CGCGCCCGGC	CTTCATTTTT	AAATGTTTTT	CCACAGACGG	19920
GGTCTCATCA	TTTCTTTGCA	ACCTCCTGTC	CCGGCGTCTC	AAAGTGCTGG	CGTGACGGG	19980
GTGAGCCACT	GCGCCTGGAC	TCCGGGGAAT	GACTCACGAC	CACCATCGCT	CTACTGATCC	20040
TTTCTTTCTT	TCTTTCTTTC	TTTCTTTCTT	TCTTTCTTTC	TTTCTTTCTT	TCTTTCTTGA	20100
TGAATTATCT	TATGATTTAT	TTGTGTACTT	ATTTTCAGAC	GGAGTCTCGC	TCTGGGCGGG	20160
GCGAGGCGAG	GCGAGGCACA	GCGCATCGCT	TTGGAAGCCG	CGGCAACGCC	TTTCAAAGCC	20220
CCATTTCGTAT	GCACAGAGCC	TTATTCCCTT	CTTGGAGTTG	GAGCTGATGC	CTTCCGTAGC	20280
CTTGGGCTTC	TCTCCATTCT	GAAGCTTGAC	AGGCGCAGGG	CCACCCAGAG	GCTGGCTGCG	20340
GCTGAGGATT	AGGGGGTGTG	TTGGGGCTGA	AAACTGGGTC	CCCTATTTTT	GATACCTCAG	20400
CCGACACATC	CCCCGACCGC	CATCGCTTGC	TCGCCCTCTG	AGATCCCCCG	CCTCCACCGC	20460
CTTGACAGGCT	CACCTCTTAC	TTTCATTTCT	TCCTTTCTTG	CGTTTGAGGA	GGGGGTGCGG	20520
GAATGAGGGT	GTGTGTGGGG	AGGGGGTGCG	GGGTGGGGAC	GGAGGGGAGC	GTCTTAAGGG	20580
TCGATTTAGT	GTCTATGCCTC	TTTCACCACC	ACCACCACCA	CCGAAGATGA	CAGCAAGGAT	20640
CGGCTAAATA	CCGCGTGTTT	TCATCTAGAA	GTGGGAACCT	ACAGATGACA	GTTCTTGAT	20700
GGGCAGAACG	AGGGGGACCG	GGGACGCGGA	AGTCTGCTTG	AGGGAGGAGG	GGTGAAGGA	20760
GAGACAGCTT	CAGGAAGAAA	ACAAAACACG	AATACTGTCT	GACACAGCAC	TGACTACCCG	20820
GGTGATGAAA	TCATCTGCAC	ACTGAACACC	CCCGTCACAA	GTTTACCTAT	GTCACAATCT	20880
TGCACATGTA	TCGCTTGAAC	GACAAATAAA	AGTTAGGGGG	GAGAAGAGAG	GAGAGAGAGA	20940
GAGAGAGAGA	GACAGAGAGA	GACAGAGAGA	GAGAGAGAGG	AGGGAGAGAG	GAAGAGAGAA	21000
CACCACCTCC	TTGACCTGAG	TCAGGGGGTT	TCTGGCCTTT	TGGGAGAACG	TTGAGCGACA	21060
ATGCAGTATT	TGGGCCCGTT	CTTTTCTTTT	CTTCTTCTTT	TCTTTCTTTT	TTTTTGGACT	21120
GAGTCTCTCT	CGCTCTGTCA	CCCAGGCTGC	GGTCGCGGTG	GCGCTCTCTC	GGCTCACTGA	21180
AACCTCTGCT	TCCCGGGTTC	CAGTGATTCT	TCTTCGGTAG	CTGGGATTAC	AGGCGCACAC	21240
CATGACGGCG	GGCTCATATT	CCTATTTTCA	GTAGAGACGG	GGTTTCTCCA	CGTTGGCCAC	21300
GCTGGTCTCG	AATCTCTGAC	CTCAAATGAT	CCGCCTCTCT	GGGCCTCCCA	AAGTCTTGGG	21360
AACGACAGCG	CTGAGCCGCC	GGGATTTTCAG	CCTTTAAAG	CGCGGCCCTG	CCACCTTTTC	21420
CTGTGGCCCT	TACGCTCAGA	ATGACGTGTC	CTCTCTGCGG	TAGGTTGACT	CCTTGAGTCC	21480
CCTAGGCCAT	TGCACTGTAG	CCTGGGCAGC	AAGAGCCAAA	CTCCGNNCCC	CCACCTCCTC	21540
GCGCACATAA	TAACCTAATA	ACAACTAATC	TAACCTAATA	AACTAATAAA	CTAATAAAAA	21600
TCTCTACACG	TCACCCATAA	GTGTGTGTTC	CCGTGAGAGT	GATTTCTAAG	AAATGGTACT	21660
GTCACTGAA	CGCAGTGGCT	CACGTCTGTC	ATCCCGAGGT	CAGGAGTTCT	AGACAGACCC	21720
GGCCAACGTG	GTGAAACCCC	GTCTCTACTG	AAAATACGAA	ATGGAGTCAG	GCGCCGTGGG	21780
GCAGGCACCT	GTAACCCAG	CTACTCGGGA	GGCTGGGGTG	GAAGAATTGC	TTGAACCTGG	21840
CAGGCGGAGG	CTGCAGTGAC	CCAAGATCGC	ACCACTGCAC	TACAGCCTGG	GCGACAGAGT	21900
GAGACCCGGT	CTCCAGATAA	ATACGTACAT	AAATAAATAC	ACACATACAT	ACATACATAC	21960
ATACATACAT	ACATACATAC	ATCCATGCAT	ACAGATATAC	AAGAAAGAAA	AAAAGAAAAG	22020
AAAAGAAAAG	GAAAATGAAA	GAAAAGGCAC	TGTATTGCTA	CTGGGCTAGG	GCCTTCTCTC	22080
TGTCTGTTTC	TCTCTGTTTC	TCTCTGTCTT	TCTCTGTCTG	TCTCTTTCTC	TCTCTGTCTG	22140
TCTCTTTCTT	TCTCTCTGTC	TCTGTCTCTG	TCTTTGTCTC	TCTCTCTCCC	TCTCTGCCTG	22200
TCTCACTGTG	TCTGTCTTCT	GTCTTACTCT	CTTTCTCTCC	CCGTCTGTCT	CTCTCTCTCT	22260
CTCTCCCTCC	CTGTTTGTTT	CTCTCTCTCC	CTCCCTGTCT	GTTTCTCTCT	CTCTCTTTCT	22320
GTCTGTTTCT	GTCTCTCTCT	GTCTGTCTAT	GTCTTTCTCT	GTCTGTCTCT	TTCTCTGTCT	22380
GTCTGCCTCT	CTCTTTCTTT	TTCTGTGTCT	CTCTGTCTGG	CTCTCTCTCT	CTGTCTGTCT	22440
TCTGTCTCTT	CTCTCTCTCT	CTCTGTGCCT	ATCTGTGTCT	TTACTCTCTT	TCTCTGCCTG	22500
TCTGTCTGTC	TCTCCCTCCC	TTTCTGTTTC	TCTCTCTCTC	TCTCTCTCTC	TCCCTCTCTC	22560
CCTGTCTGTT	TCTCTCCGTC	TCTCTCTCTT	TCTGTCTGTT	TCTCACTGTC	TCTCTCTGTC	22620
CATCTCTCTC	TCTCTCTGTC	TGTCTCTTTC	GTTCTCTCTG	TCTGTCTGTC	TCTCTCTCTC	22680
TCTCTCTCTC	TCTCTCTCTC	TCCCTGTCTG	TCTGTTTCTC	TCTATCTCTC	GCTGTCCATC	22740
TCTGTCTTTT	TATGTCTGTC	TCTTTCTCTG	TCAGTCTGTC	AGACACCCCC	GTGCCGGGTA	22800
GGGCCCTGCC	CCTTCCACGA	AAGTGAGAAG	CGCGTGCTTC	GGTGCTTAGA	GAGGCCGAGA	22860
GGAATCTAGA	CAGGCGGGCC	TTGCTGGGCT	TCCCCACTCG	GTGTATGATT	TCGGGAGGTC	22920
GAGGCCGGGT	CCCCGCTTGG	ATGCGAGGGG	CATTTTTCAGA	CTTTTCTCTC	GGTCACGTGT	22980
GGCGTCCGTA	CTTCTCCTAT	TTCCCCGATA	AGCTCCTCGA	CTTCAACATA	AACGGCGTCC	23040
TAAGGGTCGA	TTTAGTGTCA	TGCCTCTTTC	ACCGCCACCA	CCGAAGATGA	AAGCAAAGAT	23100
CGGCTAAATA	CCGCGTGTTT	TCATCTAGAA	GTGGGAACCT	ACAGATGACA	GTTCTTGAT	23160
GGGCAGAACG	AGGGGGACCG	GGNACGCGGA	AGCCTGCTTG	AGGGRGGAGG	GGYGAAGGA	23220

-218-

GAGACAGCTT	CAGGAAGAAA	ACAAAACACG	AATACTGTCTG	GACACAGCAC	TGACTACCCG	23280
GGTGATGAAA	TCATCTGCAC	ACTGAACACC	CCCGTCACAA	GTTTACCTAT	GTCACAGTCT	23340
TGCTCATGTA	TGCTTGAACG	ACAAATAAAA	GTTTCGGGGG	GAGAAGAGAG	GAGAGAGAGA	23400
GAGAGACGGG	GAGAGAGGGG	GGAGAGGGGG	GGGGAGAGAG	AGAGAGAGAG	AGAGAGAGAG	23460
AGAGAGAGAG	AGAAAGAGAA	GTAAAACCAA	CCACCACCTC	CTTGACCTGA	GTCAGGGGGT	23520
TTCTGGCCTT	TTGGGAGAAC	GTTCAGCGAC	AATGCAGTAT	TTGGGCCCGT	TCTTTTTTTC	23580
TTCTTCTTCT	TTTCTTTCTT	TTTTTTTGGG	CTGAGTCTCT	CTCGCTCTGT	CACCCAGGCT	23640
GCGGTGCGGT	GGCGCTCTCT	CGGCTCACTG	AAACCTCTGC	TTCCCGGGT	CCAGTGATTCT	23700
TTCTTCGGTA	GCTGGGATTA	CAGGTGCGCA	CCATGACGGC	CGGCTCATCG	TTCTATTTT	23760
AGTAGAGACG	GGGTTTCTCC	ACGTTGGCCA	CGCTGGTCTC	GAACCTCTGA	CCACAAATGA	23820
TCCACCTTCC	TGGGCCTCCC	AAAGTGCTGG	AAACGACAGG	CCTGAGCCGC	CGGGATTTC	23880
GCCTTTAAAA	GCGCGCGGCC	CTGCCACCTT	TCGCTGCGGC	CCTTACGCTC	AGAATGACGT	23940
GTCCTCTCTG	CCATAGGTTG	ACTCCTTGAG	TCCCCTAGGC	CATTGCACTG	TAGCCTGGGC	24000
AGCAAGAGCC	AAACTCCGTC	CCCCACCTC	CCCGCGCACA	TAATAACTAA	CTAACTAACT	24060
AACCTAATAA	AATCTCTACA	CGTCACCCAT	AAGTGTGTGT	TCCCGTGAGG	AGTGATTCT	24120
AAGAAATGGT	ACTGTACACT	GAACGCAGGC	TTCACGTCTG	TCATCCCGAG	GTCAGGAGTT	24180
CGAGACCAGC	CCGGCCCCACG	TGGTGAAACC	CCCGTCTCTA	CTGAAAATAC	GAAATGGAGT	24240
CAGGCGCCGT	GGGGCAGGCA	CCTGTAACCC	CAGCTACTCG	GGAGGCTGGG	GTGGAAGAAT	24300
TGCTTGAACC	TGGCAGGCGG	AGGCTGCAGT	GACCCAAGAT	CGCACCCTG	CACTACAGCC	24360
TGGGCGACAC	AGTGAGACCC	GGTCTCCAGA	TAAATACGTA	CATAAATAAA	TACACACATA	24420
CATACATACA	TACATACAAC	ATACATACAT	ACAGATATAC	AAAAGAAAGAA	AAAAGAAAG	24480
AAAAGAAAGA	GAAAATGAAA	GAAAAGGCAC	TGTATTGCTA	CTGGGCTAGG	GCCTTCTCTC	24540
TGTCTGTTTC	TCTCTGTTTC	TCTCTGTCTT	TCTCTCTGTG	TCTCTTTCTC	TGTCTGTCTG	24600
TCTGTCTGTC	TGTCTGTCTC	TTTCTTTCTT	TCTGTCTCTG	TCTTTGTCCC	TCTCTCTCCC	24660
TCTCTGCCCT	GTCTCACTGT	GTCTGTCTTC	TATCTTACTC	TCTTTCTCTC	CCCGTCTGTG	24720
TCTCTCTCAC	TCCCTCCCTG	TCTGTTTCTC	TCTCTCTCTC	TTTCTGTCTG	TTTCTGTCTC	24780
TCTCTGTCTG	CCTCTCTCTT	TCTCTATCTG	TCTCTTTCTC	TGTCTGTCTG	CCCTCTCTCT	24840
TCTTTTTCTG	TGTCTCTCTG	TCTGTCTCTC	TCTCTCTCTG	TGCCTATCTT	CTGTCTTACT	24900
CTCTTTCTCT	GCCTGTCTGT	CTGTCTCTCT	CTGTCTCTCC	CTCCCTTTCT	GCTTCTCTCT	24960
CTCTCTCTCT	CTCTNNNCCC	TCCCTGTCTG	TTTCTCTCTG	TCTCCCTCTC	TTTCTGTCTG	25020
TTTCTCACTG	TCTCTCTCTG	TCTGTCTGTT	TCATTCTCTC	TGTCTCTGTC	TCTGTCTCTC	25080
TCTCTCTCTG	TCTCTCCCTG	TCTGTGTGTA	TCTTTTGTCT	TACTCTCCTT	CTCTGCCTGT	25140
CCGTCTGTCT	GTCTGTCTCT	TCTCTCCCT	GTCCTCTCT	CTTTCTGTCT	GTTTCTCTCT	25200
CTCTCTCTCT	CTCTCTCTCT	CTGTCTCTGT	CTTTCTCTGT	CTGTCCCTTT	CTCTGTCTGT	25260
CTGCCTCTCT	CTTTCTCTTT	CTGTGTCTCT	CTGTCTCTCT	CTCTGTGCCT	ATCTTCTGTG	25320
TACTCTCTTT	TCTCTGCCTG	TCTATCTGTC	TGTCTCTCTC	TGTCTCTCTC	CCTGCCTTTT	25380
TGTTTCTCTC	TCTCTCCCTC	TCTCGCTCTC	TCTGTCTTTT	TCTCTTTCTC	TCTGTTTCTC	25440
TGTCTCTCTC	TGTCTGCTCT	TGTCTTTTTC	TGTCTGTCTG	TCTCTCTCTT	TCTTTCTGTG	25500
GTCTGTCTCT	GTCTCTGTCT	CTGTCTCTCT	CTCTCTCTCT	CTCCTTGTCT	CTCTCACTGT	25560
GTCTGTCTCT	TGTCTTACTC	TCCTTCTCTG	CCTGTCCATC	TGTCTGTCTG	TCTCTCTCTC	25620
TCTCTCCCTA	CCTTTCTGTT	TCTCTCTCGC	TAGCTCTCTC	TCTCTCTGCC	TGTTTCTCTC	25680
TTTCTCTCTC	TGTCTTTCTC	TGTCTGTCTC	TTTCTCTGTC	TGTCTGTCTC	TTTCTCTCTG	25740
TCTCTGTCTC	TGTCTCTCTC	TCTCTCTCTC	TCTCTCTCTC	TGCCTCTCTC	ACTGTGTCTG	25800
TCTTCTGTCT	TATTCTCTTT	CTCTCTCTGT	CTCTCTCTCT	CTCTCCTTTA	CTGTCTGTTT	25860
CTCTCTCTCT	CTCTCTCTTT	CTGCCTGTTT	CTCTCTGTCT	GTCTCTGTCT	TTCTCTGTCT	25920
GTCTGCCTCT	CTCTTTCTTT	TTCTGCGTCT	CTCTGTCTCT	CTCTCTCTCT	CTCTGTTCTT	25980
ATCTTCTGTC	TACTCTGTT	TCCTTGCCTG	CCTGCCTGTC	TGTGTGTCTG	TCTCTCTCTC	26040
TCTCTCTCTC	TCTCTCTCCC	TCCCTTTCTC	TTTCTCTGTC	TCTCTCTCTC	TTTCTGGGTG	26100
TTTCTCTCTG	TCTCTCTGTC	CATCTCTGTC	TTTCTATGTC	TGTCTCTCTC	TTTCTCTCTG	26160
TCTCTGTCTC	TGCCTCTCTC	TCTCTCTCTC	TCTCTCTCTC	TCTGTCTGTC	TCTCTCACTG	26220
TGTGTGTCTG	TCTTCTGTCT	TACTCTCCTT	CTCTGCCTGT	CCGTCTGTCT	GTCTGTCTCT	26280
CCCTCTCTCT	CCCTCCCTTT	CTGTTTCTCT	CTCTCTCTCT	TTCTGTCTGT	TTCTCTCTTT	26340
CTCTCTCTGT	CTGTCTCTTT	CTCTGTCTGT	CTGTCTCTCT	CTTTCTTTTT	CTCTGTCTCT	26400
CTGTCTCTCT	CTGTGTCTGT	CTCTCTGTCT	GTGCCTATCT	TCTGTCTTAC	TCTCTTTCTC	26460
TGGCTGTCTG	CCTGTCTCTC	TCTCTCTCTC	TGTCTGTCTC	CGTCCCTCTC	TCCCTGTCTG	26520
TCTGTTTCTC	TCTCTGCCTC	TCTCTCTCTC	TGTCTGTCTC	TTTCTCTGTC	TGTCTGTCTC	26580
TCTCTTTCTT	TTTCTCTGTC	TCTCTGTCTC	TCTCTGTGTC	TGTCTCTCTT	TGTGTGCTTA	26640
TCTTCTGTCT	TACTCTCTTT	CTCTGGCTGT	CTGCCTGTCT	CTCTCTCTCT	GCCTGTCTCC	26700
GTCCCTCCCT	CCCTGTCTGT	CTGTTTCTCT	CTCTGTCTCT	GTCTCTCTGT	CCATCTCTGT	26760
CTGTCTCTTT	CTCTTTCTCT	CTCTCTGTCT	CTGTCTCTCT	CTCTCTCTGC	CTGTCTCTCT	26820
CACTGTGTCT	GTCTTCTGTG	TACTCTCTTT	TCTCTTGCTT	GCCTCTCTGT	CTGTCTGTCT	26880

-219-

CTCTCCCTCC	ATGTCTCTCT	CTCTCTCTCA	CTCACTCTCT	CTCCGTCTCT	CTCTCTTTCT	26940
GTCTGTTTTCT	CTCTCTGTCT	GTCTCTCTCC	CTCCATGTCT	CTCTCTCTCT	CTCTCACTCA	27000
CTCTCTCTCC	GTCTCTCTCT	CTCTTTCTGT	CTGTTTCTCT	CTCTGTCTGT	CTCTCTCCCT	27060
CCATGTCTCT	CTCTCTCCCT	CTCACTCACT	CTCTCTCCGT	CTCTCTCTCT	CTTTCTGTCT	27120
GTTTCTTTGT	CTGTCTGTCT	GTCTGTCTGT	CTGTCTCTCT	CTCTCTCTCT	CTCTCTCTCT	27180
CTCTCTGTTT	GTCTTTCTCC	CTCCCTGTCT	GTCTGTCTGT	CTCTCTCTCT	CTGTCTCTGT	27240
CTCTGTCTCT	CTCTCTTTCT	CTTTCTGTCT	GTTTCTCTCT	ATCTCTCGCT	GTCCATCTCT	27300
GTCTTTCTAT	GTCTGTCTCT	TTCTCTGTCA	GTCTGTCTGT	CACACCCGTG	CCGGTAGGGC	27360
CCTGCCCTTC	CACGAGAGTG	AGAAGCGCGT	GCTTCGGTGC	TTAGAGAGGC	CGAGAGGAAT	27420
CTAGACAGGC	GGGCCTTGCT	GGGCTTCCCC	ACTCGGTGTA	CGATTTCTGG	AGGTCGAGGC	27480
CGGGTCCCCG	CTTGGATGCG	AGGGGCATTT	TCAGACTTTT	CTCTCGGTCA	CGTGTGGCGT	27540
CCGTACTTCT	CTATTTCCCT	CGATAAGTCT	CCTCGACTTC	AACATAAACT	GTTAAGGCCG	27600
GACGCCAACA	CGGCGAAACC	CCGTCTCTAC	TAAAAATACA	AAGCTGAGTC	GGGAGCGGTG	27660
GGGCAGGCCC	TGTAATGCCA	GCTCCTCGGG	AGGCTGAGGC	GGGAGAATCG	CTTGAACCAG	27720
GGAAGCGGAG	GCTGCAGGGA	GCCGAGATCG	CGCCACTGCA	CTACGGCCCA	GGCTGTAGAG	27780
TGAGTGAGAC	TCGGTCTCTA	AATAAATACG	GAAATTAATT	AATTCATTAA	TTCTTTTCCC	27840
TGCTGACGGA	CATTTGCAGG	CAGGCATCGG	TTGTCTTCTG	GCATCACCTA	GCGGCCACTG	27900
TTATTGAAAG	TCGACGTTGA	CACGGAGGGA	GGTCTCGCCG	ACTTCACCGA	CCCTAGGGCA	27960
ACGGGTTTCT	CTCTCTCCCT	TCTGGAGGCC	CCTCCCTCTC	TCCCTCGTTG	CCTAGGGAAC	28020
CTCGCCTAGG	GAACCTCCGC	CCTGGGGGCC	CTATTGTTCT	TTGATCGGCG	CTTTACTTTT	28080
CTTTGTGTTT	TGGCGCCTAG	ACTCTTCTAC	TTGGGCTTTG	GGAAGGGTCA	GTTTAATTTT	28140
CAAGTTGCCC	CCCGGCTCCC	CCCACTACCC	ACGTCCCTTC	ACCTTAATTT	AGTGAGNCGG	28200
TTAGGTGGGT	TTCCCCCAA	CCGCCCCCCC	CCCCCGCCTC	CCCAACACCC	TGCTTGGAAC	28260
CCTTCCAGAG	CCACCCCGGT	GTGCCTCCGT	CTTCTCTCCC	CTTCCCCCAC	CCCTTGCCGG	28320
CGATCTCAAT	CTTGCCAGGC	TGACATTTGC	ATCGGTGGGC	GTCAGGCCTC	ACTCGGGGGC	28380
CACCGTTTTT	GAAGATGGGG	GCGGCACGGT	CCCACTTCCC	CGGAGGCAGC	TTGGGCCGAT	28440
GGCATAGCCC	CTTGACCCGC	GTGGGCAAGC	GGGCGGGTCT	GCAGTTGTGA	GGCTTTTCCC	28500
CCCGCTGCTT	CCCGCTCAGG	CCTCCCTCCC	TAGGAAAGCT	TCACCCTGGC	TGGGTCTCGG	28560
TCACCTTTTA	TCACGATGTT	TTAGTTTCTC	CGCCCTCCGG	CCAGCAGAGT	TTACAATGTC	28620
GAAGGCGGCC	ACGGCTCTAG	TCTGGGCCTT	CTCAGACTTA	GCCCAAATA	GAAACGCTTT	28680
CTGAAACATA	ATACTTTTNC	TCACTTAAGA	TTTCCAGGTA	CGGCGCCTTG	GCCCGTGTTC	28740
GTTGGCTTGT	TTTGTTCGT	TCTGTTTTGT	TTTGTTCGTG	TTTTTCCTTT	CTCGTATGTC	28800
TTTCTTTTCA	GGTGAAGTAG	AAATCCCCAG	TTTTCCAGGA	GACGTCTATT	TTCCCCAAGA	28860
CACGTTAGCT	GCCGTTTTTT	CCTGTTGTGA	ACTAGCGCTT	TTGTGACTCT	CTCAACGCTG	28920
CAGTGAGAGC	CGGTTGATGT	TTACNATCCT	TCATCATGAC	ATCTTATTTT	CTAGAAATCC	28980
GTAGGCGAAT	GCTGTGCTG	CTCTGTGTTG	GGTTGTGTTT	GTTGTGTTG	CTAGCTGTTG	29040
TGTTGTCTCT	TGCGTTGTTG	TTGTGTTGTT	CGTTGTTTTC	AAAGTATACC	CCGGCCACCG	29100
TTTATGGGAT	CAAAAGCATT	ATAAAATATG	TGTGATTATT	TCTTGAGCAC	GCCCTTCTCT	29160
CCCTCTCTCT	TGTCTCTCTG	TCTGTCTCTG	TCTCTCTCTT	TCTGTCTCTG	TCTTCTCTCT	29220
CTCTCTCTCT	CTGTGTCTCT	CTCTCTCTGC	CTGTCTGTTT	CTCTCTCTCT	GCCTCTCTCT	29280
CTCTCTCTCT	CTCTGCCTGT	CTCTCTCACT	GTGTCTGTCT	TCTGTCTTAC	TCCCTTTCTC	29340
TCTCTGTCTG	TCGGTCTCTC	TCTCTCTCTC	TCCCTGTCTG	TATGTTTCTG	TCTGTCTCTG	29400
TCTCTCTCTC	TCTTTCTGTT	TCTCTCTCTC	CGTCTCTGTC	TTTCTCTGAC	TGTCTCTCTC	29460
TTTCTCTCTC	TCTGTCTCTC	TCTGCCTGTC	TCTCTCACTC	TGTCTTCTGT	CTTATCTCTC	29520
TCTCTGCCTG	CCTGTCTCTC	TCACTCTCTC	TCTCTGTGTG	TCTCTCTCTC	TCTTTCTGTT	29580
TCTCTCTGTC	TCTCTGTCCG	TCTCTGTCTT	TCTCTGTCTG	TCTCTTTGTC	TGTCTGTCTT	29640
TGTCTTTCCCT	TCTCTCTGTC	TCTGTCTCTC	TCACTGTGTC	TGTCTTCTGT	CTTAGTCTCT	29700
CTCTCTCTCT	CTCCCTGTCT	GTCTGTCTCT	CTCTCTCTCT	CCCCCTGTCT	GTTTCTCTCT	29760
CTCTCTCTCT	CTCTCTCTCT	CTCTGTCTTT	GTCTTTCTTT	CTGTCTCTGT	CTCTCTCTCT	29820
CTCTCTGTGT	GTCTGTCTTC	TGTCTTACTG	TCTTTCTCTG	CCTGTCTGTC	TGTCTGTCTC	29880
TCTCTGTCTG	TCTCTCTCTC	TCTCTCCCCC	TGTGCGCTGT	TTCTCTGTCT	CTGTCTGTGT	29940
CTCTCTTTCT	GTCTGTTTCT	CTCTGTCTGT	CTTTCTCTCT	CTGTCTCTTT	CTCTCTGTCT	30000
CTCTGTCTGT	CTCTGTCTCT	CTCTCTGTCT	CTCTCTCTCT	GTGGGGGTGT	GTGTGTGTGT	30060
GTGTATGTGT	GTGTGTGTGT	GTGTGTGTGT	GTGCCTCTCT	TCTTACTCTC	TTTCTCTGCC	30120
TGTCTGTCTG	CCTGTCTGTT	TGTCTCTCTC	TCTCTCCCTG	TCTCTCTCCC	TTCTGTCTCT	30180
TTTCTCTCTC	TTTCTGTTTC	TCTCTGTCTC	TGTCCATCTC	TGTCTTTCTC	CGTCTGTCTC	30240
TTTATCTGTC	TCTCTCCGTC	TGTCTCTTTA	TCTGTCTCTC	TCTCTCTTTT	TGTCTTTCTC	30300
TCTCTGTGTA	TCGTTGTCTC	TCTCTGTCTG	TCTCTGTCTC	TGTCTCTCTG	TCTCTCTCTC	30360
TCTCTCTCTC	TCTCTGTCTG	TCTGTCCGTC	TGTCTGTCTC	GGTCTCTGCG	TCTCGCTATC	30420
TCCCGCCCTC	TCTTTTTTTG	CAAAAGAAGC	TCAAGTACAT	CTAATCTAAT	CCCTTACCAA	30480
GGCCTGAATT	CTTCACTTCT	GACATCCCAG	ATTTGATCTC	CCTACAGAAT	GCTGTACAGA	30540

-220-

ACTGGCGAGT	TGATTTCTGG	ACTTGGATAC	CTCATAGAAA	CTACATATGA	ATAAAGATCC	30600
AATCCTAAAA	TCTGGGGTGG	CTTCTCCCTC	GACTGTCTCG	AAAAATCGTA	CCTCTGTTCC	30660
CCTAGGATGC	CGGAAGAGTT	TTCTCAATGT	GCATCTGCCC	GTGTCCTAAG	TGATCTGTGA	30720
CCGAGCCCTG	TCCGTCCTGT	CTCAAATATG	TACGTGCAAA	CACTTCTCTC	CATTTCCACA	30780
ACTACCCACG	GCCCCCTGTG	GAACCACTGG	CTCTTTGAAA	AAAATCCCAG	AAGTGGTTTT	30840
GGCTTTTTGG	CTAGGAGGCC	TAAGCCTGCT	GAGAACTTTC	CTGCCCAGGA	TCCTCGGGAC	30900
CATGCTTGCT	AGCGCTGGAT	GAGTCTCTGG	AAGGACGCAC	GGGACTCCGC	AAAGCTGACC	30960
TGTCCCAACG	AGGTCAAATG	GATACCTCTG	CATTGGCCCC	AGGCCTCCGA	AGTACATCAC	31020
CGTCACCAAC	CGTCACCGTC	AGCATCCTTG	TGAGCCTGCC	CAAGGCCCCG	CCTCCGGGGA	31080
GACTCTTGGG	AGCCCGGCCT	TCGTCGGCTA	AAGTCCAAAG	GGATGGTGAC	TTCCACCCAC	31140
AAGGTCCAC	TGAACGGCGA	AGATGTGGAG	CGTAGGTCAG	AGAGGGGACC	AGGAGGGGAG	31200
ACGTCCCGAC	AGGCGACGAG	TTCCCAAGGC	TCTGGCCACC	CCACCCACGC	CCCACGCCCC	31260
ACGTCCCGGG	CACCCGCGGG	ACACCGCCGC	TTTATCCCCCT	CCTCTGTCCA	CAGCCGGCCC	31320
CACCCACCA	CGCAACCCAC	GCACACACGC	TGGAGGTTCC	AAAACCACAC	GGTGTGACTA	31380
GAGCCTGACG	GAGCGAGAGC	CCATTTACAG	AGGTGGGAGG	GGTGGGGGTG	GGGTGGGTTG	31440
GGGGTTGTGG	GGTCTGTGGG	GAGCCCGATT	CTCCCTCTTG	GGTGGCTACA	GGCTAGAAAT	31500
GAATATCGCT	TCTTGGGGGG	AGGGGCTTCC	TTAGGCCATC	ACCGCTTGCG	GGACTACCTC	31560
TCAAACCCTC	CCTTGAGGCC	ACAAAATAGA	TTCCACCCCA	CCCATCGACG	TTTCCCCCGG	31620
GTGCTGGATG	TATCCTGTCA	AGAGACCTGA	GCCTGACACC	GTCTGAATTAA	ACACCTTGAC	31680
TGGCTTTGTG	TGTTTGTGTT	TTTCTGAGAT	GGAGTCTTGC	TCTGTCCCCC	AGGCTGGAGT	31740
GCAGTGGCGT	GATCTCAGCT	CACTGGAACC	TCTGCCTCCT	GGGTTCAAGT	GATTCTCCTG	31800
TCTCAGCGCG	ACCATGGCCG	GCTCATTTTT	TTTTTTTTTT	TTTTTGGTAG	ACACGGGGTT	31860
TCACCCTCTT	TCATTGGTTT	TCACTGGAGA	TTCTAGATTG	GAGCCACACC	TCATTCCGTG	31920
CCACAGAGAG	ACTTCTTTTT	TTTTTTTTTT	TTTTTAAGCG	CAACGCAACA	TGTCTGCCTT	31980
ATTTGAGTGG	CTTCCTATAT	CATTATAATT	GTGTTATAGA	TGAAGAAACG	GTATTAAACA	32040
CTGTGCTAAT	GATAGTGAAA	GTGAAGACAA	AAGAAAGGCT	ATCTATTTTG	TGGTTAGAAT	32100
AAAGTTGCTC	AGTATTTAGA	AGCTACCTAA	ATACGTCAGC	ATTTACACTC	TTCTAGTAA	32160
AAGCTGGCCG	ATCTGAATAA	TCCTCCTTTA	AACAACACAA	ATTTTTGATA	GGGTAAAGAT	32220
TTTTTTAAGA	ATGCGACTCC	TGCAAAATAG	CTGAACAGAC	GATACACATT	TAAAAAATA	32280
ACAACACAAG	GATCAACCAG	ACTTGGGAAA	AAATCGAAAA	CCACACAAGT	CTTATGAAGA	32340
ACTGAGTTCT	TAAAATAGGA	CGGAGAACGT	AGCTATCGGA	AGAGAAGGCA	GTATTGGCAA	32400
GTTGATTGTT	ACGTTGGTCA	GCAGTAGCTG	GCACTATCCT	TTTGGCCATC	TTTCGGGCAA	32460
TGTAACACTA	ACAGCAAAAT	GAGATATGAT	CCATTAAACA	ACATATTCGC	AAATCAAAAA	32520
GTGTTTCAGT	AATATAATGC	TTTCAAGTTA	GAAGCAAAATC	AAATGATAGA	ACTCCACTGC	32580
TGTAATAAGT	CACCCCAAAG	ATCACCGTAT	CTGACAAAAT	AACTACCACA	GGGTTATGAC	32640
TTCAGAATCA	TACTTTCTTC	TTGATATTTA	CTTATGTATT	TATTTTTTTT	AATTTATTTT	32700
TCTTGAGACG	CGTCTCGCTC	TGTCGCCCAG	GCTGGAGTGC	GATGGTGTGA	TCTCGGCTCA	32760
CTGCAACCGC	CACCTCCCTG	GGTTCAAGCG	ATTCTCCTGC	CTCAGCCTCC	CGAGTAGCTG	32820
GGACTACAGG	TGCCCCCAG	CACGCCACGC	TAATCTTTAT	ACTTTTAATA	GAGACGGGGT	32880
TTCACCGTGT	GGCCCCGGAT	GGTCTCGATC	TCTTGACCTC	GTGACCCGCC	CGCCTCGGCC	32940
TCCCAAAGTG	CTGGGATGAC	AGGCGTGAGC	CACTGAGCCC	GGCCTTCTCT	TGACGTTTAA	33000
ACTATGAAGT	CAGTCCAGAG	AAACGCAATA	AATGTCAACG	GTGAGGATGG	TGTTGAGGCA	33060
GAAGTAGGAC	CACACTTTTT	CCTATCTTAT	TCAGTTGATA	ACAATATGAC	CTAGGTAGTA	33120
ATTTCTTATG	TGCCTACTTA	TACACGAGTA	CAAAAGAGTA	AAACAGAGAG	ACTGCTAAAT	33180
TAAAGGGTAC	GTGAAGTTCT	TCATAGTAAC	TCCGTAAACT	GGAACACTGT	CAAAAAGCAG	33240
CAGCTAGTGA	ATTGTTTCCA	TGTATTTTTT	TATTATCCAA	TAAGTGAAC	ATGCTATTCC	33300
TTTCCAGTCT	CCCAAGCACT	TCTTGTCCTC	ATCACCACTT	CGGTGCTCGA	AGAAAAAGTA	33360
AGCAAATCAA	GGAACACAAG	CTAAAGAAAC	ACACACACAA	ACCAAAGACA	ACTACAGCGT	33420
CTGCAAAAGT	TTGCTAGAAG	ACTGAAACTG	TTGAGTATAA	GGATCTGGTA	TTCTACGATC	33480
ATGAGTTCAC	TTTCAAGGTT	GTTCAAGACA	TACGTTTCGT	AAGGAAACAT	CTTAGTTAGA	33540
AGTTATTCAG	CAGTAGGTAC	CATCCCTAAG	TATTTTTCAC	CAAATCCGTG	ACAATAAAGA	33600
GCTATCTAAC	CAGAAAAATT	AGCGAGTACG	GGCACCATCC	ATAGGGCTTT	GTCTTTACGC	33660
TTCATTAGCA	CTTACCATGC	CTTACAATGT	CTAGGATTGA	CCCTGATAGC	ATTTTCAAAA	33720
CAAGCTAATG	CTTTGTCCAG	TTCTTCAGTG	AAGACAACCTC	ACGCCCTAAT	GCGCTATAGG	33780
CATAAGCATC	ATTTGGATCC	ACTTCGAGAG	TTCTCTGGAA	GAATTGAATC	GCAATATCGT	33840
GTTCCCGTTT	GCAGACCGAA	ACAGTTTCCC	TGCAGCACAC	CAGGCCCTCTG	GCTGGCGAAT	33900
TTTATCTCAT	GTCTGTGAAG	TCTTTGGACA	GAACTGAAAAG	AGCAACCTCT	TTCGGAGGAT	33960
GCCAAAGTGT	TGTAGAGTAG	ATCTCCATGC	CTTCGACTCT	GTAATTCTCA	ATCCTCCTAA	34020
CCTCTGAGAA	TTGTCTTTCA	GCTTGCGTGG	ACTCTGAAAG	TTTACAATAG	GCCNTTTCCG	34080
ATTTGGCACA	GTACCCAACC	GGTATTGCAG	TGGTGAGAAG	CTAGATGGCT	CAAGATGCTG	34140
ATAGCTTCTT	TGCCGTGGTA	AGAACACAAA	GCTAAATAAC	CTTCCCCCTT	TTCACGAAGA	34200

AGGCTCATCA	AGCCTTCCGC	TGCTGCTTTT	TGTAGATTAA	AAGCCTGAAT	CTGAGGCGCG	34260
ATTGCGGCTA	TTTTCCCTTC	TGAAATGACG	GAAGAGTCCA	ATTTTGTCAC	TTCCAGGCTA	34320
TCACCTTATGT	TCGGTGAGT	TATTGCTCCT	TTATTAGTTT	TACTTTTGGT	TCTTCTGTTT	34380
GGGATTTTAG	GTGGAAACTT	CATTTTTAAT	TTTCTCCTAA	TTCTCCTCGG	TTGTGGAGCT	34440
GTCAC TAGTC	AAGAGTCGTG	AATTTCTTCG	AGGNCGGTGC	ATTTGGGGGA	GATGCCATAG	34500
TGGGGCTCAA	TACCTGAGGT	GTTGCCCTTG	TCGGCGGACC	AGAACTTTGT	GTTTTTGCAA	34560
GGACTGGAGT	TACCTTTCGG	CTCTTTCCCC	TCTGCGAGAA	GACAGACGGT	GTTCCGGTTT	34620
GGCCGATTCT	GGCAACAGGC	TTTTCTGAAG	GGGCTCCGGT	GGATGGCAG	TCAGTGACAG	34680
ACGGTGTCTC	ATACCAGTGC	AGTTTTGTCA	ATAGGGTCCG	TCTCCGGGAC	TTGGGGTTTC	34740
TAATGGCAAA	ATGCCAACAC	TTGGGGTTAA	TGGACTAACA	GCTGCTGGTC	CTCCTAATAA	34800
ACTTCGACCA	GTTTTTGTTT	TATGTTGAAC	CTGTTTAGAT	CATATGGAAG	TTCTGTTC	34860
CAGTGGGACA	GTATCAGGTG	AAAGGACAGC	TGAATCGATA	GAAGACACTG	GGGAGTCTGT	34920
ATTCAAGGAG	TACTTTGAAT	TGGAAGATTG	TAAATTCCAT	CCGTTTCATT	CGACGGTGTG	34980
CTGGGGTGTT	TCCGTAAGAA	CGGTCTCGGG	CTGTCTGTGA	CATAAACTAG	GACGAGTCC	35040
AAGTGTGTGT	GCGCAACACT	TGGACAGGCA	GTTGCTAAAG	CTCTCTAGAG	AGGTGAATCA	35100
AAATGTTTGG	TCAGGATCTG	GCTTTTCCCC	CCTATTTTAC	ATCATGATTG	AAAGGGACAC	35160
CAGAGGAAAG	GATTTCAACG	AAGGCTCTTT	TGGTCACATT	CTGATCCTTT	GGTAAAGCCGA	35220
TCTGTCTTGC	AATATACATG	TCCCGACGAT	GGAAGGGGAA	AGCGAGCTGA	ATCACCAAAC	35280
TCAGGAACGA	TAATATCATC	GTGGCTTTTC	TGCTTATGAA	ACACTCCACC	CGATAAGATT	35340
TGATCCCCCT	CTGCAAGCTT	GCTGAGATCA	ACACAACATT	TCGCAAGCAG	GCATTTGCAT	35400
TGCGGGGTAG	TACAACGTG	TCCTTTCAAG	AGTCTATATG	TTTTATAGGC	CTTTCCTGAG	35460
CGGTAAGAAC	AGGTGCGCAG	TAAGAACAAG	GCTTCTTCTG	AGTGTACTTC	TGCATAAAGG	35520
CGTTCTGCGG	GGGAAACCGC	ATCTCGGTAG	GCATAGTGGT	TTAGTGCTTG	CCATATAGCA	35580
GCCTGGACGG	GTCCCTGCAG	CACCGCCATC	CTCGAGGCTC	AGGCCCACTT	TCTGCAGTGC	35640
CACAGGCACC	CCCCCCCCC	CATAGCGGCT	CCGGCCCGGC	CAGCCCCGGC	TCATTTAAAG	35700
GCACCGAGCC	CCGTTACCGG	GGGATGGGGG	AGTCCGAGAC	AGAATGACTT	CTTTATCCTG	35760
CTGACTCTGG	AAAGCCCGGC	GCCTTGTTGAT	CCATTGCAAA	CCGAGAGTCA	CCTCGTGTTT	35820
AGAACACGGA	TCCACTCCCA	AGTTCAGTGG	GGGGATGTGA	GGGGTGTGGC	AGGTAGGACG	35880
AAGGACTCTC	TTCTTTCTGA	TTCCGTCTGC	ACAGTGGGGC	CTAGGGCTGG	AGCTCTCTCC	35940
GTGCGGACCG	CTGACTCCCT	CTACCTTGCG	TTCCCTCGGC	CCCACCCTGG	AACGCCGGGG	36000
CTTGCGAGAT	TCTGGCCCTT	TCTGGCCCTT	CAGTCGCTGT	CAGAAACCCC	ATCTCATGCT	36060
CGGATGCCCC	GAGTGACTGT	GGCTCGCACC	TCTCCGAAAT	CATTGGAAAT	CTTCTCTCTA	36120
CGCGCGGCCA	CCTGAAACCA	CAGGAGCTCG	GGACACACGT	GCTTTCGGGA	GAGAATGCTG	36180
AGAGTCTCTC	GCCGACTCTC	TCTTGACTTG	AGTTCCTCGT	GGGTGCGTGG	TTAAGACGTA	36240
GTGAGACCAG	ATGTATTAAC	TCAGGCCGGG	TGCTGGTGGC	TCACGCCTGT	AACCCCAACA	36300
CTTTGGGAGG	CCGAGGCCGT	AGGATCCCTC	GAGGAATCCG	CTAACCTTGG	GGAGGTAG	36360
GTTGCGAGTA	GTTAGCCATA	GTTGTGTAC	TGCTGCCAG	TCTGGGCGAA	AGACAGAAATG	36420
AGGCCCTGCC	ACAGGCAGGC	AGGCAGGCAG	GCAGGCAGAA	AGACAACAGC	TGTATTATGT	36480
TCTTCTCAGG	GTAGGAAGCA	AAAATAACAG	AATACAGCAC	TTAATTAATT	TTTTTTTTTT	36540
CCTTCGGACG	GAGTTTCACT	CTTGGTGCCC	ACGCTGGAGT	GCAGTGGCAC	CATCTCGGCT	36600
CACCGCAACC	TCCACCTCCC	GCGTTCAAGC	GATTCTCCTG	CCTCAGCCTC	CTGAGTAGCT	36660
GGGATTACAG	GGAGGAGCCA	CCACACCCAG	CTGATTTTGT	ATTGTTAGTA	GAGACGGCAT	36720
TTCTCCATGT	GGGTCAGGCT	GGTCTCGAAC	TGGCGACCCC	AGTGGATCTG	CCCGCCCCGG	36780
CCTCCCAAAG	TGCTGGGGTG	ACAGGCGTGA	GCCATCGTGA	CTGGCCGGCT	ACGTTTATTT	36840
ATTTATTTTT	TTAATTATTT	TACTTTTTTT	TAGTTTTTCCA	TTTTAATCTA	TTTATTTATT	36900
TACATTTATT	TATTTATTTA	TTTATTTACT	TATTTATTTA	TTTTTCGAGAC	AGACTCTCGC	36960
TCTGCTGCCC	AGGCTGGAGT	GCAGCGGCGT	GATCTCGGCT	CACTGCAACG	TCCGCCTCCC	37020
GGGTTACCGC	CATTCTCCTG	CCTCAGCCTC	CCAAGTAGCT	GGGACTACAG	GGCCCCGCCA	37080
CCGTGCCCGG	GTAACTTTTT	GTATTTTGAG	TAGAGTAGGG	GTTTCACTGT	GCTTACAGAG	37140
ATGGTCTCGA	TCTCCTGACC	CCGTGATCCG	TCCACCTCGG	CCTCCCAAAG	TGCTGGGATG	37200
ACAGGCGTGA	GCCACCGGCC	CCGGCCTATT	TATCTATTTA	TTAACTTTGA	GTCCAGGTTA	37260
TGAAACCAGT	TAGTTTTTGT	AATTTTTTTT	TTTTTTTTTT	TTTTTTTGAGA	CGAGGTTTCA	37320
CCGTGTTGCC	AAGGCTTGGA	CCGAGGGATC	CACCGGCCCT	CGGCCTCCCA	AAAGTGCGGG	37380
GATGACAGGC	CGCAGCCTAC	CGCGCCCGGA	CCCCCCTTTT	CCCCCTCCCC	CGCTTGTCTT	37440
CCCGACAGAC	AGTTTTCACG	CAGAGCGTTT	GGCTGGCGTG	CTTAAGTCTA	TTCTAAATAG	37500
AAATTTGGGA	CGTCAGCTTC	TGGCCTCACG	GACTCTGAGC	CGAGGAGTCC	CCTGGTCTGT	37560
CTATCACAGG	ACCGTACACG	TAAGGAGGAG	AAAAATCGTA	ACGTTCAAAG	TCAGTCATTT	37620
TGTGATACAG	AAATACACGG	ATTCACCCAA	AACACAGAAA	CCAGTCTTTT	AGAAATGGCC	37680
TTAGCCCTGG	TGTCCGTGCC	AGTGATTCTT	TTCCGGTTTG	ACCTTGACTG	AGAGGATTCC	37740
CAGTCGGTCT	CTCGTCTCTG	GACGGAAGTT	CCAGATGATC	CGATGGGTGG	GGGACTTAGG	37800
CTGCGTCCCC	CCAGGAGCCC	TGGTTCGATTA	GTTGTGGGGA	TCGCCTTGA	GGGCGCGGTG	37860

-222-

ACCCACTGTG	CTGTGGGAGC	CTCCATCCTT	CCCCCACC	CCTCCCCAGG	GGGATCCCAA	37920
TTCATTCCGG	GCTGACACGC	TCACTGGCAG	GCGTCGGGCA	TCACCTAGCG	GTCAGTGTTA	37980
CTCTGAAAAC	GGAGGCCTCA	CAGAGGAAGG	GAGCACCAGG	CCGCCTGCGC	ACAGCCTGGG	38040
GCAACTGTGT	CTTCTCCACC	GCCCCGCCC	CCACCTCCAA	GTTCCTCCCT	CCCTTGTTC	38100
CTAGGAAATC	GCCACTTTGA	CGACCGGGTC	TGATTGACCT	TTGATCAGGC	AAAAACGAAC	38160
AAACAGATAA	ATAAATAAAA	TAACACAAAA	GTAATAACT	AAATAAAATA	AGTCAATACA	38220
ACCCATTACA	ATACAATAAG	ATACGATACG	ATAGGATGCG	ATAGGATACG	ATAGGATACA	38280
ATACAATAGG	ATACGATACA	ATACAATACA	ATACAATACA	ATACAATACA	ATACAATACA	38340
ATACAATACA	ATACAATACG	CCGGGCGCGG	TGGCTCATGC	CTGTCATCCC	GTCACCTTGG	38400
GATGCCGAGG	TGGACGCATC	ACCTGAAGTC	GGGAGTTGGA	GACAAGCCCC	ACCAACATGG	38460
AGAAATCCCC	TCTCAATTGA	AAATACAAAA	CTAGCCGGGC	GCGGTGGCAC	ATGCCTATAA	38520
TCCCAGCTCG	TAGGAAGGCT	GAGGCAGGAG	AATCGCTTGA	ACCTGGGAAG	CGGAGGTTGC	38580
AGTGAGCCGA	GATTGCGCCA	TCGCACTCCA	GTCTGAGCAA	CAAGAGCGAA	ACTCCGTCTC	38640
AAAAATAAAT	ACATAAATAA	ATACATACAT	ACATACATAC	ATACATACAT	ACATACATAC	38700
ATAAATTAAA	ATAAATAAAT	AAAATAAAAT	AAATAAATGG	GCCCTGCGCG	GTGGCTCAAG	38760
CCTGTCTATC	CCTCACTTTG	GGAGGCCAAG	GCCGGTGGAT	CAAGAGGCGG	TCAGACCAAC	38820
AGGGCCAGTA	TGGTGAAACC	CCGTCTCTAC	TCACAATACA	CAACATTAGC	CGGGCGCTGT	38880
GCTGTGCTGT	ACTGTCTGTA	ATCCCAGCTA	CTCGGGAGGC	CGAGCTGAGG	CAGGAGAATC	38940
GCTTGAACCT	GGGAGGCGGA	GGTTGCAGTG	AGCCGAGATC	GCGCCACTGC	AACCCAGCCT	39000
GGGCGACAGA	GCGAGACTCC	GTCTCCAAAA	AATGAAAATG	AAAATGAAAC	GCAACAAAAT	39060
AATTAAAAAG	TGAGTTTCTG	GGGAAAAAGA	AGAAAAGAAA	AAAGAAAAAA	ACAACAAAAC	39120
AGAACAACCC	CACCGTGACA	TACACGTACG	CTTCTCGCCT	TTGAGGCCT	CAAACACGTT	39180
AGGAATTATG	CGTGATTCT	TTTTTTAACT	TCATTTTATG	TTATTATCAT	GATTGATGTT	39240
TCGAGACGGA	GTCTCGGAGG	CCCGCCCTCC	CTGGTTGCC	AGACAACCCC	GGGAGACAGA	39300
CCCTGGCTGG	GCCCGATTGT	TCTTCTCCTT	GGTCAGGGGT	TTCTTGTCT	TTCTTCGTGT	39360
CTTTAACCCG	CGTGGACTCT	TCCGCCTCGG	GTTTGACAGA	TGGCAGCTCC	ACTTTAGGCC	39420
TTGTTGTTGT	TGGGGACTTT	CCTGATTCTC	CCCAGATGTA	GTGAAAGCAG	GTAGATTGCC	39480
TTGCCTGGCC	TTGCCTGGCC	TTGCCTTTTC	TTTCTTTCTT	TCTTTCTTTA	TTACTTTCTC	39540
TTTTCTTCT	TCTTCTTCTT	CTTTTCTTTG	AGACAGAGTT	TCACTCTTGT	TGCCCAGGCT	39600
AGAGGGCAAT	GGCGCGATCT	CGGCTCACCG	CACCCCTCCG	CTCCCAGGTT	CAAGCGATT	39660
TCCTGCCTCA	GCCTCCTGAT	TAGCTGGGAT	TACAGGCATG	GGCCACCGTG	CTGGCTGATG	39720
TTTGTACTTT	TAGTAGAGAC	GGTGTTTTTT	CATGTTGGTC	AGGCTGGTCT	CCCACTCCCA	39780
ACCTCAGGTG	GTCCGCCTGC	CTTAGCCTCC	CAAAGTGCTG	GGATGACAGG	CGTGCAACCG	39840
CGCCCAGCCT	CTCTCTCTCT	CTCTCTCTCT	CTCGCTCGCT	TGCTTGCTTG	CTTTCGTGCT	39900
TTCTTGCTTT	CCCGTTTTCT	TGCTTTCTTT	CTTCTTTCTG	TTTCTTTCTT	GCTTGCTTTT	39960
TTGCTTGCTT	GCTTGCTTTT	GTGCTTTCTT	GCTTTCTCTG	TTTCTTTCTT	TCTTTCTTTT	40020
TTTCTTTCTT	TTGTTTCTTT	CTTGCTTGCT	TTCTTGCTTG	CTTGCTTGCT	TTCTTGCTTT	40080
CTTGCTTTCC	TGTTTCTTTT	CTTCTTTCTT	TTCTTTCTTT	TCTTTCTTGC	TTGCTTTCTT	40140
GCTTGCTTGC	TTTCTGCTTT	TCTTGTTTTT	TCGATTTCTT	TCTTTCTTTT	GTTTCTTTCT	40200
TGCTTGCTTT	CTTGCTTGCT	TGCTTTCTG	CTTCTTGCTT	TCCTGTTTTT	TTTCTTTCTT	40260
TCTTTCTTTT	GTTTCTTTCT	TGCTTGCTTT	CTTGCTTGCT	TGCTTTCTG	CTGCTTGCTT	40320
TCTCGATTTT	TTTCTTTCTT	TTGTTTCTTT	CCTGCTTGCT	TTCTTGCTTG	ATTGCTTTCT	40380
TGCTTTCTTG	CTTTCTTGTT	TTCTTTCTTT	CTTTTGTTTT	TTTCTTTCTT	GCTTCCTTGT	40440
TTTCTTGCTT	TCTTGCTTGC	TTGCTTTCTG	GCTTTCTTGT	TTTCTTGCTT	TCTTTCTTTT	40500
GTTTCTTTCT	TGCTTGCTTT	CTTGCTTTCT	TGTTTTCTTG	CTTTCTTGCT	TGCTTGCTTT	40560
CGTGTGTTCT	TTCTTGCTTT	CTTTTCTTTT	TTTCTTTCTT	TTTCTTTCTT	TTCTTGCTTT	40620
CTTTTCTTTT	ATCATCATCT	TTCTTTCTTT	CCTTTCTTTT	TTTCTTTCTT	TCTATCTTTT	40680
TTTCTTTCTT	TCTTTCTTTT	TTTCTTTCTT	TCTTTCTGTT	TCGTCCTTTT	GAGACAGAGT	40740
TTCACTCTTG	TTTCCACGGC	TAGAGTGCAA	TGGCGCGATC	TTGGCTCACC	GCACCTTCCG	40800
CCTCCCGGGT	TCGAGCGCTT	CTCCTGCCTC	CAGCCTCCCG	ATTAGCGGGG	ATTGACAGGG	40860
AGGCACCCCC	ACGCCTGGCT	TGGCTGATGT	TTGTGTTTTT	AGTAGGCACG	CCGTGTCTCT	40920
CCATGTTGCT	CAGGCTGGTC	TCCAACCTCC	GACCTCCTGT	GATGCGCCCA	CCTCGGCCCT	40980
TCGAAGTGCT	GGGATGACGG	GCGTGACGAC	CGTGCCCGGC	CTGTTGACTC	ATTTCGCTTT	41040
TTTATTTCTT	TCGTTTCCAC	GCGTTTACTT	ATATGTATTA	ATGTAAACGT	TTCTGTACGC	41100
TTATATGCAA	ACAACGACAA	CGTGTATCTC	TGCATTGAAT	ACTCTTGCGT	ATGGTAAATA	41160
CGTATCGGTT	GTATGGAAAT	AGACTTCTGT	ATGATAGATG	TAGGTGTCTG	TGTTATACAA	41220
ATAAATACAC	ATCGCTCTAT	AAAGAAGGGA	CCGTCGATAA	AGACGTTTAT	TTTACGTATG	41280
AAAAGCGTCG	TATTTATGTG	TGTAAATGAA	TCCAGCGTAC	GTAGTTATCT	CTGTTTCTTT	41340
TCTTCCTCTC	CTTTCGTGTT	TTCTTCTTCT	CTTCTTCTCT	TTCTTCTCTT	CTTTAGGTTT	41400
TTCTTCTCTT	CTTCTTCTCT	TTCTTCTCTT	CTTCTTCTCT	TTTTTCTCTT	CGTGCTTTAT	41460
TTCTTCTCTG	TTCTTCTCTT	TTCTTCTCTT	TTCTTCTCTT	CTCTGTTTCT	TTTTTCTCTT	41520

-223-

TTTCCTTCGT	TTCTTTCCTC	ATTCTTTCTC	TCTTTTTCGT	TGTTTCTTTC	CTTCCCCTCT	41580
GTCTTTTAAA	AAATTGGAGT	GTTTCAGAAG	TTTACTTTGT	GTATCTACGT	TTTCTAAATT	41640
GTCTCTCTTT	TCTCCATTTT	CTTCCTCCCT	CCCTCCCTCC	CTCCCTGCTC	CCTTCCCTCC	41700
CTCCTTCCCT	TTCGCCATCT	GTCTCTTTTC	CCCACTCCCC	TCCCCCGTCT	TGTCTCTGCG	41760
TGGATTCCGG	AAGAGCCTAC	CGATTCTGCC	TCTCCGTGTG	TCTGCAGCGA	CCCCGCGACC	41820
GAGTCCTTGT	GTGTTCTTTC	TCCCTCCCTC	CCTCCCTCCC	TCCCTCCCTC	CCTCCCTGCT	41880
TCCGAGAGGC	ATCTCCAGAG	ACCGCGCCGT	GGGTTGTCTT	CTGACTCTGT	CGCGGTCGAG	41940
GCAGAGACGC	GTTTTGGGCA	CCGTTTGTGT	GGGGTTGGGG	CAGAGGGGCT	GCGTTTTTCGG	42000
CCTCGGGAAG	AGCTTCTCGA	CTCACGGTTT	CGCTTTCGCG	GTCCACGGGC	CGCCCTGCCA	42060
GCCGGATCTG	TCTCGCTGAC	GTCCGCGGCG	GTTGTCGGGC	TCCATCTGGC	GGCCGCTTTG	42120
AGATCGTGCT	CTCGGCTTCC	GGAGCTGCGC	TGGCAGCTGC	CGAGGGAGGG	GACCGTCCCC	42180
GCTGTGAGCT	AGGCAGAGCT	CCGGAAGGCC	CGCGTTCGTC	AGCCCGGCTG	GCCCGGTGGC	42240
GCCAGAGCTG	TGGCCGCTCG	CTTGTGAGTC	ACAGCTCTGG	CGTGCAGGTT	TATGTGGGGG	42300
AGAGGCTGTC	GCTGCGCTTC	TGGGCCCCGCG	GCGGGCGTGG	GGCTGCCCGG	GCCGGTCGAC	42360
CAGCGCGCCG	TAGCTCCCGA	GGCCCGAGCC	GCGACCCGGC	GGACCCGCCG	CGCGTGGCGG	42420
AGGCTGGGGA	CGCCCTTCCC	GGCCCGGTCT	CGGTCCGCTC	ATCCTGGCCG	TCTGAGGCGG	42480
CGGCCGAATT	CGTTTCCGAG	ATCCCCGTGG	GGAGCCGGGG	ACCGTCCCGC	CCCCGTCCCC	42540
CGGGTGCCGG	GGAGCGGTCC	CCGGGCCGGG	CCGCGTCCC	TCTGCCGCGA	TCCTTTCTGG	42600
CGAGTCCCCG	TGGCCAGTCG	GAGAGCGCTC	CCTGAGCCGG	TGCGGCCCGA	GAGGTCGCGC	42660
TGGCCGGCCT	TCGGTCCCTC	GTGTGTCCCG	GTCGTAGGAG	GGGCCGGCCG	AAAATGCTTC	42720
CGGCTCCCGC	TCTGGAGACA	CGGGCCGGCC	CCTGCGTGTG	GCCAGGGCGG	CCGGGAGGGC	42780
TCCCCGGCCC	GGCGCTGTCC	CCGCGTGTGT	CCTTGGGTTG	ACCAGAGGGA	CCCCGGGCGC	42840
TCCGTGTGTG	GCTGCGATGG	TGGCGTTTTT	GGGGACAGGT	GTCCGTGTCC	GTGTCGCGCG	42900
TCGCCTGGGC	CGGCGGCGTG	GTCGGTGACG	CGACCTCCCG	GCCCCGGGGG	AGGTATATCT	42960
TTCGCTCCGA	GTCGGCAATT	TTGGGCCGCC	GGGTTATAT			42999

(2) INFORMATION FOR SEQ ID NO:18:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 175 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTCCCCGCGC	GCCCCCGTGT	TCGCCGTTCC	CGTGGCGCGG	ACAATGCGGT	TGTGCGTCCA	60
CGTGTGCGTG	TCCGTGCAGT	GCCGTTGTGG	AGTGCCTCGC	TCTCCTCCTC	CTCCCCGGCA	120
GCGTTCCAC	GGTTGGGGAC	CACCGGTGAC	CTCGCCCTCT	TCGGGCCTGG	ATCCG	175

(2) INFORMATION FOR SEQ ID NO:19:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 755 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

-224-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

GGTCTGGTGG	GAATTGTTGA	CCTCGCTCTC	GGGTGCGGCC	TTTGGGGAAC	GGCGGGGTCG	60
GTCGTGCCCC	GCGCCGACG	TGTGTCGGGG	CCCACTTCCC	GCTCGAGGGT	GGCGGTGGCG	120
GCGGCGTTGG	TAGTCTCCCG	TGTTGCGTCT	TCCCGGGCTC	TTGGGGGGGG	TGCCGTCGTT	180
TTCCGGGGCCG	GCGTTGCTTG	GCTTACGCAG	GCTTGGTTTG	GGACTGCCTC	AGGAGTCGTG	240
GGCGGTGTGA	TTCCCGCCGG	TTTTGCCTCG	CGTCTGCCTG	CTTTGCCTCG	GGTTTGCTTG	300
GTTCTGTCT	CGGGAGCGGT	GGTTTTTTTT	TTTTTCGGGT	CCCGGGGAGA	GGGGTTTTTC	360
CGGGGGACGT	TCCCGTCGCC	CCCTGCCGCC	GGTGGGTTTT	CGTTTCGGGC	TGTGTTCTGTT	420
TCCCTTCCC	CGTTTCGCCG	TCGGTTCCTC	CCGGTCGGTC	GGCCCTCTCC	CCGGTCGGTC	480
GCCCGGCCGT	GCTGCCGGAC	CCCCCTTCT	GGGGGGGATG	CCCGGGCACG	CACGCGTCCG	540
GGCGGCCACT	GTGGTCCGGG	AGCTGCTCGG	CAGGCGGGTG	AGCCAGTTGG	AGGGGCGTCA	600
TGCCCCCGCG	GGTCCCCGTG	GCCGACGCGG	CGTGTTCTTT	GGGGGGGCCT	GTCCGTGCGG	660
GAAGGCTGCG	CACGTTGTG	GTCCTTGCGA	GGGAAAGAGG	CTTTTTTTTT	TTAGGGGGTC	720
GTCCTTCGTC	GTCCCGTCGG	CGGTGGATCC	GGCCT			755

(2) INFORMATION FOR SEQ ID NO:20:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 463 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

GGCCGAGGTG	CGTCTGCGGG	TTGGGGGCTCG	TCCGGCCCCG	TCGTCCTCCG	GGAAGGCGTT	60
TAGCGGGTAC	CGTCGCCGCG	CCGAGGTGGG	CGCACGTCGG	TGAGATAACC	CCGAGCGTGT	120
TTCTGGTTGT	TGGCGGCGGG	GGCTCCGGTC	GATGTCTTCC	CCTCCCCCTC	TCCCCGAGGC	180
CAGGTACAGC	TCCGCCTGTG	GGCTTCGTCG	GCCGTCTCCC	CCCCCCTCAC	GTCCCTCGCG	240
AGCGAGCCCG	TCCGTTTCGAC	CTTCCCTCCG	CCTTCCCCCC	ATCTTTCCGC	GCTCCGTTGG	300
CCCCGGGGTT	TTACAGGCGC	CCCCACGCT	CCTCCGCCTC	TCCGCCCCGTG	GTTTGGACGC	360
CTGGTTCCGG	TCTCCCCGCC	AAACCCCGGT	TGGGTTGGTC	TCCGGCCCCG	GCTTGCTCTT	420
CGGGTCTCCC	AACCCCCGGC	CGGAAGGGTT	CGGGGGTTCC	GGG		463

(2) INFORMATION FOR SEQ ID NO:21:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 378 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

GGATTCTTCA	GGATTGAAAC	CCAAACCGGT	TCAGTTTCCT	TTCCGGCTCC	GGCCGGGGGG	60
GGCGGCCCCG	GGCGGTTTGG	TGAGTTAGAT	AACCTCGGGC	CGATCGCACG	CCCCCGTGG	120
CGGCGACGAC	CCATTGGAAC	GTCTGCCCTA	TCAACTTTCG	ATGGTAGTCG	ATGTGCCTAC	180
CATGGTGACC	ACGGGTGACG	GGGAATCAGG	GTTCGATTCC	GGAGAGGGAG	CCTGAGAAAC	240

-225-

GGCTACCACA	TCCAAGGAAG	GCAGCAGGCG	CGCAAATTAC	CCACTCCCGA	CCCGGGGAGG	300
TAGTGACGAA	AAATAACAAT	ACAGGACTCT	TTCGAGGCC	TGTAATTGGA	ATGAGTCCAC	360
TTTAAATCCT	TTAAGCAG					378

(2) INFORMATION FOR SEQ ID NO:22:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 378 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

GATCCATTGG	AGGGCAAGTC	TGGTGCCAGC	AGCCGCGGTA	ATTCCAGCTC	CAATAGCGTA	60
TATTAAAGTT	GCTGCAGTTA	AAAAGCTCGT	AGTTGGATCT	TGGGAGCGGG	CGGGCGGTCC	120
GCCGCGAGGC	GAGTCACCGC	CCGTCCCCGC	CCCTTGCCCTC	TCGGCGCCCC	CTCGATGCTC	180
TTAGCTGAGT	TGTCCCGCGG	GGCCCGAAGC	GTTTACTTTG	AAAAAATTAG	AGTTGTTTCA	240
AAGCAGGCCC	GAGCCGCCTG	GATACCGCCA	GCTAGGAAAT	AATGGAATAG	GACCGCGGTT	300
CCTATTTTGT	TTGGTTTTTCG	GAAGTGAGCC	CATGATTAAG	GGAAACGGCC	GGGGGCATTC	360
CCTTATTGCG	CCCCCCTA					378

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 719 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

GGATCTTTCC	CGCTCCCCGT	TCCTCCCCGGC	CCCTCCACCC	GCGCGTCTCC	CCCCTTCTTT	60
TCCCCTCTCC	GGAGGGGGGG	GAGGTGGGGG	CGCGTGGGCG	GGGTGCGGGG	TGGGGTCGGC	120
GGGGGACCGC	CCCCGGCCGG	CAAAAGGCCG	CCGCCGGGCG	CACTTCAACC	GTAGCGGTGC	180
GCCGCGACCG	GCTACGAGAC	GGCTGGGAAG	GCCCCACGGG	GAATGTGGCT	CGGGGGGGGC	240
GGCGCGTCTC	AGGGCGCGCC	GAACCACCTC	ACCCCGAGTG	TTACAGCCCT	CCGGCCGCGC	300
TTTCGCGGAA	TCCCGGGGCC	GAGGGGAAGC	CCGATACCCG	TCGCCGCGCT	TTTCCCCTCC	360
CCCCGTCCGC	CTCCCGGGCG	GGCGTGGGGG	TGGGGGCCCG	GCCGCCCTC	CCACGCCCGT	420
GGTTTCTCTC	TCTCCCGGTC	TCGGCCGGTT	TGGGGGGGGG	AGCCCGGTTG	GGGGCGGGGC	480
GGACTGTCCT	CAGTGCGCC	CGGGCGTCGT	CGCGCCGTG	GGCCCGGGG	GTCTCTCGG	540
TCACGCCGCC	CCGACGAAG	CCGAGCGCAC	GGGGTCGGCG	GCGATGTCGG	CTACCCACCC	600
GACCCGTCTT	GAAACACGGA	CCAAGGAGTC	TAACGCGTGC	GCGAGTCAGG	GGCTCGCACG	660
AAAGCCGCCG	TGGCGCAATG	AAGGTGAAGG	GCCCCGTCCG	GGGGCCCCGAG	GTGGGATCC	719

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 685 base pairs

-226-

- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

- (ii) MOLECULE TYPE: Genomic DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CGAGGCCTCT	CCAGTCCGCC	GAGGGCGCAC	CACCGGCCCG	TCTCGCCCCG	CGCGTCGGGG	60
AGGTGGAGCA	CGAGCGTACG	CGTTAGGACC	CGAAAGATGG	TGAACTATGC	CTGGGCAGGG	120
CGAAGCCAGA	GGAAACTCTG	GTGGAGGTCC	GTAGCGGTCC	TGACGTGCAA	ATCGGTCGTC	180
CGACCTGGGT	ATAGGGGCGA	AAGACTAATC	GAACCATCTA	GTAGCTGGTT	CCCTCCGAAG	240
TTCCTTCAG	GATAGCTGGC	GCTCTCGCAA	CCTTCGGAAG	CAGTTTTATC	CGGGTAAAGG	300
CGGAATGGAT	TAGGAGGTCT	TGGGGCCGGA	AACGATCTCA	AACTATTTCT	CAAACTTTAA	360
ATGGGTAAGG	AAGCCCGGCT	CGCTGGCGTG	GAGCCGGGCG	TGGAATGCCA	GTGCCTAGTG	420
GGCCACTTTT	GGTAAGCAGA	ACTGGCGCTG	CGGGATGAAC	CGAACGCCCG	GTTAAGGCGC	480
CCGATGCCGA	CGTTCATCAG	ACCCAGAAA	AGGTGTTGGT	TGATATAGAC	AGCAGGACGG	540
TGGCCATGGA	AGTCGGAATC	CGCTAAGGAG	TGTGTAACAA	CTCACCTGCC	GAATCAACTA	600
GCCCTGAAAA	TGGATGGCGC	TGGAGCGTCG	GGCCCATACC	CGGCCGTCGC	CGGCAGTCGG	660
AACGGGACGG	GACGGGAGCG	GCCGC				685

(2) INFORMATION FOR SEQ ID NO:25:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 33 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Genomic DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GAGGAATTCC	CCTATCCCTA	ATCCAGATTG	GTG	33
------------	------------	------------	-----	----

(2) INFORMATION FOR SEQ ID NO:26:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 35 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: Genomic DNA
- (iii) HYPOTHETICAL: NO
- (iv) ANTISENSE: NO
- (v) FRAGMENT TYPE:
- (vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

AAACTGCAGG	CCGAGCCACC	TCTCTTCTGT	GTTTG	35
------------	------------	------------	-------	----

-227-

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 33 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

AGGAATTCAC AGAAGAGAGG TGGCTCGGCC TGC

33

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 34 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

AGCCTGCAGG AAGTCATACC TGGGGAGGTG GCCC

34

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 80 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

AAACTGCAGG TTAATTAACC CTAACCCTAA CCCTAACCCT AACCCCTAACC CTAACCCTAA
CCCTAACCCT AACCCGGGAT

60

80

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 19 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single

-228-

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

TTGGGCCCTA GGCTTAAGG

19

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 25 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

GCCAGGGTTT TCCAGTCAC GACGT

25

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

GCTGCAAGGC GATTAAGTTG GGTAAC

26

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 26 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

-229-

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

TATGTTGTGT GGAATTGTGA GCGGAT

26

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: Genomic DNA

(iii) HYPOTHETICAL: NO

(iv) ANTISENSE: NO

(v) FRAGMENT TYPE:

(vi) ORIGINAL SOURCE:

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

GGGTTTAAAC AGATCTCTGC A

21

-230-

WHAT IS CLAIMED:

1. A method for producing an artificial chromosome, comprising:
introducing a DNA fragment into a cell, wherein the DNA
fragment comprises a selectable marker;
5 growing the cell under selective conditions to produce cells
that have incorporated the DNA fragment into their genomic DNA; and
selecting a cell that comprises a satellite artificial
chromosome [SATAC].
2. The method of claim 1, wherein the DNA fragment is
10 introduced into or adjacent to an amplifiable region of a chromosome in
the cell.
3. The method of claim 2, wherein the amplifiable region
comprises rDNA.
4. The method of claim 2, wherein the amplifiable region
15 comprises heterochromatin.
5. The method of claim 1 or claim 2, wherein the DNA is
introduced into pericentric heterochromatin in a chromosome of the cell.
6. The method of claim 1, wherein the cell is a mammalian cell.
7. The method of claim 1 or claim 2, further comprising,
20 isolating the SATAC.
8. The method of any of claims 1-7, wherein the DNA fragment
comprises a sequence of nucleotides that targets the fragment to the
heterochromatic region of a chromosome.
9. The method of claim 8, wherein the targeting sequence of
25 nucleotides comprises satellite DNA.
10. The method of any of claims 1-9, wherein the cell is a
human cell.
11. The method of any of claims 1-5 and 7- 9, wherein the cell is
a fish, insect, reptile, amphibian, arachnid, or rodent cell.

-231-

12. A SATAC produced by the method of any of claims 1-11.
13. An isolated substantially pure satellite artificial chromosome [SATAC].
14. The SATAC of claim 13 that is a megachromosome,
5 comprising about 50 to about 450 megabases [Mb].
15. The SATAC of claim 13, comprising about 250 to about 400 Mb.
16. The SATAC of claim 13, comprising about 150 to about 200 Mb.
- 10 17. The SATAC of claim 13, comprising about 90 to about 120 Mb.
18. The SATAC of claim 13, comprising about 15 to about 60 Mb.
19. A cell containing an artificial chromosome, wherein the
15 artificial chromosome is produced by the method of any of claims 1-11.
20. A cell containing the SATAC of any of claims 12-19.
21. The cell of claim 19 or claim 20 that is a mammalian cell.
22. The method of any of claims 1-11, wherein the SATAC is a megachromosome, and the method further comprises:
20 introducing a fragmentation vector, whereby the megachromosomes in the cells are reduced in size,
and identifying cells that contain SATACs that are about 15 to about 60 Mb.
23. The method of any of claims 1-11, wherein the SATAC is a
25 megachromosome, and the method further comprises, exposing the cells to conditions, whereby cells that contain truncated megachromosomes are produced.
24. The method of claim 23, wherein the conditions are selected from among exposure to X-rays, growth in the presence of an agent that
30 destabilizes base pairing in the chromosome.

-232-

25. The method of claim 24, wherein the agent is bromodeoxyuridine.

26. The method of any of claims 22-25, further comprising selecting a cell that comprises a satellite artificial chromosome [SATAC]
5 that comprises about 15 to about 60 Mb.

27. A cell containing an artificial chromosome, wherein the artificial chromosome is produced by the method of any of claims 22-25.

28. The cell of any of claims 19-21, 25-27, wherein the artificial chromosome is a SATAC comprising about 10 to about 60 Mb.

10 29. An isolated substantially pure satellite artificial chromosome [SATAC] of claim 13 that comprises about 10 to about 60 Mb.

30. The method of any of claims 1-11 and 22-26, further comprising isolating the SATAC from the cell.

15 31. The method of claim 30, wherein isolation is effected by:
isolating metaphase chromosomes;
distinguishing SATACs from endogenous chromosomes; and
separating the SATACs from endogenous chromosomes.

32. The method of claim 31, wherein:
the SATACs are distinguished from endogenous chromosomes by
20 staining the chromosomes with DNA sequence-specific dyes; and
separation is effected by flow cell sorter.

33. A method for producing an artificial chromosome, comprising:

25 introducing a DNA fragment into a cell, wherein the DNA fragment comprises a selectable marker,
growing the cell under selective conditions to produce cells that have incorporated the DNA fragment into their genomic DNA,
selecting from among those cells, a cell that comprises a *de novo* centromere.

-233-

34. The method of claim 33, further comprising isolating that cell with the chromosome that comprises the *de novo* centromere, and growing the cell under conditions whereby a cell with a sausage chromosome is produced.

5 35. The method of claim 34, further comprising isolating the cell with the sausage chromosome; and growing the cell under conditions whereby a first SATAC is produced.

36. The method of claim 35, wherein the DNA fragment is introduced into or adjacent to an amplifiable region of a chromosome in
10 the cell.

37. The method of claim 36, wherein the amplifiable region comprises rDNA.

38. The method of claim 36, wherein the amplifiable region comprises heterochromatin.

15 39. The method of claim 35 or claim 36, wherein the DNA is introduced into pericentric heterochromatin in a chromosome of the cell.

40. The method of any of claims 33-39, further comprising:
introducing a fragmentation vector that is targeted to the first SATAC; growing the cells; and selecting a cell that comprises a second
20 SATAC, wherein the second SATAC is smaller than the first SATAC.

41. The method of claim 40, wherein the selected cell has a dicentric chromosome comprising the *de novo* centromere.

42. The method of claim 40, wherein the selected cell has a formerly dicentric chromosome and a minichromosome comprising the *de*
25 *nov* centromere.

43. The method of claim 40, wherein the selected cell has a formerly dicentric chromosome.

-234-

44. A method for producing an artificial chromosome, comprising
introducing a DNA fragment into a cell, wherein the DNA
fragment comprises a selectable marker;
growing the cell under selective conditions to produce cells
5 that have incorporated the DNA fragment into their genomic DNA;
selecting from among those cells a cell that has produced a
dicentric chromosome; and
growing that cell under selective conditions, whereby a cell
that contains a chromosome comprising a heterochromatic arm is
10 produced.
45. The method of claim 44, further comprising selecting the cell
with the chromosome comprising the heterochromatic arm and growing it
in the presence of an agent that destabilizes the chromosome.
46. The method of claim 45, further comprising identifying cells
15 that contain a heterochromatic chromosome that is about 50 to about
400 Mb.
47. The method of any of claims 44-46, wherein the DNA
fragment is introduced into or adjacent to an amplifiable region of a
chromosome in the cell.
- 20 48. The method of claim 47, wherein the amplifiable region
comprises rDNA.
49. The method of claim 47, wherein the amplifiable region
comprises heterochromatin.
50. The method of claim 47, wherein the DNA is introduced into
25 pericentric heterochromatin in a chromosome of the cell.
51. A method for producing a transgenic (non-human) animal,
comprising introducing a satellite artificial chromosome [SATAC] into an
embryonic cell.
52. The method of claim 51, wherein the embryonic cell is a
30 stem cell.

-235-

53. The method of claim 51, wherein the embryonic cell is in an embryo.

54. The method of any of claims 51-53, wherein the SATAC comprises heterologous DNA that encodes a gene product.

5 55. The method of any of claims 51-54, wherein the SATAC comprises heterologous DNA that encodes a therapeutic product.

56. The method of claim 55, wherein the anti-HIV ribozyme is an anti-*gag* ribozyme, and the tumor suppressor gene is p53.

57. The method of claim 54, wherein the product comprises an
10 antigen that upon expression induces a immunoprotective response against a pathogen in the transgenic (non-human) animal.

58. The method of claim 54, wherein the product comprises a plurality of antigens that upon expression induce an immunoprotective response against a plurality of pathogens.

15 59. The method of any of claims 51-58, wherein the transgenic (non-human) animal is a fish, insect, reptile, amphibian, arachnid or mammal.

60. The method of any of claims 51-59, wherein the SATAC is introduced by cell fusion, lipid-mediated transfection by a carrier system,
20 microinjection, microcell fusion, electroporation, microprojectile, nuclear transfer, bombardment or direct DNA transfer.

61. A transgenic (non-human) animal produced by the method of any of claims 51-60.

62. The transgenic animal that is a fish, insect, reptile, amphibian,
25 arachnid, or mammal.

63. A method of for producing a transgenic plant or animal, comprising:

introducing a DNA fragment into a cell, wherein the DNA fragment comprises a selectable marker;

-236-

growing the cell under selective conditions to produce cells that have incorporated the DNA fragment into their genomic DNA; and

selecting a cell that comprises a minichromosome that is about 10 Mb to about 50 Mb that comprises the selectable marker and
5 euchromatin;

isolating the minichromosome and introducing it into a plant or animal cell.

64. The method of claim 63, wherein: after selecting the cell, DNA encoding a gene product or products is introduced into the cell, and
10 the cell is grown under selective conditions, whereby cells comprising minichromosomes comprising the DNA encoding the gene product(s) are produced.

65. The method of claim 64, wherein: after selecting the cell, DNA encoding a gene product or products is introduced into the cell, and the cell is grown under selective conditions, whereby cells comprising
15 SATACS that comprise the DNA encoding the gene product(s) are produced.

66. A method for cloning a centromere from an animal or plant, comprising:
20 preparing a library of DNA fragments that comprise the genome of the plant or animal;

introducing each of the fragments into mammalian satellite artificial chromosomes [SATACs], wherein:

each SATAC comprises a centromere from a different
25 species from the selected plant or animal, and a selectable marker;

introducing each of the SATACs into the cells and growing the cells under selective conditions;

identifying cells that have a SATAC; and

-237-

selecting from among those cells any that have a SATAC comprising a centromere that differs from the centromeres in the original SATAC.

67. A cell line having the identifying characteristics of any of
5 TF1004G19C5, 19C5xHa4, H1D3 and G3D5, which have been deposited at the ECACC under Accession Nos. 96040926, 96040927, 96040929, and 96040928, respectively.

68. A cell line, comprising a megachromosome that comprises about 50-400 Mb.

10 69. A cell line of claim 68, wherein the megachromosome comprises 250 to about 400 Mb.

70. A cell line of claim 68, wherein the megachromosome comprises about 150 to about 200 Mb.

15 71. A cell line of claim 68, wherein the megachromosome comprises about 90 to about 120 Mb.

72. A cell line of claim 68, wherein the megachromosome comprises about 60 to about 100 Mb.

73. A method for gene therapy, comprising:
introducing a SATAC that comprises DNA therapeutic product into
20 a target cell; and
introducing the resulting target cells into a host animal.

74. The method of claim 73, wherein the target cells are lymphocytes, stem cells, nerve cells, insect cells, chicken cells or muscle cells.

25 75. The method of claim 73, wherein the minichromosome is the minichromosome present in the cell line EC3/7C5.

76. The method of claim 73, wherein the chromosome is the λ neo-chromosome in the cell line KE1 2/4.

30 77. The artificial chromosome of claim 76 that is between about 20 Mb and about 200 Mb.

-238-

78. The artificial chromosome of claim 76 that is between about 100 Mb and about 200 Mb.

79. The artificial chromosome of claim 76 that is between about 20 Mb and about 200 Mb.

5 80. The artificial chromosome of claim 76 that is between about 1 Mb and about 15 Mb.

81. The method of claim 51, wherein the animal is a mammal or a oviparous animal and the SATAC includes proteins and regulatory elements for expression of genes in the milk of the animal or in the egg of
10 the animal.

82. The method of claim 81, wherein the animal is selected from among cows, goats, oxen, pigs and sheep.

83. The method of claim 81, wherein the animal is selected from among fowl.

15 84. The method of claim 51, wherein the SATAC includes DNA encoding genes for expression of human cell surface proteins, wher by the organs of the animal express the human proteins and will not be rejected upon transplantation into a human.

85. Isolated DNA, comprising the DNA having the sequence set
20 forth in SEQ ID NO. 13, 14 or 15.

86. An isolated DNA, comprising the DNA having the sequence set forth in SEQ ID NO. 13, 14 or 15.

87. An isolated DNA fragment, comprising a sequence of nucleotides set forth in any of SEQ ID Nos. 18-24.

25 88. A SATAC of claim 14, comprising a sequence of nucleotides set forth in any of SEQ ID Nos. 18-24.

89. A SATAC of claim 13, comprising a sequence of nucleotides set forth in any of SEQ ID Nos. 18-24.

30 90. A SATAC of claim 13, comprising a sequence of nucleotides set forth in any of SEQ ID Nos. 18-24.

-239-

91. A cellular production system, comprising a cell containing an artificial chromosome [AC], wherein the AC comprises multiple copies of a heterologous gene or a plurality of heterologous genes.

92. The cellular production system of claim 91, wherein the AC
5 is a SATAC.

93. The system of claim 91 or claim 92, wherein the heterologous genes encode proteins that comprise a metabolic pathway.

94. A method of expression of a product that is produced upon expression of a metabolic pathway, comprising culturing the system of
10 claim 93 under conditions whereby the proteins comprising the pathway are expressed to produce the product.

95. The method of claim 94, wherein the product is a vitamin, a hormone, a nucleotide, an amino acid, a protein or a peptide.

96. The method of claim 51, wherein the animal is oviparous.

15 97. The method of claim 51, wherein animal is a chicken.

98. The method of claim 51, wherein the animal is an insect.

99. A method for producing a transgenic plant, comprising introducing a satellite artificial chromosome [SATAC] of any of claims 13-18 or 88-90 into a plant cell; and culturing the cell under conditions
20 whereby a plant is generated.

100. The method of claim 99, wherein the SATAC is introduced by protoplast fusion, microinjection, microcell fusion, lipid-mediated gene transfer, electroporation, microprojectile bombardment or direct DNA transfer.

25 101. A method for producing a gene product(s), comprising introducing a satellite artificial chromosome [SATAC] of any of claims 13-18 or 88-90 into a cell; and culturing the cell under conditions whereby the gene product(s) is (are) expressed.

102 The method of claim 102, wherein the gene product is
30 produced by expression of a series of genes that encode proteins that

-240-

comprise a metabolic pathway; and the SATAC comprises each of these genes.

103. An in vitro synthesized artificial mammalian chromosome (ISMAC), comprising a centromere, a telemere, a megareplicator, and a
5 selectable marker, wherein the centromere is derived from a SATAC of any of claims 12-18 and 88-90.

104. An in vitro synthesized artificial mammalian chromosome (ISMAC), comprising a centromere, a telemere, a megareplicator, and a
10 selectable marker, wherein the centromere is derived from a SATAC of any of claims 12-18, and 88-90.

105. The ISMAC of claim 103 or claim 104, further comprising heterochromatin.

106. The ISMAC of any of claims 103-105, wherein the megareplicator comprises rDNA.

15 107. The ISMAC of any of claims 103-106, wherein the centromere is a human centromere.

108. The ISMAC of any of claims 103-107, wherein the centromere is derived from a megachromosome.

109. The ISMAC of any of claims 103, 105, 106 or 108, wherein
20 the centromere is derived from a cell line having all of the identifying characteristics of the cell line deposited under at the European Collection of Animal Cell Culture (ECACC) under Accession No. 96040929.

110. The method of claim 54, wherein the product is a hormone, antibody, cytokine, growth factor, regulatory protein, secreted proteins.

25 111. The method of claim 54, wherein the product is the cystic fibrosis transmembrane regulatory protein [CFTR], an anti-HIV ribozyme, or a tumor suppressor gene.

112. The method of claim 66, wherein the animal is a human.

113. A method of producing an in vitro synthesized artificial
30 mammalian chromosome (ISMAC) comprising, combining a centromere, a

-241-

telomere, a megareplicator, and a selectable marker to produce a replicable ISMAC, herein the centromere is derived from a SATAC of any of claims 12-18, and 88-90.

114. The method of claim 113, further comprising including rDNA
5 in the ISMAC.

115. The method of claim 113, wherein the telomere comprises a plurality of repeats of SEQ ID No. 29.

116. The method of claim 115, wherein the telomere is about 1 kB up to about 1 Mb, preferably about 1 kB up to about 500 kB.

10 117. The ISMAC of claim 103-108, wherein the telomere comprises a plurality of repeats of SEQ ID No. 29.

118. The ISMAC of claim 117, wherein the telomere is about 1 kB up to about 1 Mb, preferably about 1 kB up to about 500 kB.

119. A method for producing an artificial chromosome,
15 comprising:

introducing a DNA fragment into a cell, wherein the DNA fragment comprises a selectable marker;

growing the cell under selective conditions to produce cells that have incorporated the DNA fragment into their genomic DNA,
20 wherein the DNA fragment is introduced into or adjacent to an amplifiable region of a chromosome in the cell, whereby a minichromosome comprising the DNA derived from the amplifiable region is produced, wherein the minichromosome is an artificial chromosome that contains more euchromatin than heterochromatin.

25 120. The method of claim 119, wherein the amplifiable region is rDNA.

121. The method of claim 119 or 120, further comprising isolating the minichromosome.

122. A minichromosome produced by the methods of any of
30 claims 119-121.

-242-

123. The method of claim 1, further comprising introducing DNA encoding a gene product, wherein:

the DNA encoding the gene is on the fragment that comprises the selectable marker or is on a second DNA fragment; and the resulting

5 SATAC comprises the heterologous DNA that encodes a gene product.

1 / 5





FIG. 1

2 / 5



3 / 5




FIG. 3




FIG. 4




FIG. 5



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification⁶ : C12N 15/90, 15/85</p>	<p>A3</p>	<p>(11) International Publication Number: WO 97/40183 (43) International Publication Date: 30 October 1997 (30.10.97)</p>									
<p>(21) International Application Number: PCT/US97/05911 (22) International Filing Date: 10 April 1997 (10.04.97) (30) Priority Data: <table border="0"> <tr> <td>629,822</td> <td>10 April 1996 (10.04.96)</td> <td>US</td> </tr> <tr> <td>682,080</td> <td>15 July 1996 (15.07.96)</td> <td>US</td> </tr> <tr> <td>695,191</td> <td>7 August 1996 (07.08.96)</td> <td>US</td> </tr> </table> <p>(71) Applicants (for all designated States except US): THE BIOLOGICAL RESEARCH CENTER OF THE HUNGARIAN ACADEMY OF SCIENCES [HU/HU]; P.O. Box 521, H-6701 Szeged (HU). LOMA LINDA UNIVERSITY [US/US]; Loma Linda, CA 92350 (US). AMERICAN GENE THERAPY, INC. [CA/CA]; 5022 154th Street, Edmonton, Alberta T6H 5P6 (CA). (72) Inventors; and (75) Inventors/Applicants (for US only): HADLACZKY, Gyula [HU/HU]; Szamos U.I.A. IX. 36., H-6723 Szeged (HU). SZALAY, Aladar, A. [US/US]; 7327 Fairwood, Highland, CA 92346 (US). (74) Agent: SEIDMAN, Stephanie, L.; Brown, Martin, Haller & McClain, 1660 Union Street, San Diego, CA 92101-2926 (US).</p> </p>		629,822	10 April 1996 (10.04.96)	US	682,080	15 July 1996 (15.07.96)	US	695,191	7 August 1996 (07.08.96)	US	<p>(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</p> <p>Published <i>With a revised version of the international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i></p> <p>(88) Date of publication of the international search report: 5 February 1998 (05.02.98) (88) Date of publication of the revised version of the international search report: 30 April 1998 (30.04.98)</p>
629,822	10 April 1996 (10.04.96)	US									
682,080	15 July 1996 (15.07.96)	US									
695,191	7 August 1996 (07.08.96)	US									

(54) Title: **ARTIFICIAL CHROMOSOMES, USES THEREOF AND METHODS FOR PREPARING ARTIFICIAL CHROMOSOMES**

(57) Abstract

Methods for preparing cell lines that contain artificial chromosomes, methods for preparation of artificial chromosomes, methods for purification of artificial chromosomes, methods for targeted insertion of heterologous DNA into artificial chromosomes, and methods for delivery of the chromosomes to selected cells and tissues are provided. Also provided are cell lines for use in the methods, and cell lines and chromosomes produced by the methods. In particular, satellite artificial chromosomes that, except for inserted heterologous DNA, are substantially composed of heterochromatin, are provided. Methods for use of the artificial chromosomes, including for gene therapy, production of gene products and production of transgenic plants and animals are also provided.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

A. CLASSIFICATION F SUBJECT MATTER
IPC 6 C12N15/90 C12N15/85

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	G. HADLACZKY ET AL.: "Centromere formation in mouse cells cotransformed with human DNA and a dominant marker gene" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 88, 1991, WASHINGTON, DC, US, pages 8106-8110, XP002037028 cited in the application see the whole document --- -/--	1-50,75, 76

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

Z document member of the same patent family

Date of the actual completion of the international search

24 February 1998

Date of mailing of the international search report

10.03.98

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Mateo Rosell, A.M.

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 97/05911

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	EP 0 532 050 A (BIOLOGICAL RESEARCH CENTER OF THE HUNGARIAN ACADEMY OF SCIENCES) 17 March 1993	33-50, 75,76
Y	see the whole document, specially example 6	1-32
A	& US 5 288 625 A cited in the application	63-65
Y	--- K. FATYOL ET AL.: "Cloning and molecular characterization of a novel chromosome specific centromere sequence of Chinese hamster" NUCLEIC ACIDS RESEARCH, vol. 22, no. 18, 1994, OXFORD, GB, pages 3728-3736, XP002037027 cited in the application see the whole document	1-32
X	--- T. PRAZNOVSZKY ET AL.: "De novo chromosome formation in rodent cells" PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES, vol. 88, 1991, WASHINGTON, DC, US, pages 11042-11046, XP002037029 see the whole document	33-50, 75,76
X	--- F. RAYNAL ET AL.: "Complete nucleotide sequence of mouse 18S rRNA gene : comparison with other available homologs" FEBS LETTERS, vol. 167, 1984, AMSTERDAM, NL, pages 263-268, XP002037024 see the whole document, specially figure 2	87-90, 99-101, 103-108, 113
X	--- R.M. TORCZYNSKI ET AL.: "Cloning and sequencing of a human 18S ribosomal RNA gene" DNA, vol. 4, no. 4, 1985, NEW YORK, NY,US, pages 283-291, XP002037025 see the whole document, specially figure 3	87-90, 99-108, 113
X	--- S. CROSS ET AL.: "The structure of a subterminal repeated sequence present on many human chromosomes" NUCLEIC ACIDS RESEARCH, vol. 18, 1990, OXFORD, GB, pages 6649-6657, XP002037026 see the whole document, specially figure 3	115,117

-/--

INTERNATIONAL SEARCH REPORT

Application No

CT/US 97/05911

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO 94 23049 A (THE JOHNS HOPKINS UNIVERSITY) 13 October 1994 see the whole document ---	51-61, 81-84, 96-98, 110,111
A	EP 0 473 253 A (BIOLOGICAL RESEARCH CENTER OF THE HUNGARIAN ACADEMY OF SCIENCES) 4 March 1992 cited in the application see abstract see examples 1,3 ---	66-72, 112
A	WO 95 32297 A (CANCER RESEARCH CAMPAIGN TECHNOLOGY LIMITED) 30 November 1995 see abstract see page 5-6 see example 6 ---	51-65, 73-84, 99,100
A	A. SCHEDL ET AL.: "A method for generation of YAC transgenic mice by pronuclear microinjection" NUCLEIC ACIDS RESEARCH, vol. 21, no. 20, 1993, OXFORD, GB, pages 4783-4787, XP000616418 cited in the application see the whole document ---	51-62
A	WO 94 24300 A (TRANSGENE SA) 27 October 1994 see abstract ---	3,37, 119-122
A	WO 92 17582 A (THE REGENTS OF THE UNIVERSITY OF MICHIGAN) 15 October 1992 see page 11-19 see abstract & US 5 240 840 A cited in the application ---	13-32, 67-72
A	WO 95 29992 A (THE REGENTS OF THE UNIVERSITY OF MICHIGAN) 9 November 1995 see page 12-14 ---	13-32, 67-72
P,X	J. KERESO ET AL.: "De novo chromosome formations by large-scale amplification of the centromeric region of mouse chromosomes" CHROMOSOME RESEARCH, vol. 4, no. 3, 5 June 1996, OXFORD, GB, pages 226-239, XP002037022 see the whole document ---	1-21, 23-50, 75,76, 78, 119-123

-/--

INTERNATIONAL SEARCH REPORT

Application No

PCT/US 97/05911

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>GY. HOLLO ET AL: "Evidence for a megareplicon covering megabases of centromeric chromosome segments" CHROMOSOME RESEARCH, vol. 4, no. 3, 5 June 1996, OXFORD, GB, pages 240-247, XP002037023 see the whole document -----</p>	<p>1-21, 23-32, 35-40</p>

INTERNATIONAL SEARCH REPORT

information on patent family members

Application No

PCT/US 97/05911

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0532050 A	17-03-93	US 5288625 A CA 2078189 A JP 7177881 A	22-02-94 14-03-93 18-07-95
WO 9423049 A	13-10-94	NONE	
EP 0473253 A	04-03-92	CA 2042093 A JP 6121685 A US 5712134 A	10-11-91 06-05-94 27-01-98
WO 9532297 A	30-11-95	AU 2534395 A ZA 9504300 A	18-12-95 24-01-96
WO 9424300 A	27-10-94	FR 2703996 A AU 6571994 A CA 2160697 A EP 0694072 A JP 8508878 T	21-10-94 08-11-94 27-10-94 31-01-96 24-09-96
WO 9217582 A	15-10-92	US 5240840 A	31-08-93
WO 9529992 A	09-11-95	US 5635376 A	03-06-97